

(19) World Intellectual Property  
Organization  
International Bureau



(43) International Publication Date  
16 September 2004 (16.09.2004)

PCT

(10) International Publication Number  
**WO 2004/078169 A1**

(51) International Patent Classification<sup>7</sup>: **A61K 31/185**,  
31/415, 31/433, A61P 19/00, 9/02, 1/04, 27/06

**Timothy, Michael** [US/US]; Pfizer Global Research &  
Development, 2800 Plymouth Road, Ann Arbor, MI 48105  
(US).

(21) International Application Number:  
PCT/IB2004/000553

(74) Agents: **LUMB, J., Trevor** et al.; Pfizer Inc., C/o  
Lawrence, Jackie, Eastern Point Road, Mailstop  
8260-1615, Groton, CT 06340 (US).

(22) International Filing Date: 23 February 2004 (23.02.2004)

(25) Filing Language: English

(81) Designated States (*unless otherwise indicated, for every  
kind of national protection available*): AE, AG, AL, AM,  
AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN,  
CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI,  
GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE,  
KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD,  
MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG,  
PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM,  
TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM,  
ZW.

(26) Publication Language: English

(30) Priority Data:  
60/451,889 4 March 2003 (04.03.2003) US

(71) Applicant (*for all designated States except US*): **PFIZER  
PRODUCTS INC.** [US/US]; Eastern Point Road, Groton,  
CT 06340 (US).

(72) Inventors; and

(84) Designated States (*unless otherwise indicated, for every  
kind of regional protection available*): ARIPO (BW, GH,  
GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),  
Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), Euro-  
pean (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR,  
GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK,  
TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW,  
ML, MR, NE, SN, TD, TG).

(75) Inventors/Applicants (*for US only*): **CONSTAN,  
Alexander, Angelo** [US/US]; Pfizer Global Research &  
Development, Eastern Point Road, Groton, CT 06340  
(US). **KESHARY, Prakash, Raj** [US/US]; Pfizer Global  
Research & Development, Eastern Point Road, Groton,  
CT 06340 (US). **MACLEAN, David, Burton** [US/US];  
479 Wolcott Avenue, Middletown, RI 02842 (US). **PAR-  
ALKAR, Vishwas, Madhav** [US/US]; Pfizer Global  
Research & Development, Eastern Point Road, Groton,  
CT 06340 (US). **ROMAN, Doina, Cosma** [US/US];  
Pfizer Global Research & Development, 2800 Plymouth  
Road, Ann Arbor, MI 48105 (US). **THOMPSON, David,  
Duane** [US/US]; Pfizer Global Research & Development,  
Eastern Point Road, Groton, CT 06340 (US). **WRIGHT,**

Published:

— *with international search report*

*For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.*

(54) Title: USE OF EP2 SELECTIVE RECEPTOR AGONISTS IN MEDICAL TREATMENT

(57) Abstract: The present invention relates to methods of treating pulmonary hypertension, facilitating joint fusion, facilitating tendon and ligament repair, reducing the occurrence of secondary fracture, treating avascular necrosis, facilitating cartilage repair, facilitating bone healing after limb transplantation, facilitating liver regeneration, facilitating wound healing, reducing the occurrence of gastric ulceration, treating hypertension, facilitating the growth of tooth enamel or finger or toe nails, treating glaucoma, treating ocular hypertension, and repairing damage caused by metastatic bone disease using an EP<sub>2</sub> selective receptor agonist.



WO 2004/078169 A1

-1-

USE OF EP<sub>2</sub> SELECTIVE RECEPTOR AGONISTS IN MEDICAL TREATMENT**Field of the Invention**

5           The present invention relates to methods of treating pulmonary hypertension, facilitating joint fusion, facilitating tendon and ligament repair, reducing the occurrence of secondary fracture, treating avascular necrosis, facilitating cartilage repair, facilitating bone healing after limb transplantation, facilitating liver regeneration, facilitating wound healing, reducing the occurrence of gastric ulceration, 10 treating hypertension, facilitating the growth of tooth enamel or finger or toe nails, treating glaucoma, treating ocular hypertension, and repairing damage caused by metastatic bone disease using an EP<sub>2</sub> selective receptor agonist.

**Background of the Invention**

15           Compounds that are prostaglandin receptor ligands are known to be useful to treat various diseases such as osteoporosis. A variety of natural prostaglandins such as PGE, PGD and PDF are associated with skeletal metabolism. PGE<sub>2</sub> has been reported to stimulate bone formation, increase bone mass and bone strength in animal models of osteoporosis when administered systemically or locally. However, 20 there are severe side effects associated with PGE<sub>2</sub> such as diarrhea, gastrointestinal bleeding, decreased food consumption, dehydration, weight loss and decreased physical activity. Accordingly, PGE<sub>2</sub> has not found widespread use in humans because of these side effects. Recently, four different subtypes of PGE<sub>2</sub> receptors have been cloned. The four subtypes have been named EP<sub>1</sub>, EP<sub>2</sub>, EP<sub>3</sub> and EP<sub>4</sub>, and 25 research to better understand the pharmacology of the receptor subtypes is presently being conducted.

          The present invention provides methods of treating pulmonary hypertension, facilitating joint fusion, facilitating tendon and ligament repair, reducing the occurrence of secondary fracture, treating avascular necrosis, facilitating cartilage 30 repair, facilitating bone healing after limb transplantation, facilitating liver regeneration, facilitating wound healing, reducing the occurrence of gastric ulceration, treating hypertension, facilitating the growth of tooth enamel or finger or toe nails, treating glaucoma, treating ocular hypertension, and repairing damage caused by metastatic bone disease using an EP<sub>2</sub> selective receptor agonist. Certain EP<sub>2</sub>


-2-

selective receptor agonists are known in the art. See, for example, U.S. Patent Number 6,498,172.

### Summary of the Invention

The present invention provides methods of treating pulmonary hypertension, facilitating joint fusion, facilitating tendon and ligament repair, reducing the occurrence of secondary fracture, treating avascular necrosis, facilitating cartilage repair, facilitating bone healing after limb transplantation, facilitating liver regeneration, facilitating wound healing, reducing the occurrence of gastric ulceration, treating hypertension, facilitating the growth of tooth enamel or finger or toe nails, treating glaucoma, treating ocular hypertension, and repairing damage caused by metastatic bone disease, the methods comprising administering to a patient in need thereof a therapeutically effective amount of an EP<sub>2</sub> selective receptor agonist.

The present invention also provides such methods wherein the EP<sub>2</sub> selective receptor agonist is a compound of Formula I



Formula I

or a prodrug thereof, or a pharmaceutically acceptable salt thereof, wherein

A is SO<sub>2</sub> or CO;

G is Ar, Ar<sup>1</sup>-V-Ar<sup>2</sup>, Ar-(C<sub>1</sub>-C<sub>6</sub>)alkylene, Ar-CONH-(C<sub>1</sub>-C<sub>6</sub>)alkylene, R<sup>1</sup>R<sup>2</sup>-amino, oxy(C<sub>1</sub>-C<sub>6</sub>)alkylene, amino substituted with Ar, or amino substituted with Ar(C<sub>1</sub>-C<sub>4</sub>)alkylene and R<sup>11</sup>, wherein R<sup>11</sup> is H or (C<sub>1</sub>-C<sub>8</sub>)alkyl, R<sup>1</sup> and R<sup>2</sup> may be taken separately and are independently selected from H and (C<sub>1</sub>-C<sub>8</sub>)alkyl, or R<sup>1</sup> and R<sup>2</sup> are taken together with the nitrogen atom of the amino group to form a five- or six-membered azacycloalkyl, said azacycloalkyl optionally containing an oxygen atom and optionally mono-, di- or tri-substituted independently with up to two oxo, hydroxy, (C<sub>1</sub>-C<sub>4</sub>)alkyl, fluoro or chloro;

B is N or CH;

Q is

-(C<sub>2</sub>-C<sub>6</sub>)alkylene-W-(C<sub>1</sub>-C<sub>3</sub>)alkylene-, said alkylenes each optionally substituted with up to four substituents independently selected from fluoro or (C<sub>1</sub>-C<sub>4</sub>)alkyl,

-3-

-(C<sub>4</sub>-C<sub>8</sub>)alkylene-, said alkylene optionally substituted with up to four substituents independently selected from fluoro or (C<sub>1</sub>-C<sub>4</sub>)alkyl,

-X-(C<sub>1</sub>-C<sub>5</sub>)alkylene-, said alkylene optionally substituted with up to four substituents independently selected from fluoro or (C<sub>1</sub>-C<sub>4</sub>)alkyl,

5       -(C<sub>1</sub>-C<sub>5</sub>)alkylene-X-, said alkylene optionally substituted with up to four substituents independently selected from fluoro or (C<sub>1</sub>-C<sub>4</sub>)alkyl,

-(C<sub>1</sub>-C<sub>3</sub>)alkylene-X-(C<sub>1</sub>-C<sub>3</sub>)alkylene-, said alkylenes each optionally substituted with up to four substituents independently selected from fluoro or (C<sub>1</sub>-C<sub>4</sub>)alkyl,

10       -(C<sub>2</sub>-C<sub>4</sub>)alkylene-W-X-(C<sub>0</sub>-C<sub>3</sub>)alkylene-, said alkylenes each optionally substituted with up to four substituents each independently selected from fluoro or (C<sub>1</sub>-C<sub>4</sub>)alkyl,

-(C<sub>0</sub>-C<sub>4</sub>)alkylene-X-W-(C<sub>1</sub>-C<sub>3</sub>)alkylene-, said alkylenes each optionally substituted with up to four substituents each independently selected from fluoro or (C<sub>1</sub>-C<sub>4</sub>)alkyl,

15       -(C<sub>2</sub>-C<sub>5</sub>)alkylene-W-X-W-(C<sub>1</sub>-C<sub>3</sub>)alkylene-, wherein the two occurrences of W are independent of each other, said alkylenes each optionally substituted with up to four substituents each independently selected from fluoro or (C<sub>1</sub>-C<sub>4</sub>)alkyl,

-(C<sub>1</sub>-C<sub>4</sub>)alkylene-ethenylene-(C<sub>1</sub>-C<sub>4</sub>)alkylene-, said alkylenes and said ethenylene each optionally substituted with up to four substituents each independently selected from fluoro or (C<sub>1</sub>-C<sub>4</sub>)alkyl,

20       -(C<sub>1</sub>-C<sub>4</sub>)alkylene-ethenylene-(C<sub>0</sub>-C<sub>2</sub>)alkylene-X-(C<sub>0</sub>-C<sub>5</sub>)alkylene-, said alkylenes and said ethenylene each optionally substituted with up to four substituents each independently selected from fluoro or (C<sub>1</sub>-C<sub>4</sub>)alkyl,

-(C<sub>1</sub>-C<sub>4</sub>)alkylene-ethenylene-(C<sub>0</sub>-C<sub>2</sub>)alkylene-X-W-(C<sub>1</sub>-C<sub>3</sub>)alkylene-, said alkylenes and said ethenylene optionally each substituted with up to four substituents each independently selected from fluoro or (C<sub>1</sub>-C<sub>4</sub>)alkyl,

25       -(C<sub>1</sub>-C<sub>4</sub>)alkylene-ethynylene-(C<sub>1</sub>-C<sub>4</sub>)alkylene-, said alkylenes and said ethynylene each optionally substituted with up to four substituents each independently selected from fluoro or (C<sub>1</sub>-C<sub>4</sub>)alkyl, or

30       -(C<sub>1</sub>-C<sub>4</sub>)alkylene-ethynylene-X-(C<sub>0</sub>-C<sub>3</sub>)alkylene-, said alkylenes and said ethynylene each optionally substituted with up to four substituents each independently selected from fluoro or (C<sub>1</sub>-C<sub>4</sub>)alkyl;

Z is carboxyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonyl, tetrazolyl, 1,2,4-oxadiazolyl, 5-oxo-1,2,4-oxadiazolyl, 5-oxo-1,2,4-thiadiazolyl, (C<sub>1</sub>-C<sub>4</sub>)alkylsulfonyl/carbamoyl or phenylsulfonyl/carbamoyl;

-4-

K is a bond, (C<sub>1</sub>-C<sub>9</sub>)alkylene, thio(C<sub>1</sub>-C<sub>4</sub>)alkylene, (C<sub>1</sub>-C<sub>4</sub>)alkylenethio(C<sub>1</sub>-C<sub>4</sub>)alkylene, (C<sub>1</sub>-C<sub>4</sub>)alkyleneoxy(C<sub>1</sub>-C<sub>4</sub>)alkylene or oxy(C<sub>1</sub>-C<sub>4</sub>)alkylene, said (C<sub>1</sub>-C<sub>9</sub>)alkylene optionally mono-unsaturated and wherein, when K is not a bond, K is optionally mono-, di- or tri-substituted independently with chloro, fluoro, hydroxy or methyl;

M is -Ar<sup>3</sup>, -Ar<sup>4</sup>-V<sup>1</sup>-Ar<sup>5</sup>, -Ar<sup>4</sup>-S-Ar<sup>5</sup>, -Ar<sup>4</sup>-SO-Ar<sup>5</sup>, -Ar<sup>4</sup>-SO<sub>2</sub>-Ar<sup>5</sup> or -Ar<sup>4</sup>-O-Ar<sup>5</sup>;

Ar is a partially saturated or fully unsaturated five to eight membered ring optionally having one to four heteroatoms selected independently from oxygen, sulfur and nitrogen, or a bicyclic ring consisting of two fused independently partially saturated, fully saturated or fully unsaturated five or six membered rings, taken independently, optionally having one to four heteroatoms selected independently from nitrogen, sulfur and oxygen, or a tricyclic ring consisting of three fused independently partially saturated, fully saturated or fully unsaturated five or six membered rings, optionally having one to four heteroatoms selected independently from nitrogen, sulfur and oxygen, said partially or fully saturated ring, bicyclic ring or tricyclic ring optionally having one or two oxo groups substituted on carbon or one or two oxo groups substituted on sulfur; or Ar is a fully saturated five to seven membered ring having one or two heteroatoms selected independently from oxygen, sulfur and nitrogen;

Ar<sup>1</sup> and Ar<sup>2</sup> are each independently a partially saturated, fully saturated or fully unsaturated five to eight membered ring optionally having one to four heteroatoms selected independently from oxygen, sulfur and nitrogen, or a bicyclic ring consisting of two fused independently partially saturated, fully saturated or fully unsaturated five or six membered rings, taken independently, optionally having one to four heteroatoms selected independently from nitrogen, sulfur and oxygen, or a tricyclic ring consisting of three fused independently partially saturated, fully saturated or fully unsaturated five or six membered rings, optionally having one to four heteroatoms selected independently from nitrogen, sulfur and oxygen, said partially or fully saturated ring, bicyclic ring or tricyclic ring optionally having one or two oxo groups substituted on carbon or one or two oxo groups substituted on sulfur;

said Ar, Ar<sup>1</sup> and Ar<sup>2</sup> moieties are optionally substituted on carbon or nitrogen, on one ring if the moiety is monocyclic, on one or both rings if the moiety is bicyclic, or on one, two or three rings if the moiety is tricyclic, with up to three substituents per

-5-

- moiety independently selected from  $R^3$ ,  $R^4$  and  $R^5$  wherein  $R^3$ ,  $R^4$  and  $R^5$  are independently hydroxy, nitro, halo, carboxy, (C<sub>1</sub>-C<sub>7</sub>)alkoxy, (C<sub>1</sub>-C<sub>4</sub>)alkoxy(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxycarbonyl, (C<sub>1</sub>-C<sub>7</sub>)alkyl, (C<sub>2</sub>-C<sub>7</sub>)alkenyl, (C<sub>2</sub>-C<sub>7</sub>)alkynyl, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl(C<sub>1</sub>-C<sub>4</sub>)alkanoyl, formyl, (C<sub>1</sub>-C<sub>8</sub>)alkanoyl, (C<sub>1</sub>-C<sub>6</sub>)alkanoyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkanoylamino, (C<sub>1</sub>-C<sub>4</sub>)alkoxycarbonylamino, hydroxysulfonyl, aminocarbonylamino or mono-N-, di-N,N-, di-N,N'- or tri-N,N,N'-(C<sub>1</sub>-C<sub>4</sub>)alkyl substituted aminocarbonylamino, sulfonamido, (C<sub>1</sub>-C<sub>4</sub>)alkylsulfonamido, amino, mono-N- or di-N,N-(C<sub>1</sub>-C<sub>4</sub>)alkylamino, carbamoyl, mono-N- or di-N,N-(C<sub>1</sub>-C<sub>4</sub>)alkylcarbamoyl, cyano, thiol, (C<sub>1</sub>-C<sub>6</sub>)alkylthio, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfinyl, (C<sub>1</sub>-C<sub>4</sub>)alkylsulfonyl or mono-N- or di-N,N-(C<sub>1</sub>-C<sub>4</sub>)alkylaminosulfinyl;
- $Ar^3$ ,  $Ar^4$  and  $Ar^5$  are each independently a partially saturated, fully saturated or fully unsaturated five to eight membered ring optionally having one to four heteroatoms selected independently from oxygen, sulfur and nitrogen, or a bicyclic ring consisting of two fused independently partially saturated, fully saturated or fully unsaturated five or six membered rings, taken independently, optionally having one to four heteroatoms selected independently from nitrogen, sulfur and oxygen, or a tricyclic ring consisting of three fused independently partially saturated, fully saturated or fully unsaturated five or six membered rings, optionally having one to four heteroatoms selected independently from nitrogen, sulfur and oxygen, said partially or fully saturated ring, bicyclic ring or tricyclic ring optionally having one or two oxo groups substituted on carbon or one or two oxo groups substituted on sulfur; said  $Ar^3$ ,  $Ar^4$  and  $Ar^5$  moieties are optionally substituted on carbon or nitrogen, on one ring if the moiety is monocyclic, on one or both rings if the moiety is bicyclic, or on one, two or three rings if the moiety is tricyclic, with up to three substituents per moiety independently selected from  $R^{31}$ ,  $R^{41}$  and  $R^{51}$  wherein  $R^{31}$ ,  $R^{41}$  and  $R^{51}$  are independently hydroxy, nitro, halo, carboxy, (C<sub>1</sub>-C<sub>7</sub>)alkoxy, (C<sub>1</sub>-C<sub>4</sub>)alkoxy(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxycarbonyl, (C<sub>1</sub>-C<sub>7</sub>)alkyl, (C<sub>2</sub>-C<sub>7</sub>)alkenyl, (C<sub>2</sub>-C<sub>7</sub>)alkynyl, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl(C<sub>1</sub>-C<sub>4</sub>)alkanoyl, formyl, (C<sub>1</sub>-C<sub>8</sub>)alkanoyl, (C<sub>1</sub>-C<sub>6</sub>)alkanoyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkanoylamino, (C<sub>1</sub>-C<sub>4</sub>)alkoxycarbonylamino, hydroxysulfonyl, aminocarbonylamino or mono-N-, di-N,N-, di-N,N'- or tri-N,N,N'-(C<sub>1</sub>-C<sub>4</sub>)alkyl substituted aminocarbonylamino, sulfonamido, (C<sub>1</sub>-C<sub>4</sub>)alkylsulfonamido, amino, mono-N- or di-N,N-(C<sub>1</sub>-C<sub>4</sub>)alkylamino, carbamoyl, mono-

-6-

N- or di-N,N-(C<sub>1</sub>-C<sub>4</sub>)alkylcarbamoyl, cyano, thiol, (C<sub>1</sub>-C<sub>6</sub>)alkylthio, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfinyl, (C<sub>1</sub>-C<sub>4</sub>)alkylsulfonyl or mono-N- or di-N,N-(C<sub>1</sub>-C<sub>4</sub>)alkylaminosulfinyl;

W is oxy, thio, sulfinio, sulfonyl, aminosulfonyl-, -mono-N-(C<sub>1</sub>-C<sub>4</sub>)alkyleneaminosulfonyl-, sulfonylamino, N-(C<sub>1</sub>-C<sub>4</sub>)alkylenesulfonylamino, carboxamido, N-(C<sub>1</sub>-C<sub>4</sub>)alkylenecarboxamido, carboxamidooxy, N-(C<sub>1</sub>-C<sub>4</sub>)alkylenecarboxamidooxy, carbamoyl, -mono-N-(C<sub>1</sub>-C<sub>4</sub>)alkylenecarbamoyl, carbamoyloxy, or -mono-N-(C<sub>1</sub>-C<sub>4</sub>)alkylenecarbamoyloxy, wherein said W alkyl groups are optionally substituted on carbon with one to three fluorines;

X is a five or six membered aromatic ring optionally having one or two heteroatoms selected independently from oxygen, nitrogen, and sulfur; said ring optionally mono-, di- or tri-substituted independently with halo, (C<sub>1</sub>-C<sub>3</sub>)alkyl, trifluoromethyl, trifluoromethyloxy, difluoromethyloxy, hydroxyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy, or carbamoyl;

R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, R<sup>4</sup>, R<sup>5</sup>, R<sup>11</sup>, R<sup>31</sup>, R<sup>41</sup> and R<sup>51</sup>, when containing an alkyl, alkylene, alkenylene or alkynylene moiety, are optionally mono-, di- or tri-substituted on carbon independently with halo or hydroxy; and

V and V<sup>1</sup> are each independently a bond, thio(C<sub>1</sub>-C<sub>4</sub>)alkylene, (C<sub>1</sub>-C<sub>4</sub>)alkylenethio, (C<sub>1</sub>-C<sub>4</sub>)alkyleneoxy, oxy(C<sub>1</sub>-C<sub>4</sub>)alkylene or (C<sub>1</sub>-C<sub>3</sub>)alkylene optionally mono- or di-substituted independently with hydroxy or fluoro;

with the provisos that:

a. when K is (C<sub>2</sub>-C<sub>4</sub>)alkylene and M is Ar<sup>3</sup> and Ar<sup>3</sup> is cyclopent-1-yl, cyclohex-1-yl, cyclohept-1-yl or cyclooct-1-yl then said (C<sub>5</sub>-C<sub>8</sub>)cycloalkyl substituents are not substituted at the one position with hydroxy; and

b. when K is a bond; G is phenyl, phenylmethyl, substituted phenyl or substituted phenylmethyl; Q is (C<sub>3</sub>-C<sub>8</sub>)alkylene; and M is Ar<sup>3</sup> or Ar<sup>4</sup>-Ar<sup>5</sup>, then A is sulfonyl.

The present invention also provides methods of treating pulmonary hypertension, facilitating joint fusion, facilitating tendon and ligament repair, reducing the occurrence of secondary fracture, treating avascular necrosis, facilitating cartilage repair, facilitating bone healing after limb transplantation, facilitating liver regeneration, facilitating wound healing, reducing the occurrence of gastric ulceration, treating hypertension, facilitating the growth of tooth enamel or finger or toe nails, treating glaucoma, treating ocular hypertension, and repairing damage caused by

-7-

metastatic bone disease, the methods comprising administering to a patient in need thereof a therapeutically effective amount of (3-(((4-tert-butyl-benzyl)-(pyridine-3-sulfonyl)-amino)-methyl)-phenoxy)-acetic acid or a pharmaceutically acceptable salt thereof.

5

### Detailed Description of the Invention

Examples of EP<sub>2</sub> selective receptor agonists are set forth in U.S. Patent Number 6,498,172. A preferred EP<sub>2</sub> selective receptor agonist that can be used in the present methods is a compound of Formula I as defined above.

A preferred group of compounds designated the A Group, comprises those  
 10 compounds having the Formula I as shown above, prodrugs thereof and pharmaceutically acceptable salts of said compounds and said prodrugs, wherein B is N; Z is carboxyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonyl or tetrazolyl; Ar is phenyl, furyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, pyridyl, pyridazinyl, pyrimidinyl, pyrazinyl, 2H-pyrrolyl, 3H-pyrrolyl, pyrrolyl, 2-pyrrolinyl, 3-  
 15 pyrrolinyl, pyrrolidinyl, 1,3-dioxolanyl, 2H-imidazolyl, 2-imidazolinyl, imidazolidinyl, 2-pyrazolinyl, pyrazolidinyl, 1,2,3-oxadiazolyl, 1,2,4-oxadiazolyl, 1,2,5-oxadiazolyl, 1,3,4-oxadiazolyl, 1,2,3-triazolyl, 1,2,4-triazolyl, 2H-pyranyl, 4H-pyranyl, pyridyl, piperidinyl, 1,4-dioxanyl, morpholinyl, 1,4-dithianyl, thiomorpholinyl, piperazinyl, 1,3,5-triazinyl, 1,2,4-triazinyl, azepinyl, oxepinyl, thiepinyl, cyclopentenyl, cyclohexenyl,  
 20 benzo(b)thienyl, benzoxazolyl, benzimidazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, phthalazinyl, quinazolinyl, quinoxalyl, naphthyl, tetralinyl, decalinyl, 2H-1-benzopyranyl and 1,4-benzodioxan; Ar<sup>1</sup>, Ar<sup>2</sup>, Ar<sup>3</sup>, Ar<sup>4</sup> and Ar<sup>5</sup> are each independently cyclopentyl, cyclohexyl, cycloheptyl, cyclooctyl, phenyl, furyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, pyridyl, pyridazinyl,  
 25 pyrimidinyl, pyrazinyl, 2H-pyrrolyl, 3H-pyrrolyl, pyrrolyl, 2-pyrrolinyl, 3-pyrrolinyl, pyrrolidinyl, 1,3-dioxolanyl, 2H-imidazolyl, 2-imidazolinyl, imidazolidinyl, 2-pyrazolinyl, pyrazolidinyl, 1,2,3-oxadiazolyl, 1,2,4-oxadiazolyl, 1,2,5-oxadiazolyl, 1,3,4-oxadiazolyl, 1,2,3-triazolyl, 1,2,4-triazolyl, 2H-pyranyl, 4H-pyranyl, pyridyl, piperidinyl, 1,4-dioxanyl, morpholinyl, 1,4-dithianyl, thiomorpholinyl, piperazinyl, 1,3,5-triazinyl,  
 30 1,2,4-triazinyl, azepinyl, oxepinyl, thiepinyl, 1,2,4-diazepinyl, cyclopentenyl, cyclohexenyl, cycloheptenyl, cyclooctenyl, cyclooctadienyl, indolizinyl, indolyl, isoindolyl, 3H-indolyl, 1H-isoindolyl, indolinyl, cyclopenta(b)pyridinyl, pyrano(3,4-b)pyrrolyl, benzofuryl, isobenzofuryl, benzo(b)thienyl, benzo(c)thienyl, 1H-indazolyl,



-8-

indoxaziny, benzoxazolyl, anthranilyl, benzimidazolyl, benzthiazolyl, puriny, 4H-quinoliziny, quinoliny, isoquinoliny, cinnoliny, phthalaziny, quinazoliny, quinoxaliny, 1,8-naphthyridiny, pteridiny, indenyl, isoindenyl, naphthyl, tetraliny, decaliny, 2H-1-benzopyrany, 1,4-benzodioxan, pyrido(3,4-b)-pyridiny, pyrido(3,2-b)-pyridiny, 5 pyrido(4,3-b)-pyridiny, 2H-1,3-benzoxaziny, 2H-1,4-benzoxaziny, 1H-2,3-benzoxaziny, 4H-3,1-benzoxaziny, 2H-1,2-benzoxaziny and 4H-1,4-benzoxaziny; and X is tetrahydrofurany, phenyl, thiazolyl, thienyl, pyridyl, pyrrazolyl, furanyl or pyrimidyl, wherein X is optionally mono-, di- or tri-substituted independently with chloro, fluoro, methoxy, difluoromethoxy, trifluoromethoxy, trifluoromethyl or methyl; 10 and wherein each of said Ar, Ar<sup>1</sup> and Ar<sup>2</sup> groups are optionally substituted on carbon or nitrogen with up to three substituents independently selected from R<sup>3</sup>, R<sup>4</sup> and R<sup>5</sup>; each of said Ar, Ar<sup>1</sup> and Ar<sup>2</sup> groups are optionally substituted independently on carbon or sulfur with one or two oxo groups; each of said Ar<sup>3</sup>, Ar<sup>4</sup> and Ar<sup>5</sup> groups are optionally substituted on carbon or nitrogen independently with up to three R<sup>31</sup>, R<sup>41</sup> 15 and R<sup>51</sup> groups and each of said Ar<sup>3</sup>, Ar<sup>4</sup> and Ar<sup>5</sup> groups are optionally substituted independently on carbon or sulfur with one or two oxo groups.

A group of compounds within the A Group, designated the B Group, comprises those compounds, prodrugs thereof and pharmaceutically acceptable salts of said compounds and said prodrugs, wherein A is CO; G is oxy(C<sub>1</sub>-C<sub>6</sub>)alkylene; Q is 20 -(C<sub>2</sub>-C<sub>6</sub>)alkylene-O-(C<sub>1</sub>-C<sub>3</sub>)alkylene-, -(C<sub>4</sub>-C<sub>8</sub>)alkylene-, said -(C<sub>4</sub>-C<sub>8</sub>)alkylene- optionally substituted with up to four substituents independently selected from fluoro or (C<sub>1</sub>-C<sub>4</sub>)alkyl, -X-(C<sub>2</sub>-C<sub>5</sub>)alkylene-, -(C<sub>1</sub>-C<sub>5</sub>)alkylene-X-, 25 -(C<sub>1</sub>-C<sub>3</sub>)alkylene-X-(C<sub>1</sub>-C<sub>3</sub>)alkylene-, -(C<sub>2</sub>-C<sub>4</sub>)alkylene-O-X-(C<sub>0</sub>-C<sub>3</sub>)alkylene-, or -(C<sub>0</sub>-C<sub>4</sub>)alkylene-X-O-(C<sub>1</sub>-C<sub>3</sub>)alkylene-; and X is phenyl, thienyl, furanyl or thiazolyl, wherein X is optionally mono-, di- or tri-substituted with chloro, fluoro, methoxy, difluoromethoxy, trifluoromethoxy, trifluoromethyl or methyl.

30 Another group of compounds which is preferred within the A Group, designated the C Group, comprises those compounds, prodrugs thereof and pharmaceutically acceptable salts of said compounds and said prodrugs, wherein A is CO; G is Ar; Q is

-9-

-(C<sub>2</sub>-C<sub>6</sub>)alkylene-O-(C<sub>1</sub>-C<sub>3</sub>)alkylene-,  
 -(C<sub>4</sub>-C<sub>8</sub>)alkylene-, said -(C<sub>4</sub>-C<sub>8</sub>)alkylene- optionally substituted with up to four  
 substituents independently selected from fluoro or (C<sub>1</sub>-C<sub>4</sub>)alkyl,  
 -X-(C<sub>2</sub>-C<sub>5</sub>)alkylene-,  
 5       -(C<sub>1</sub>-C<sub>5</sub>)alkylene-X-,  
 -(C<sub>1</sub>-C<sub>3</sub>)alkylene-X-(C<sub>1</sub>-C<sub>3</sub>)alkylene-,  
 -(C<sub>2</sub>-C<sub>4</sub>)alkylene-O-X-(C<sub>0</sub>-C<sub>3</sub>)alkylene-, or  
 -(C<sub>0</sub>-C<sub>4</sub>)alkylene-X-O-(C<sub>1</sub>-C<sub>3</sub>)alkylene-; and X is phenyl, thienyl, furanyl or  
 thiazolyl, wherein X is optionally mono-, di- or tri-substituted with chloro, fluoro,  
 10       methoxy, difluoromethoxy, trifluoromethoxy, trifluoromethyl or methyl.

Another group of compounds which is preferred within the A Group,  
 designated the D Group, comprises those compounds, prodrugs thereof and  
 pharmaceutically acceptable salts of said compounds and said prodrugs, wherein A  
 is CO; G is R<sup>1</sup>R<sup>2</sup>-amino or amino substituted with Ar, or amino substituted with Ar(C<sub>1</sub>-  
 15       C<sub>4</sub>)alkylene and R<sup>11</sup>, wherein R<sup>11</sup> is H; Q is

-(C<sub>2</sub>-C<sub>6</sub>)alkylene-O-(C<sub>1</sub>-C<sub>3</sub>)alkylene-,  
 -(C<sub>4</sub>-C<sub>8</sub>)alkylene-, said -(C<sub>4</sub>-C<sub>8</sub>)alkylene- optionally substituted with up to four  
 substituents independently selected from fluoro or (C<sub>1</sub>-C<sub>4</sub>)alkyl,  
 -X-(C<sub>2</sub>-C<sub>5</sub>)alkylene-,  
 20       -(C<sub>1</sub>-C<sub>5</sub>)alkylene-X-,  
 -(C<sub>1</sub>-C<sub>3</sub>)alkylene-X-(C<sub>1</sub>-C<sub>3</sub>)alkylene-,  
 -(C<sub>2</sub>-C<sub>4</sub>)alkylene-O-X-(C<sub>0</sub>-C<sub>3</sub>)alkylene-, or  
 -(C<sub>0</sub>-C<sub>4</sub>)alkylene-X-O-(C<sub>1</sub>-C<sub>3</sub>)alkylene-; and X is phenyl, thienyl, furanyl or  
 thiazolyl, wherein X is optionally mono-, di- or tri-substituted with chloro, fluoro,  
 25       methoxy, difluoromethoxy, trifluoromethoxy, trifluoromethyl or methyl; and

wherein R<sup>1</sup> and R<sup>2</sup> may be taken separately and are independently selected  
 from H and (C<sub>1</sub>-C<sub>6</sub>)alkyl, or R<sup>1</sup> and R<sup>2</sup> are taken together to form a five- or six-  
 membered azacycloalkyl, said azacycloalkyl optionally containing an oxygen atom.

Another group of compounds which is preferred within the G Group,  
 30       designated the E Group, comprises those compounds, prodrugs thereof and  
 pharmaceutically acceptable salts of said compounds and said prodrugs, wherein A  
 is SO<sub>2</sub>; G is R<sup>1</sup>R<sup>2</sup>-amino, or amino substituted with Ar and R<sup>11</sup>; Q is

-(C<sub>2</sub>-C<sub>6</sub>)alkylene-O-(C<sub>1</sub>-C<sub>3</sub>)alkylene-,

-10-

-(C<sub>4</sub>-C<sub>8</sub>)alkylene-, said -(C<sub>4</sub>-C<sub>8</sub>)alkylene- optionally substituted with up to four substituents independently selected from fluoro or (C<sub>1</sub>-C<sub>4</sub>)alkyl,

-X-(C<sub>2</sub>-C<sub>5</sub>)alkylene-,

-(C<sub>1</sub>-C<sub>5</sub>)alkylene-X-,

5       -(C<sub>1</sub>-C<sub>3</sub>)alkylene-X-(C<sub>1</sub>-C<sub>3</sub>)alkylene-,

-(C<sub>2</sub>-C<sub>4</sub>)alkylene-O-X-(C<sub>0</sub>-C<sub>3</sub>)alkylene-, or

-(C<sub>0</sub>-C<sub>4</sub>)alkylene-X-O-(C<sub>1</sub>-C<sub>3</sub>)alkylene-; and X is phenyl, thienyl, furanyl or thiazolyl, wherein X is optionally mono-, di- or tri-substituted with chloro, fluoro, methoxy, difluoromethoxy, trifluoromethoxy, trifluoromethyl or methyl; and

10       wherein R<sup>1</sup> and R<sup>2</sup> may be taken separately and are independently selected from H and (C<sub>1</sub>-C<sub>8</sub>)alkyl, or R<sup>1</sup> and R<sup>2</sup> are taken together to form a five- or six-membered azacycloalkyl, said azacycloalkyl optionally containing an oxygen atom.

Another group of compounds which is preferred within the A Group,

designated the F Group, comprises those compounds, prodrugs thereof and

15       pharmaceutically acceptable salts of said compounds and said prodrugs, wherein A is SO<sub>2</sub>; G is Ar, Ar(C<sub>1</sub>-C<sub>2</sub>)alkylene or Ar<sup>1</sup>-V-Ar<sup>2</sup>; Q is

-(C<sub>2</sub>-C<sub>6</sub>)alkylene-O-(C<sub>1</sub>-C<sub>3</sub>)alkylene-,

-(C<sub>4</sub>-C<sub>8</sub>)alkylene-, said -(C<sub>4</sub>-C<sub>8</sub>)alkylene- optionally substituted with up to four substituents independently selected from fluoro or (C<sub>1</sub>-C<sub>4</sub>)alkyl,

20       -X-(C<sub>2</sub>-C<sub>5</sub>)alkylene-,

-(C<sub>1</sub>-C<sub>5</sub>)alkylene-X-,

-(C<sub>1</sub>-C<sub>3</sub>)alkylene-X-(C<sub>1</sub>-C<sub>3</sub>)alkylene-,

-(C<sub>2</sub>-C<sub>4</sub>)alkylene-O-X-(C<sub>0</sub>-C<sub>3</sub>)alkylene-, or

25       -(C<sub>0</sub>-C<sub>4</sub>)alkylene-X-O-(C<sub>1</sub>-C<sub>3</sub>)alkylene-; and X is phenyl, pyrimidyl, pyridyl,



thienyl, tetrahydrofuranyl, furanyl or thiazolyl, wherein X is optionally mono-, di- or tri-substituted with chloro, fluoro, methoxy, difluoromethoxy, trifluoromethoxy, trifluoromethyl or methyl.

A particularly preferred group of compounds within the F Group, designated the FA Group, comprises those compounds, prodrugs thereof and pharmaceutically

30       acceptable salts of said compounds and said prodrugs, wherein G is Ar or Ar-(C<sub>1</sub>-C<sub>2</sub>)-alkylene; Ar is phenyl, furyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, pyridyl, pyridazinyl, pyrimidinyl, pyrazinyl, isothiazolyl, 1,2,3-triazolyl, 1,2,4-triazolyl or 1,3,4-thiadiazolyl wherein each of said Ar groups is optionally substituted

-11-

on carbon or nitrogen with  $R^1$ ,  $R^2$  or  $R^3$ ;  $Ar^4$  is cyclopentyl, cyclohexyl, cycloheptyl, cyclooctyl, phenyl, furyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, pyridyl, pyridazinyl, pyrimidinyl, pyrazinyl, pyrrolidinyl, 1,2,3-triazolyl, 1,2,4-triazolyl, pyranyl, thiomorpholinyl, piperazinyl, 1,3,5-triazinyl, 1,2,4-triazinyl, 1,2,3-triazinyl, azepinyl, oxepinyl or thiepinyl wherein each of said  $Ar^4$  groups is optionally mono- di- or tri-substituted on carbon or nitrogen with  $R^{31}$ ,  $R^{41}$  or  $R^{51}$ ;  $Ar^5$  is cyclopentyl, cyclohexyl, cycloheptyl, cyclooctyl, phenyl, furyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, pyridyl, pyridazinyl, pyrimidinyl, pyrazinyl, pyrrolidinyl, 1,2,3-triazolyl, 1,2,4-triazolyl, pyranyl, 1,4-dioxanyl, thiomorpholinyl, piperazinyl, 1,3,5-triazinyl, 1,2,4-triazinyl, 1,2,3-triazinyl, azepinyl, oxepinyl or thiepinyl wherein each of said  $Ar^5$  groups is optionally mono- di- or tri-substituted on carbon or nitrogen with  $R^{31}$ ,  $R^{41}$  or  $R^{51}$ ; Q is  $-(C_5-C_7)$ -alkylene-,  $-(C_1-C_2)$ -alkylene-X- $(C_1-C_2)$ -alkylene-,  $-(C_1-C_2)$ -X-O- $(C_1-C_2)$ -alkylene-,  $-(C_2-C_4)$ -alkylene-thienyl-,  $-(C_2-C_4)$ -alkylene-furanyl- or  $-(C_2-C_4)$ -alkylene-thiazolyl-; X is phenyl, pyridyl, pyrimidyl or thienyl; and said X groups are optionally mono-, di- or tri- substituted with chloro, fluoro, methoxy, difluoromethoxy, trifluoromethoxy, trifluoromethyl or methyl; said  $-(C_2-C_4)$ -alkylene-furanyl- and  $-(C_2-C_4)$ -alkylene-thienyl- having a 2,5-substitution pattern, e.g.,



A preferred group of compounds within the FA Group, designated the FB Group, comprises those compounds, prodrugs thereof and pharmaceutically acceptable salts of said compounds and said prodrugs, wherein K is methylene, M is  $Ar^4-Ar^5$ ,  $Ar^4-O-Ar^5$  or  $Ar^4-S-Ar^5$  and Ar is phenyl, pyridyl, pyrazolyl, imidazolyl, pyrimidyl, thienyl or thiazolyl, wherein Ar is optionally mono-, di- or tri-substituted on carbon or nitrogen with  $R^3$ ,  $R^4$  or  $R^5$ .

A preferred group of compounds within the FB Group, designated the FC Group, comprises those compounds, prodrugs thereof and pharmaceutically acceptable salts of said compounds and said prodrugs, wherein M is  $Ar^4-Ar^5$ ; Ar is phenyl, pyridyl or imidazolyl;  $Ar^4$  is phenyl, furanyl or pyridyl; and  $Ar^5$  is cyclopentyl, cyclohexyl, cycloheptyl, phenyl, pyridyl, imidazolyl, pyrimidyl, thienyl, pyridazinyl, pyrazinyl, imidazolyl, pyrazolyl or thiazolyl, wherein Ar,  $Ar^4$  and  $Ar^5$  are optionally

-12-

mono, -di- or tri-substituted on carbon or nitrogen independently with chloro, fluoro, methyl, methoxy, difluoromethoxy, trifluoromethyl or trifluoromethoxy.

An especially preferred group of compounds within the FC Group, designated the FD Group, comprises those compounds, prodrugs thereof and pharmaceutically acceptable salts of said compounds and said prodrugs, wherein Q is -(C<sub>5</sub>-C<sub>7</sub>)alkylene-.

Another especially preferred group of compounds within the FC Group, designated the FE Group, comprises those compounds, prodrugs thereof and pharmaceutically acceptable salts of said compounds and said prodrugs, wherein Q is CH<sub>2</sub>-X-CH<sub>2</sub>- and X is metaphenylene optionally mono- or di- substituted with chloro, fluoro, methoxy, difluoromethoxy, trifluoromethoxy, trifluoromethyl or methyl.

A preferred group of compounds within the FE Group are those compounds, and pharmaceutically acceptable salts and prodrugs thereof, selected from (3-(((pyridine-3-sulfonyl)-(4-pyrimidin-5-yl-benzyl)-amino)-methyl)-phenyl)-acetic acid; (3-(((5-phenyl-furan-2-ylmethyl)-(pyridine-3-sulfonyl)-amino)-methyl)-phenyl)-acetic acid; (3-(((pyridine-3-sulfonyl)-(4-pyrimidin-2-yl-benzyl)-amino)-methyl)-phenyl)-acetic acid; (3-(((pyridine-3-sulfonyl)-(4-thiazol-2-yl-benzyl)-amino)-methyl)-phenyl)-acetic acid; and (3-(((4-pyrazin-2-yl-benzyl)-(pyridine-3-sulfonyl)-amino)-methyl)-phenyl)-acetic acid.

An especially preferred compound within the FE Group is the compound wherein Ar is pyrid-3-yl; Z is carboxy; M is Ar<sup>4</sup>-Ar<sup>5</sup> wherein Ar<sup>4</sup> is a furanyl ring and Ar<sup>5</sup> is phenyl wherein said phenyl moiety is substituted at the 5-position of said furanyl ring; and Q is -CH<sub>2</sub>-X-CH<sub>2</sub>- wherein X is metaphenylene.

Another especially preferred compound within the FE Group is the compound wherein Ar is pyrid-3-yl; Z is carboxy; M is Ar<sup>4</sup>-Ar<sup>5</sup> wherein Ar<sup>4</sup> is phenyl and Ar<sup>5</sup> is pyrimid-2-yl and said pyrimid-2-yl moiety is substituted at the 4-position of said phenyl ring; and Q is -CH<sub>2</sub>-X-CH<sub>2</sub>- wherein X is metaphenylene.

Yet another especially preferred compound within the FE Group is the compound wherein Ar is pyrid-3-yl; Z is carboxy; M is Ar<sup>4</sup>-Ar<sup>5</sup> wherein Ar<sup>4</sup> is phenyl and Ar<sup>5</sup> is thiazol-2-yl and said thiazol-2-yl moiety is substituted at the 4-position of said phenyl ring; and Q is -CH<sub>2</sub>-X-CH<sub>2</sub>- wherein X is metaphenylene.

Yet another especially preferred compound within the FE Group is the compound wherein Ar is pyrid-3-yl; Z is carboxy; M is Ar<sup>4</sup>-Ar<sup>5</sup> wherein Ar<sup>4</sup> is phenyl

-13-

and Ar<sup>5</sup> is pyrimid-5-yl and said pyrimid-5-yl moiety is substituted at the 4-position of said phenyl ring; and Q is -CH<sub>2</sub>-X-CH<sub>2</sub>- wherein X is metaphenylene.

Yet another especially preferred compound within the FE Group is the compound wherein Ar is pyrid-3-yl; Z is carboxy; M is Ar<sup>4</sup>-Ar<sup>5</sup> wherein Ar<sup>4</sup> is phenyl  
5 and Ar<sup>5</sup> is pyrazin-2-yl and said pyrazin-2-yl is substituted at the 4-position of said phenyl ring; and Q is -CH<sub>2</sub>-X-CH<sub>2</sub>- wherein X is metaphenylene.

A preferred group of compounds within the FC Group, designated the G Group, comprises those compounds, prodrugs thereof and pharmaceutically acceptable salts of said compounds and said prodrugs, wherein Q is -(C<sub>2</sub>-C<sub>4</sub>)-  
10 alkylene-thienyl-, -(C<sub>2</sub>-C<sub>4</sub>)-alkylene-furanyl- or -(C<sub>2</sub>-C<sub>4</sub>)-alkylene-thiazolyl-.

An especially preferred compound within the G Group is 5-(3-((pyridine-3-sulfonyl)-(4-thiazol-2-yl-benzyl)-amino)-propyl)-thiophene-2-carboxylic acid.

An especially preferred compound within the G Group is the compound, prodrugs thereof and pharmaceutically acceptable salts of said compounds and said  
15 prodrugs, wherein Q is n-propylenyl; X is thienyl; Z is carboxy; Ar is 3-pyridyl; Ar<sup>4</sup> is phenyl; and Ar<sup>5</sup> is 2-thiazolyl; said 2-thiazolyl being substituted at the 4-position of said phenyl.

Another especially preferred group of compounds within the FC Group, designated the H Group, comprises those compounds, prodrugs thereof and  
20 pharmaceutically acceptable salts of said compounds and said prodrugs, wherein Q is -CH<sub>2</sub>-X-O-CH<sub>2</sub>-; Ar<sup>4</sup> is phenyl or pyridyl; said phenyl and pyridyl are optionally substituted with chloro, fluoro, methoxy, difluoromethoxy, trifluoromethoxy, trifluoromethyl and methyl; and X is metaphenylene.

A preferred group of compounds within the H Group are (3-(((4-cyclohexyl-  
25 benzyl)-(pyridine-3-sulfonyl)-amino)-methyl)-phenoxy)-acetic acid; (3-(((pyridine-3-sulfonyl)-(4-pyridin-2-yl-benzyl)-amino)-methyl)-phenoxy)-acetic acid; (3-(((pyridine-3-sulfonyl)-(4-pyridin-3-yl-benzyl)-amino)-methyl)-phenoxy)-acetic acid; (3-(((pyridine-3-sulfonyl)-(4-pyridin-4-yl-benzyl)-amino)-methyl)-phenoxy)-acetic acid; and (3-  
(((pyridine-3-sulfonyl)-(4-thiazol-2-yl-benzyl)-amino)-methyl)-phenoxy)-acetic acid.

30 An especially preferred compound within the H Group is the compound, prodrugs thereof and pharmaceutically acceptable salts of said compounds and said prodrugs, wherein Ar is pyrid-3-yl; Z is carboxy; Ar<sup>4</sup> is phenyl; Ar<sup>5</sup> is cyclohexyl; and said cyclohexyl moiety is substituted at the 4-position of said phenyl ring.

Another especially preferred compound within the H Group is the compound wherein Ar is pyrid-3-yl; Z is carboxy; Ar<sup>4</sup> is phenyl; Ar<sup>5</sup> is thiazol-2-yl; and said thiazol-2-yl moiety is substituted at the 4-position of said phenyl ring.

Yet another especially preferred compound within the H Group is the  
 5 compound wherein Ar is pyrid-3-yl; Z is carboxy; Ar<sup>4</sup> is phenyl; Ar<sup>5</sup> is 2-pyridyl; and said 2-pyridyl moiety is substituted at the 4-position of said phenyl ring.

Yet another especially preferred compound within the H Group is the compound wherein Ar is pyrid-3-yl; Z is carboxy; Ar<sup>4</sup> is phenyl; Ar<sup>5</sup> is 3-pyridyl; and said 3-pyridyl moiety is substituted at the 4-position of said phenyl ring.

10 Yet another especially preferred compound within the H Group is the compound wherein Ar is pyrid-3-yl; Z is carboxy; Ar<sup>4</sup> is phenyl; Ar<sup>5</sup> is 4-pyridyl; and said 4-pyridyl moiety is substituted at the 4-position of said phenyl ring.

A preferred group of compounds within the FA Group, designated the I Group, comprises those compounds, prodrugs thereof and pharmaceutically acceptable salts  
 15 of said compounds and said prodrugs, wherein K is methylene, G is Ar; Ar is phenyl, pyridazinyl, pyrazolyl, pyrazinyl, pyridyl, imidazolyl, pyrimidyl, thienyl or thiazolyl, Ar is optionally mono-, di- or tri-substituted with R<sup>3</sup>, R<sup>4</sup> or R<sup>5</sup>, and M is Ar<sup>3</sup>, wherein said Ar<sup>3</sup> is cyclopentyl, cyclohexyl, phenyl, thienyl, pyridazinyl, pyrimidinyl, pyrazinyl, indolyl, benzofuryl, benzo(b)thienyl, benzoxazolyl, benzthiazolyl, quinoliny, isoquinoliny,  
 20 naphthyl, tetralinyl, 2H-1-benzopyranyl or 1,4-benzodioxan and is optionally mono-, di- or tri-substituted with R<sup>31</sup>, chloro, fluoro, methyl, methoxy, difluoromethoxy, trifluoromethyl or trifluoromethoxy.

An especially preferred group of compounds within the I Group are (3-(((2,3-dihydro-benzo[1,4]dioxin-6-ylmethyl)-(pyridine-3-sulfonyl)-amino)-methyl)-phenyl)-  
 25 acetic acid; and (3-((benzofuran-2-ylmethyl)-(pyridine-3-sulfonyl)-amino)-methyl)-phenyl)-acetic acid.

An especially preferred compound within the I Group is the compound, prodrugs thereof and pharmaceutically acceptable salts of said compound and prodrugs, wherein Ar is pyrid-3-yl; Z is carboxy; M is 6-(1,4-benzodioxan); and Q is -  
 30 CH<sub>2</sub>-X-CH<sub>2</sub>- wherein X is metaphenylene.

Another especially preferred compound within the I Group is the compound wherein Ar is pyrid-3-yl; Z is carboxy; M is 2-benzofuryl; and Q is -CH<sub>2</sub>-X-CH<sub>2</sub>- wherein X is metaphenylene.

Another especially preferred group of compounds within the I Group, designated the J Group, comprises those compounds, prodrugs thereof and pharmaceutically acceptable salts of said compounds and said prodrugs, wherein Ar is phenyl, pyridyl or imidazolyl, said phenyl, pyridyl and imidazolyl optionally substituted independently with chloro, fluoro, methyl, methoxy, difluoromethoxy, trifluoromethyl or trifluoromethoxy; Ar<sup>3</sup> is phenyl substituted with R<sup>31</sup>, wherein R<sup>31</sup> is (C<sub>1</sub>-C<sub>7</sub>)alkyl, mono-N- or di-N, N-(C<sub>1</sub>-C<sub>4</sub>)alkylamine, or (C<sub>1</sub>-C<sub>5</sub>)alkoxy, said (C<sub>1</sub>-C<sub>7</sub>)alkyl or (C<sub>1</sub>-C<sub>5</sub>)alkoxy optionally mono-, di- or tri-substituted independently with hydroxy or fluoro; and Ar<sup>3</sup> is further optionally mono- or di-substituted with chloro, fluoro, methyl, methoxy, difluoromethoxy, trifluoromethoxy or trifluoromethyl.

A preferred group of compounds within the J Group, designated the K Group, comprises those compounds, prodrugs thereof and pharmaceutically acceptable salts of said compounds and said prodrugs, wherein Q is -(C<sub>5</sub>-C<sub>7</sub>)alkylene-.

Another preferred group of compounds within the J Group, designated the L Group, comprises those compounds, prodrugs thereof and pharmaceutically acceptable salts of said compounds and said prodrugs, wherein Q is -CH<sub>2</sub>-X-CH<sub>2</sub>- and X is phenyl optionally mono-, di- or tri- substituted with chloro, fluoro, methoxy, difluoromethoxy, trifluoromethoxy, trifluoromethyl or methyl.

An especially preferred group of compounds within the L Group are (3-(((4-butyl-benzyl)-(pyridine-3-sulfonyl)-amino)-methyl)-phenyl)-acetic acid; (3-(((benzenesulfonyl-(4-butyl-benzyl)-amino)-methyl)-phenyl)-acetic acid; (3-(((4-butyl-benzyl)-(1-methyl-1H-imidazole-4-sulfonyl)-amino)-methyl)-phenyl)-acetic acid; and (3-(((4-dimethylamino-benzyl)-(pyridine-3-sulfonyl)-amino)-methyl)-phenyl)-acetic acid.

An especially preferred compound within the L Group is the compound, prodrugs thereof and pharmaceutically acceptable salts of said compounds and said prodrugs, wherein Ar is pyrid-3-yl; Z is carboxy; M is phenyl substituted at the 4-position with n-butyl; and Q is -CH<sub>2</sub>-X-CH<sub>2</sub>- wherein X is metaphenylene.

Another especially preferred compound within the L Group is the compound, prodrugs thereof and pharmaceutically acceptable salts of said compounds and said prodrugs, wherein Ar is phenyl; Z is carboxy; M is phenyl substituted at the 4-position with n-butyl; and Q is -CH<sub>2</sub>-X-CH<sub>2</sub>- wherein X is metaphenylene.



-16-

Yet another especially preferred compound within the L Group is the compound, prodrugs thereof and pharmaceutically acceptable salts of said compounds and said prodrugs, wherein Ar is 4-(1-methyl-imidazolyl); Z is carboxy; M is phenyl substituted at the 4-position with n-butyl; and Q is -CH<sub>2</sub>-X-CH<sub>2</sub>- wherein X is metaphenylene.

Yet another especially preferred compound within the L Group is the compound, prodrugs thereof and pharmaceutically acceptable salts of said compounds and said prodrugs, wherein Ar is pyrid-3-yl; Z is carboxy; M is phenyl substituted at the 4-position with dimethylamino; and Q is -CH<sub>2</sub>-X-CH<sub>2</sub>- wherein X is metaphenylene.

Another preferred group of compounds within the J Group comprises those compounds, prodrugs thereof and pharmaceutically acceptable salts of said compounds and said prodrugs, wherein Q is -(C<sub>2</sub>-C<sub>4</sub>)alkylene-thienyl, -(C<sub>2</sub>-C<sub>4</sub>)alkylene-furanyl or -(C<sub>2</sub>-C<sub>4</sub>)alkylene-thiazolyl.

A preferred group of compounds within the J Group, designated the M Group, comprises those compounds, prodrugs thereof and pharmaceutically acceptable salts of said compounds and said prodrugs, wherein Q is -(C<sub>1</sub>-C<sub>2</sub>)-X-O-(C<sub>1</sub>-C<sub>2</sub>)alkylene- and X is metaphenylene, said X being optionally mono-, di- or tri-substituted with chloro, fluoro, methoxy, difluoromethoxy, trifluoromethoxy, trifluoromethyl or methyl.

An especially preferred group of compounds within the M Group are (3-(((4-dimethylamino-benzyl)-(pyridine-3-sulfonyl)-amino)-methyl)-phenoxy)-acetic acid and (3-(((4-tert-butyl-benzyl)-(pyridine-3-sulfonyl)-amino)-methyl)-phenoxy)-acetic acid.

An especially preferred compound within the M Group is the compound, prodrugs thereof and pharmaceutically acceptable salts of said compounds and said prodrugs, wherein Ar is pyrid-3-yl; Z is carboxy; M is phenyl substituted at the 4-position with dimethylamino; and Q is -CH<sub>2</sub>-X-O-CH<sub>2</sub>- wherein X is metaphenylene.

Another especially preferred compound within the M Group is the compound, prodrugs thereof and pharmaceutically acceptable salts of said compounds and said prodrugs, wherein Ar is pyrid-3-yl; Z is carboxy; M is phenyl substituted at the 4-position with *tert*-butyl; and Q is -CH<sub>2</sub>-X-O-CH<sub>2</sub>- wherein X is metaphenylene.

Another preferred group of compounds within the FA Group, designated the N Group, comprises those compounds, prodrugs thereof and pharmaceutically acceptable salts of said compounds and said prodrugs, wherein G is Ar; K is (C<sub>2</sub>-C<sub>4</sub>)

alkylene or n-propenylene; Ar is phenyl, pyrazolyl, pyridazinyl, pyrazinyl, pyridyl, imidazolyl, pyrimidyl, thienyl or thiazolyl, wherein Ar is optionally mono-, di- or tri-substituted with R<sup>3</sup>, R<sup>4</sup> or R<sup>5</sup>; and M is Ar<sup>3</sup>, optionally mono-, di- or tri-substituted with chloro, fluoro, methyl, methoxy, difluoromethoxy, trifluoromethoxy and trifluoromethyl.

5 An especially preferred compound within the N Group is *trans*-(3-(((3-(3,5-dichloro-phenyl)-allyl)-(pyridine-3-sulfonyl)-amino)-methyl)-phenyl)-acetic acid.

An especially preferred compound within the N Group is the compound, prodrugs thereof and pharmaceutically acceptable salts of said compounds and said prodrugs, wherein K is *trans*-n-propenylene, said M group being attached to the 1-  
10 position of the n-propenylene and said N atom being attached to the 3-position of the n-propenylene; Ar is pyrid-3-yl; M is phenyl 3,5-disubstituted with chloro; Z is carboxy; and Q is CH<sub>2</sub>-X-CH<sub>2</sub>- wherein X is metaphenylene.

A preferred group of compounds within the N Group, designated the O Group, comprises those compounds, prodrugs thereof and pharmaceutically acceptable salts  
15 of said compounds and said prodrugs, wherein Ar<sup>3</sup> is phenyl optionally substituted with chloro, fluoro, methyl, methoxy, difluoromethoxy, trifluoromethoxy or trifluoromethyl.

A preferred group of compounds within the O Group, designated the P Group, comprises those compounds, prodrugs thereof and pharmaceutically acceptable salts  
20 of said compounds and said prodrugs, wherein Q is -(C<sub>5</sub>-C<sub>7</sub>)alkylene-.

Another group of compounds within the O Group, designated the Q Group, comprises those compounds, prodrugs thereof and pharmaceutically acceptable salts of said compounds and said prodrugs, wherein Q is -CH<sub>2</sub>-X-CH<sub>2</sub>- and X is metaphenylene.

25 Yet another group of compounds within the O Group, designated the R Group, comprises those compounds, prodrugs thereof and pharmaceutically acceptable salts of said compounds and said prodrugs, wherein Q is -(C<sub>2</sub>-C<sub>4</sub>)alkylene-X- and X is furanyl, thienyl or thiazolyl.

Yet another preferred group of compounds within the O Group, designated  
30 the S Group, comprises those compounds, prodrugs thereof and pharmaceutically acceptable salts of said compounds and said prodrugs, wherein Q is -(C<sub>1</sub>-C<sub>2</sub>)-X-O-(C<sub>1</sub>-C<sub>2</sub>)alkylene- and X is metaphenylene.

Another preferred group of compounds within the FA Group, designated the T Group, comprises those compounds, prodrugs thereof and pharmaceutically acceptable salts of said compounds and said prodrugs, wherein G is Ar; K is thioethylene or oxyethylene, Ar is phenyl, pyrazolyl, pyridazinyl, pyrazinyl, pyridyl, imidazolyl, pyrimidyl, thienyl or thiazolyl, wherein Ar is optionally substituted with up to three R<sup>3</sup>, R<sup>4</sup> or R<sup>5</sup>; and M is Ar<sup>3</sup>, optionally mono-, di- or tri-substituted with chloro, fluoro, methyl, difluoromethoxy, trifluoromethoxy or trifluoromethyl.

A preferred group of compounds within the T Group, designated the U Group, comprises those compounds, prodrugs thereof and pharmaceutically acceptable salts of said compounds and said prodrugs, wherein Ar<sup>3</sup> is phenyl.

A preferred group of compounds within the U Group, designated the V Group, comprises those compounds, prodrugs thereof and pharmaceutically acceptable salts of said compounds and said prodrugs, wherein Q is -(C<sub>5</sub>-C<sub>7</sub>)alkylene-.

Another preferred group of compounds within the U Group, designated the W Group, comprises those compounds, prodrugs thereof and pharmaceutically acceptable salts of said compounds and said prodrugs, wherein Q is -CH<sub>2</sub>-X-CH<sub>2</sub>- and X is metaphenylene.

Another preferred group of compounds within the U Group, designated the X Group, comprises those compounds, prodrugs thereof and pharmaceutically acceptable salts of said compounds and said prodrugs, wherein Q is -(C<sub>2</sub>-C<sub>4</sub>)alkylene-X- and X is furanyl, thienyl or thiazolyl.

Another preferred group of compounds within the U Group, designated the Y Group, comprises those compounds, prodrugs thereof and pharmaceutically acceptable salts of said compounds and said prodrugs, wherein Q is -(C<sub>1</sub>-C<sub>2</sub>)-X-O-(C<sub>1</sub>-C<sub>2</sub>)alkylene- and X is metaphenylene.

An especially preferred compound within the Y Group is (3-(((2-(3,5-dichlorophenoxy)-ethyl)-(pyridine-3-sulfonyl)-amino)-methyl)-phenoxy)-acetic acid.

An especially preferred compound within the Y Group is the compound, prodrugs thereof and pharmaceutically acceptable salts of said compounds and said prodrugs, wherein K is ethylenyloxy; said M group being attached to the oxygen atom of the ethylenyloxy group and said N atom being attached to the 2-position of the ethylenyloxy group; Ar is pyrid-3-yl; M is phenyl 3,5-disubstituted with chloro; Z is carboxy and Q is -CH<sub>2</sub>-X-O-CH<sub>2</sub>- wherein X is a second phenyl ring and said CH<sub>2</sub> and

OCH<sub>2</sub> substituents are situated in a meta substitution pattern on said second phenyl ring.

Another preferred group of compounds, designated the Z Group, comprises those compounds of Formula I, prodrugs thereof and pharmaceutically acceptable salts of said compounds and said prodrugs, wherein B is CH.

A preferred group of compounds within the Z Group comprises those compounds, prodrugs thereof and pharmaceutically acceptable salts of said compounds and said prodrugs, wherein A is CO; G is Ar, K is methylenyl, propylenyl, propenylenyl or oxyethylenyl; M is Ar<sup>3</sup> or Ar<sup>4</sup>-Ar<sup>5</sup>; Ar<sup>3</sup> is phenyl or pyridyl; Ar<sup>4</sup> is phenyl, thienyl, pyridyl or furanyl; Ar<sup>5</sup> is (C<sub>5</sub>-C<sub>7</sub>) cycloalkyl, phenyl, pyridyl, imidazolyl, pyrimidyl, thienyl, pyridazinyl, pyrazinyl, imidazolyl, pyrazolyl or thiazolyl; Ar is phenyl, pyrazolyl, pyridazinyl, pyrazinyl, pyridyl, imidazolyl, pyrimidyl, thienyl or thiazolyl, wherein Ar, Ar<sup>3</sup>, Ar<sup>4</sup> and Ar<sup>5</sup> are optionally substituted independently with up to three chloro, fluoro, methyl, difluoromethoxy, trifluoromethoxy or trifluoromethyl.

Another especially preferred group of compounds within the Z Group comprises those compounds, prodrugs thereof and pharmaceutically acceptable salts of said compounds and said prodrugs, wherein A is CO; G is Ar, K is methylenyl, propylenyl, propenylenyl or oxyethylenyl; M is Ar<sup>3</sup> or Ar<sup>4</sup>-Ar<sup>5</sup>; Ar<sup>3</sup> is phenyl or pyridyl; Ar<sup>4</sup> is phenyl, thienyl, pyridyl or furanyl; Ar<sup>5</sup> is (C<sub>5</sub>-C<sub>7</sub>) cycloalkyl, phenyl, pyridyl, imidazolyl, pyrimidyl, thienyl, pyridazinyl, pyrazinyl, imidazolyl, pyrazolyl or thiazolyl; Ar is phenyl, pyrazolyl, pyridazinyl, pyrazinyl, pyridyl, imidazolyl, pyrimidyl, thienyl or thiazolyl, wherein Ar, Ar<sup>3</sup>, Ar<sup>4</sup> and Ar<sup>5</sup> are optionally substituted independently with up to three chloro, fluoro, methyl, difluoromethoxy, trifluoromethoxy or trifluoromethyl. Exemplary five to six membered aromatic rings optionally having one or two heteroatoms selected independently from oxygen, nitrogen and sulfur (i.e., X rings) are phenyl, furyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, pyridyl, pyridiazinyl, pyrimidinyl and pyrazinyl.

Exemplary partially saturated, fully saturated or fully unsaturated five to eight membered rings optionally having one to four heteroatoms selected independently from oxygen, sulfur and nitrogen (i.e., Ar, Ar<sup>1</sup> and Ar<sup>2</sup>) are cyclopentyl, cyclohexyl, cycloheptyl, cyclooctyl and phenyl. Further exemplary five membered rings are furyl, thienyl, 2H-pyrrolyl, 3H-pyrrolyl, pyrrolyl, 2-pyrrolinyl, 3-pyrrolinyl, pyrrolidinyl, 1,3-dioxolanyl, oxazolyl, thiazolyl, imidazolyl, 2H-imidazolyl, 2-imidazolynyl, imidazolidinyl,

-20-

pyrazolyl, 2-pyrazoliny, pyrazolidinyl, isoxazolyl, isothiazolyl, 1,2-dithiolyl, 1,3-dithiolyl, 3H-1,2-oxathiolyl, 1,2,3-oxadiazolyl, 1,2,4-oxadiazolyl, 1,2,5-oxadiazolyl, 1,3,4-oxadiazolyl, 1,2,3-triazolyl, 1,2,4-triazolyl, 1,3,4-thiadiazolyl, 1,2,3,4-oxatriazolyl, 1,2,3,5-oxatriazolyl, 3H-1,2,3-dioxazolyl, 1,2,4-dioxazolyl, 1,3,2-dioxazolyl, 1,3,4-dioxazolyl, 5H-1,2,5-oxathiazolyl and 1,3-oxathiolyl.

Further exemplary six membered rings are 2H-pyranyl, 4H-pyranyl, pyridyl, piperidinyl, 1,2-dioxinyl, 1,3-dioxinyl, 1,4-dioxanyl, morpholinyl, 1,4-dithianyl, thiomorpholinyl, pyridazinyl, pyrimidinyl, pyrazinyl, piperazinyl, 1,3,5-triazinyl, 1,2,4-triazinyl, 1,2,3-triazinyl, 1,3,5-trithianyl, 4H-1,2-oxazinyl, 2H-1,3-oxazinyl, 6H-1,3-oxazinyl, 6H-1,2-oxazinyl, 1,4-oxazinyl, 2H-1,2-oxazinyl, 4H-1,4-oxazinyl, 1,2,5-oxathiazinyl, 1,4-oxazinyl, o-isoxazinyl, p-isoxazinyl, 1,2,5-oxathiazinyl, 1,2,6-oxathiazinyl, 1,4,2-oxadiazinyl and 1,3,5,2-oxadiazinyl.

Further exemplary seven membered rings are azepinyl, oxepinyl, thiepinyl and 1,2,4-diazepinyl.

Further exemplary eight membered rings are cyclooctyl, cyclooctenyl and cyclooctadienyl.

Exemplary bicyclic rings consisting of two fused independently partially saturated, fully saturated or fully unsaturated five and/or six membered rings, taken independently, optionally having one to four heteroatoms selected independently from nitrogen, sulfur and oxygen are indolizinyl, indolyl, isoindolyl, 3H-indolyl, 1H-isoindolyl, indolinyl, cyclopenta(b)pyridinyl, pyrano(3,4-b)pyrrolyl, benzofuryl, isobenzofuryl, benzo(b)thienyl, benzo(c)thienyl, 1H-indazolyl, indoxazinyl, benzoxazolyl, anthranilyl, benzimidazolyl, benzthiazolyl, purinyl, 4H-quinolizinyl, quinolinyl, isoquinolinyl, cinnolinyl, phthalazinyl, quinazolinyl, quinoxalinyl, 1,8-naphthyridinyl, pteridinyl, indenyl, isoindenyl, naphthyl, tetralinyl, decalinyl, 2H-1-benzopyranyl, 1,4-benzodioxan, pyrido(3,4-b)-pyridinyl, pyrido(3,2-b)-pyridinyl, pyrido(4,3-b)-pyridinyl, 2H-1,3-benzoxazinyl, 2H-1,4-benzoxazinyl, 1H-2,3-benzoxazinyl, 4H-3,1-benzoxazinyl, 2H-1,2-benzoxazinyl and 4H-1,4-benzoxazinyl.

Exemplary tricyclic rings consisting of three fused independently partially saturated, fully saturated or fully unsaturated five and/or six membered rings, taken independently, optionally having one to four heteroatoms selected independently from nitrogen, sulfur and oxygen are indacenyl, biphenylenyl, acenaphthylenyl, fluorenyl, phenalenyl, phenanthrenyl, anthracenyl, naphthothienyl, thianthrenyl,

xanthenyl, phenoxathiinyl, carbazolyl, carbolinyl, phenanthridinyl, acridinyl, perimidinyl, phenanthrolinyl, phenazinyl, phenothiazinyl and phenoxazinyl. It will be understood that the fully saturated and all partially unsaturated forms of these rings are within the scope of this invention. Further, it will be understood that nitrogen may  
5 be substituted as the heteroatom at any position, including a bridgehead position, in the heterocyclic rings. Further still, it will be understood that sulfur and oxygen may be substituted as the heteroatom at any non-bridgehead position within the heterocyclic rings.

By alkylene is meant saturated hydrocarbon (straight chain or branched)  
10 wherein a hydrogen atom is removed from each of the terminal carbons. Exemplary of such groups (assuming the designated length encompasses the particular example) are methylene, ethylene, propylene, butylene, pentylene, hexylene and heptylene.

By alkenylene is meant a hydrocarbon containing monounsaturations in the form of one double bond wherein said hydrocarbon is straight chain or branched and  
15 wherein a hydrogen atom is removed from each of the terminal carbons. Exemplary of such groups (assuming the designated length encompasses the particular example) are ethenylene (or vinylene), propenylene, butenylene, pentenylene, hexenylene and heptenylene.

By alkynylene is meant a hydrocarbon containing di-unsaturations in the form of one triple bond wherein said hydrocarbon is straight chain or branched and  
20 wherein a hydrogen atom is removed from each of the terminal carbons. Exemplary of such groups (assuming the designated length encompasses the particular example) are ethynylene, propynylene, butynylene, pentynylene, hexynylene and heptynylene.

25 By halo is meant chloro, bromo, iodo, or fluoro.

By alkyl is meant straight chain saturated hydrocarbon or branched saturated hydrocarbon. Exemplary of such alkyl groups (assuming the designated length encompasses the particular example) are methyl, ethyl, propyl, isopropyl, butyl, sec-butyl, tertiary butyl, pentyl, isopentyl, neopentyl, tertiary pentyl, 1-methylbutyl, 2-  
30 methylbutyl, 3-methylbutyl, hexyl, isohexyl, heptyl and octyl.

By alkoxy is meant straight chain saturated alkyl or branched saturated alkyl bonded through an oxy. Exemplary of such alkoxy groups (assuming the designated length encompasses the particular example) are methoxy, ethoxy, propoxy,

-22-

isopropoxy, butoxy, isobutoxy, tertiary butoxy, pentoxy, isopentoxy, neopentoxy, tertiary pentoxy, hexoxy, isohexoxy, heptoxy and octoxy.

As used herein, the term mono-N- or di-N,N-(C<sub>1</sub>-C<sub>x</sub>)alkyl... refers to the (C<sub>1</sub>-C<sub>x</sub>)alkyl moiety taken independently when it is di-N,N-(C<sub>1</sub>-C<sub>x</sub>)alkyl... (x refers to integers and is taken independently when two (C<sub>1</sub>-C<sub>x</sub>)alkyl groups are present, e.g., methylethylamino is within the scope of di-N,N-(C<sub>1</sub>-C<sub>x</sub>)alkyl).

Unless otherwise stated the "M" moieties defined above are optionally substituted (e.g., the mere listing of a substituent such as R<sup>1</sup> in a subgenus or dependent claim does not mean that M is always substituted with the R<sup>1</sup> moiety unless it is stated that the M moiety is substituted with R<sup>1</sup>). However, in the compounds of Formula I, when K is a bond and M is phenyl, said phenyl group is substituted with one to three substituents. Additionally, in the compounds of Formula I, when Ar or Ar<sup>1</sup> is a fully saturated five to eight membered ring, said ring is unsubstituted.

It is to be understood that if a carbocyclic or heterocyclic moiety may be bonded or otherwise attached to a designated substrate, through differing ring atoms without denoting a specific point of attachment, then all possible points are intended, whether through a carbon atom or, for example, a trivalent nitrogen atom. For example, the term "pyridyl" means 2-, 3-, or 4-pyridyl, the term "thienyl" means 2-, or 3-thienyl, and so forth.

A particularly preferred compound of Formula I is (3-(((4-tert-butyl-benzyl)-(pyridine-3-sulfonyl)-amino)-methyl)-phenoxy)-acetic acid, or a pharmaceutically acceptable salt or prodrug thereof, or a salt of a prodrug. A particularly preferred salt is the sodium salt.

Other EP<sub>2</sub> selective receptor agonists that can be used in the present invention include the prostaglandin receptor agonists disclosed in U.S. patent numbers 6,288,120; and 6,124,314; and PCT published patent application WO 98/58911 (PCT/IB98/00866). A preferred EP<sub>2</sub> compound disclosed in U.S. 6,288,120 is 7-[(4-butyl-benzyl)-methanesulfonyl-amino]-heptanoic acid or a pharmaceutically acceptable salt or prodrug thereof, or a salt of a prodrug. A preferred salt of 7-[(4-butyl-benzyl)-methanesulfonyl-amino]-heptanoic acid is the monosodium salt.

Other EP<sub>2</sub> selective receptor agonists that can be used in the present invention include the compounds disclosed in the following: Burk, Robert M.;

- Holoboski, Mark; Posner, Mari F., Preparation of prostaglandin E2 analogs as EP2-receptor agonists-US patent application no. 2002187961; Burk, Robert M.; Holoboski, Mark; Posner, Mari F., Preparation of prostaglandin E2 analogs as EP2-receptor agonists- US patent 6,376,533; Duckworth, N.; Marshall, K.; Clayton, J. K., An
- 5 investigation of the effect of the prostaglandin EP2 receptor agonist, butaprost, on the human isolated myometrium from pregnant and non-pregnant women, *Journal of Endocrinology* (2002), 172(2), 263-269; Tani, Kousuke; Naganawa, Atsushi; Ishida, Akiharu; Egashira, Hiromu; Odagaki, Yoshihiko; Miyazaki, Toru; Hasegawa, Tomoyuki; Kawanaka, Yasufumi; Nakai, Hisao; Ohuchida, Shuichi; Toda, Masaaki.
- 10 Synthesis of a highly selective EP2-receptor agonist, *Synlett* (2002), (2), 239-242; Tani, Kousuke; Naganawa, Atsushi; Ishida, Akiharu; Egashira, Hiromu; Sagawa, Kenji; Harada, Hiroyuki; Ogawa, Mikio; Maruyama, Takayuki; Ohuchida, Shuichi; Nakai, Hisao; Kondo, Kigen; Toda, Masaaki. Development of a highly selective EP2-receptor agonist. Part 2. Identification of 16-Hydroxy-17,17-trimethylene 9b-chloro
- 15 PGF derivatives, *Bioorganic & Medicinal Chemistry* (2002), 10(4), 1107-1114; Tani, Kousuke; Naganawa, Atsushi; Ishida, Akiharu; Sagawa, Kenji; Harada, Hiroyuki; Ogawa, Mikio; Maruyama, Takayuki; Ohuchida, Shuichi; Nakai, Hisao; Kondo, Kigen; Toda, Masaaki, Development of a highly selective EP2-receptor agonist. Part 1. Identification of 16-hydroxy-17,17-trimethylene PGE2 derivatives, *Bioorganic &*
- 20 *Medicinal Chemistry* (2002), 10(4), 1093-1106; Michelet, Jean-Francois; Mahe, Yann; Bernard, Bruno, Use of non-prostanoid agonists of EP-2 and/or EP-4 prostaglandin receptors as cosmetic agent for reducing or stopping hair loss- European patent application EP 1175891 A1; Tani, K.; Naganawa, A.; Ishida, A.; Egashira, H.; Sagawa, K.; Harada, H.; Ogawa, M.; Maruyama, T.; Ohuchida, S.; Nakai, H.; Kondo, K.; Toda,
- 25 M., Design and Synthesis of a Highly Selective EP2-Receptor Agonist, *Bioorganic & Medicinal Chemistry Letters* (2001), 11(15), 2025-2028; Crider, J. Y.; Sharif, N. A., Functional pharmacological evidence for EP2 and EP4 prostanoid receptors in immortalized human trabecular meshwork and non-pigmented ciliary epithelial cells., *International Journal of Environmental Studies* (2000), 58(1), 35-46; Crider, J. Y.;
- 30 Sharif, N., A. Functional pharmacological evidence for EP2 and EP4 prostanoid receptors in immortalized human trabecular meshwork and nonpigmented ciliary epithelial cells. *Journal of Ocular Pharmacology and Therapeutics* (2001), 17(1), 35-46; Klimko, Peter G.; Sharif, Najam A.; Griffin, Brenda W. Prostaglandin E agonists



- for treatment of glaucoma- WO 0038667 A2; Woodward, David F., EP2 receptor agonists as neuroprotective agents for the eye-US 5877211; Regan, John W.; Gil, Daniel W.; Woodward, David F., Cloning of a novel human prostaglandin receptor with characteristics of the pharmacologically defined EP2 subtype- US 5716835;
- 5 Woodward, David F. EP2-receptor agonists as agents for lowering intraocular pressure-US 5698598; Woodward, David F. EP2-receptor agonists as agents for lowering intraocular pressure.-WO 9519964; Woodward, D. F.; Bogardus, A. M.; Donello, J. E.; Fairbairn, C. E.; Gil, D. W.; Kedzie, K. M.; Burke, J. A.; Kharlamb, A.; Runde, E.; et al., Molecular characterization and ocular hypotensive properties of the
- 10 prostanoid EP2 receptor, *Journal of Ocular Pharmacology and Therapeutics* (1995), 11(3), 447-54; Nials, Anthony T.; Vardey, Christopher J.; Denyer, Lois H.; Thomas, Malcolm; Sparrow, Susan J.; Shepherd, Gillian D.; Coleman, Robert A., AH13205, a selective prostanoid EP2-receptor agonist, *Cardiovascular Drug Reviews* (1993), 11(2), 165-79; and Woodward, D. F.; Protzman, C. E.; Krauss, A. H. P.; Williams, L.
- 15 S., Identification of 19(R)-OH prostaglandin E2 as a selective prostanoid EP2-receptor agonist, *Prostaglandins* (1993), 46(4), 371-83.

The present methods can be used to treat pulmonary hypertension. Pulmonary hypertension, also known as primary pulmonary hypertension, is a disease of unknown origin that involves the medium and small pulmonary arteries

20 and results in right ventricular failure or fatal syncope, typically 2 to 5 years after detection. Intimal hyperplasia and consequent narrowing of the vessel lumen are always present. Areas of medial hypertrophy and hyperplasia, irreversible plexiform lesions, and necrotizing arteritis (plexogenic arteriopathy) occur in more advanced cases. Those skilled in the art are familiar with the identification of patients having

25 pulmonary hypertension.

The present methods can also be used in facilitating joint fusion. Examples of joint fusions include fusion of bones in the wrist or ankle as well as other joints. In a joint fusion, two or more bones are fused together.

The present methods can also be used to facilitate tendon and/or ligament

30 repair. The repair can comprise the strengthening of a tendon or ligament or can comprise the reconstruction of a damaged portion of a tendon and/or ligament. Another aspect of tendon and ligament repair is to strengthen or repair the attachment of a tendon or ligament to a bone.

The present methods can also be used to reduce the occurrence of secondary fractures. A secondary fracture is a fracture subsequent to a primary fracture. Once a fracture has occurred, the present methods can be used to prevent another fracture from occurring or reduce the magnitude or complexity of any secondary fracture. For example, if a patient has a hip fracture, the present method may be used to help avoid or ameliorate the extent of a second fracture in the hip either on the same side of the hip as the first fracture or on the other side of the hip. The prevention or amelioration of damage caused by secondary fractures is also important in cases where the risk of secondary fractures is enhanced, such as in response to chemotherapy. Also, the prevention of secondary fractures in the spine and spinal stabilization are important.

The present invention also provides methods of treating avascular necrosis. Avascular necrosis is characterized by cell death in bone resulting from a compromised blood supply. The hip, femur and shoulder are commonly affected bones. Various conditions have been associated with avascular necrosis including fracture of the femoral neck, hip dislocation, decompression sickness, sickle cell disease, radiotherapy, Gaucher's disease, and corticosteroid high-dose therapy. Other conditions that have been associated with avascular necrosis include systemic lupus erythmatosus, renal transplantation, polycythemia versa, Cushing's syndrome, Diabetes mellitus, atherosclerosis, cytotoxic chemotherapy, alcohol abuse, fatty liver, psoriasis, pancreatitis, pancreatic cancer and gout. Thus, the presence of a condition associated with avascular necrosis can be an indicator to apply the methods of the present invention to prevent or ameliorate the occurrence of avascular necrosis in a patient.

The present invention also be used to facilitate cartilage repair, facilitate bone healing after limb transplantation, facilitate liver regeneration, facilitate wound healing, reduce the occurrence of gastric ulceration, treat hypertension, facilitate the growth of tooth enamel or finger or toe nails, treat glaucoma, treat ocular hypertension, or repair damage caused by metastatic bone disease, which conditions are well known to those skilled in the art. Patients having such conditions are easily identified by those skilled in the art.

The term "facilitating", "facilitate", and the like, with regard to the present methods, means to make the methods less difficult or improve the speed of the

-26-

methods. For example, facilitating bone fusion means to make the procedure less difficult to accomplish or proceed more rapidly in the presence of an EP<sub>2</sub> selective receptor agonist than in the absence of an EP<sub>2</sub> selective receptor agonist.

5 A preferred dosage is about 0.001 to 100 mg/kg/day of an EP<sub>2</sub> selective receptor agonist, such as a compound of Formula I. An especially preferred dosage is about 0.01 to 10 mg/kg/day of an EP<sub>2</sub> selective receptor agonist, such as a compound of Formula I.

The present invention is also concerned with pharmaceutical compositions comprising an EP<sub>2</sub> selective receptor agonist, such as a compound of Formula I, and  
10 a carrier, solvent, diluent and the like.

Another aspect of this invention is directed to combinations of an EP<sub>2</sub> selective receptor agonist, such as a compound of Formula I, and other therapeutically useful compounds.

In one embodiment, the combinations of this invention comprise a  
15 therapeutically effective amount of a first compound, said first compound being an EP<sub>2</sub> selective receptor agonist, such as a compound of Formula I; and a therapeutically effective amount of a second compound, the second compound being an anti-resorptive agent such as an estrogen agonist/antagonist or a bisphosphonate. An estrogen agonist/antagonist is also called a selective estrogen receptor  
20 modulator (SERM).

It is noted that when compounds are discussed herein, it is contemplated that the compounds may be administered to a patient as a therapeutically acceptable salt, prodrug, or a salt of a prodrug. All such variations are intended to be included in the invention.

25 Another aspect of this invention is a kit comprising:

a. an amount of an EP<sub>2</sub> selective receptor agonist, such as a compound of Formula I, and a pharmaceutically acceptable carrier or diluent in a first unit dosage form;

b. an amount of an anti-resorptive agent, and a pharmaceutically acceptable  
30 carrier or diluent in a second unit dosage form; and

c. a container.

Yet another aspect of this invention is directed to pharmaceutical compositions or kits comprising an EP<sub>2</sub> selective receptor agonist, such as a

-27-

compound of Formula I, and another bone anabolic agent (although the other bone anabolic agent may be another EP<sub>2</sub> selective receptor agonist, such as a different compound of Formula I). Such compositions comprise a therapeutically effective amount of a first compound, said first compound being an EP<sub>2</sub> selective receptor agonist, such as a compound of Formula I; and a therapeutically effective amount of a second compound, said second compound being another bone anabolic agent.

The first compound can administered at the same time as the second compound in the same dosage form or in different dosage forms. Alternatively, the first compound and second compound can be administered at different times.

Moreover, the combinations of the present invention can comprise more than two compounds. For example, two EP<sub>2</sub> selective receptor agonists and an anti-repsorptive or bone anabolic compound can be administered to a patient.

Another aspect of this invention is a kit comprising:

- a. an amount of an EP<sub>2</sub> receptor selective agonist, such as a compound of Formula I, in a first unit dosage form;
- b. an amount of a second compound, said second compound being another bone anabolic agent in a second unit dosage form; and
- c. a container.

Preferred bone anabolic agents include IGF-1, prostaglandins, prostaglandin agonists/antagonists, sodium fluoride, parathyroid hormone (PTH), active fragments of parathyroid hormone, parathyroid hormone related peptides and active fragments and analogues of parathyroid hormone related peptides, growth hormones' or growth hormone secretagogues and the pharmaceutically acceptable salts or prodrugs thereof or a salt of a prodrug.

Since the present invention has an aspect that relates to a combination of active ingredients, which may be administered separately, the invention also relates to combining separate pharmaceutical compositions in kit form. The kit comprises two separate pharmaceutical compositions: an EP<sub>2</sub> selective receptor agonist, such as a compound of Formula I and a second compound as described above. The kit comprises container means for containing the separate compositions such as a divided bottle or a divided foil packet, however, the separate compositions may also be contained within a single, undivided container. Typically, the kit comprises directions for the administration of the separate components. The kit form is

-28-

particularly advantageous when the separate components are preferably administered in different dosage forms (e.g., oral and parenteral), are administered at different dosage intervals, or when titration of the individual components of the combination is desired by the prescribing physician.

5           An example of such a kit is a so-called blister pack. Blister packs are well known in the packaging industry and are being widely used for the packaging of pharmaceutical unit dosage forms (tablets, capsules, and the like). Blister packs generally consist of a sheet of relatively stiff material covered with a foil of a preferably transparent plastic material. During the packaging process recesses are  
10           formed in the plastic foil. The recesses have the size and shape of the tablets or capsules to be packed. Next, the tablets or capsules are placed in the recesses and the sheet of relatively stiff material is sealed against the plastic foil at the face of the foil which is opposite from the direction in which the recesses were formed. As a result, the tablets or capsules are sealed in the recesses between the plastic foil and  
15           the sheet. Preferably the strength of the sheet is such that the tablets or capsules can be removed from the blister pack by manually applying pressure on the recesses whereby an opening is formed in the sheet at the place of the recess. The tablet or capsule can then be removed via said opening.


          It may be desirable to provide a memory aid on the kit, e.g., in the form of  
20           numbers next to the tablets or capsules whereby the numbers correspond with the days of the regimen which the dosage form so specified should be ingested. Another example of such a memory aid is a calendar printed on the card e.g., as follows "First Week, Monday, Tuesday, ...etc.... Second Week, Monday, Tuesday,..." etc. Other variations of memory aids will be readily apparent. A "daily dose" can be a single  
25           tablet or capsule or several tablets or capsules to be taken on a given day. Also, a daily dose of a Formula I compound, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug can consist of one tablet or capsule while a daily dose of the second compound can consist of several tablets or capsules and vice versa. The memory aid should reflect this.

30           In another specific embodiment of the invention, a dispenser designed to dispense the daily doses one at a time in the order of their intended use is provided. Preferably, the dispenser is equipped with a memory-aid, so as to further facilitate compliance with the regimen. An example of such a memory-aid is a mechanical

-29-

counter which indicates the number of daily doses that has been dispensed. Another example of such a memory-aid is a battery-powered micro-chip memory coupled with a liquid crystal readout, or audible reminder signal which, for example, reads out the date that the last daily dose has been taken and/or reminds one when the next dose is to be taken.

Preferred estrogen agonists / antagonists of the present invention include the compounds described in U.S. patent no. 5,552,412. Those compounds are described by the formula designated herein as formula (I) given below:



wherein:

A is selected from CH<sub>2</sub> and NR;

B, D and E are independently selected from CH and N;

30 Y is

(a) phenyl, optionally substituted with 1-3 substituents independently selected from R<sup>4</sup>;

(b) naphthyl, optionally substituted with 1-3 substituents independently selected from R<sup>4</sup>;

35 (c) C<sub>3</sub>-C<sub>8</sub> cycloalkyl, optionally substituted with 1-2 substituents independently selected from R<sup>4</sup>;

(d) C<sub>3</sub>-C<sub>8</sub> cycloalkenyl, optionally substituted with 1-2 substituents independently selected from R<sup>4</sup>;

40 (e) a five membered heterocycle containing up to two heteroatoms selected from the group consisting of -O-, -NR<sup>2</sup>- and -S(O)<sub>n</sub>-, optionally substituted with 1-3 substituents independently selected from R<sup>4</sup>;

-30-

(f) a six membered heterocycle containing up to two heteroatoms selected from the group consisting of -O-, -NR<sup>2</sup>- and -S(O)<sub>n</sub>- optionally substituted with 1-3 substituents independently selected from R<sup>4</sup>; or


- 5 (g) a bicyclic ring system consisting of a five or six membered heterocyclic ring fused to a phenyl ring, said heterocyclic ring containing up to two heteroatoms selected from the group consisting of -O-, -NR<sup>2</sup>- and -S(O)<sub>n</sub>-, optionally substituted with 1-3 substituents independently selected from R<sup>4</sup>;

Z<sup>1</sup> is

- 10 (a) -(CH<sub>2</sub>)<sub>p</sub> W(CH<sub>2</sub>)<sub>q</sub>-;  
 (b) -O(CH<sub>2</sub>)<sub>p</sub> CR<sup>5</sup>R<sup>6</sup>-;  
 (c) -O(CH<sub>2</sub>)<sub>p</sub>W(CH<sub>2</sub>)<sub>q</sub>-;  
 (d) -OCHR<sup>2</sup>CHR<sup>3</sup>-; or  
 (e) -SCHR<sup>2</sup>CHR<sup>3</sup>-;

G is


- 15 (a) -NR<sup>7</sup>R<sup>8</sup>;



- 20 wherein n is 0, 1 or 2; m is 1, 2 or 3; Z<sup>2</sup> is -NH-, -O-, -S-, or -CH<sub>2</sub>-; optionally fused on adjacent carbon atoms with one or two phenyl rings and, optionally independently substituted on carbon with one to three substituents and, optionally, independently on nitrogen with a chemically suitable substituent selected from R<sup>4</sup>; or

- 25 (c) a bicyclic amine containing five to twelve carbon atoms, either bridged or fused and optionally substituted with 1-3 substituents independently selected from R<sup>4</sup>; or

Z<sup>1</sup> and G in combination may be




30

W is

-31-


5

- (a)  $-\text{CH}_2-$ ;
- (b)  $-\text{CH}=\text{CH}-$ ;
- (c)  $-\text{O}-$ ;
- (d)  $-\text{NR}^2-$ ;
- (e)  $-\text{S}(\text{O})_n-$ ;



- (g)  $-\text{CR}^2(\text{OH})-$ ;
- (h)  $-\text{CONR}^2-$ ;
- (i)  $-\text{NR}^2\text{CO}-$ ;

10

R is hydrogen or  $\text{C}_1$ - $\text{C}_6$  alkyl; $\text{R}^2$  and  $\text{R}^3$  are independently

15

- (a) hydrogen; or
- (b)  $\text{C}_1$ - $\text{C}_4$  alkyl;

 $\text{R}^4$  is

20

- (a) hydrogen;
- (b) halogen;
- (c)  $\text{C}_1$ - $\text{C}_6$  alkyl;
- (d)  $\text{C}_1$ - $\text{C}_4$  alkoxy;
- (e)  $\text{C}_1$ - $\text{C}_4$  acyloxy;
- (f)  $\text{C}_1$ - $\text{C}_4$  alkylthio;
- (g)  $\text{C}_1$ - $\text{C}_4$  alkylsulfinyl;
- (h)  $\text{C}_1$ - $\text{C}_4$  alkylsulfonyl;
- (i) hydroxy ( $\text{C}_1$ - $\text{C}_4$ )alkyl;
- (j) aryl ( $\text{C}_1$ - $\text{C}_4$ )alkyl;
- (k)  $-\text{CO}_2\text{H}$ ;
- (l)  $-\text{CN}$ ;
- (m)  $-\text{CONHOR}$ ;
- (n)  $-\text{SO}_2\text{NHR}$ ;

30



-32-

- (o)  $-\text{NH}_2$ ;  
(p)  $\text{C}_1\text{-C}_4$  alkylamino;  
(q)  $\text{C}_1\text{-C}_4$  dialkylamino;  
(r)  $-\text{NHSO}_2\text{R}$ ;  
5 (s)  $-\text{NO}_2$ ;  
(t) -aryl; or  
(u)  $-\text{OH}$ ;

$\text{R}^5$  and  $\text{R}^6$  are independently  $\text{C}_1\text{-C}_8$  alkyl or together form a  $\text{C}_3\text{-C}_{10}$  carbocyclic ring;

10  $\text{R}^7$  and  $\text{R}^8$  are independently

- (a) phenyl;  
(b) a  $\text{C}_3\text{-C}_{10}$  carbocyclic ring, saturated or unsaturated;  
(c) a  $\text{C}_3\text{-C}_{10}$  heterocyclic ring containing up to two heteroatoms,  
selected from -O-, -N- and -S-;  
15 (d) H;  
(e)  $\text{C}_1\text{-C}_6$  alkyl; or  
(f) form a 3 to 8 membered nitrogen containing ring with  $\text{R}^5$  or  $\text{R}^6$ ;

$\text{R}^7$  and  $\text{R}^8$  in either linear or ring form may optionally be substituted with up to three substituents independently selected from  $\text{C}_1\text{-C}_6$  alkyl, halogen, alkoxy, hydroxy and carboxy;

20 a ring formed by  $\text{R}^7$  and  $\text{R}^8$  may be optionally fused to a phenyl ring;

e is 0, 1 or 2;

m is 1, 2 or 3;

n is 0, 1 or 2;


25 p is 0, 1, 2 or 3;

q is 0, 1, 2 or 3;


and optical and geometric isomers thereof; and nontoxic pharmaceutically acceptable acid addition salts, N-oxides, esters, quaternary ammonium salts and prodrugs thereof.

30 Additional preferred estrogen agonists/antagonists are disclosed in U.S. patent no. 5,552,412 and are described by the formula designated herein as formula (IA):

-33-



5 wherein G is



10  $R^4$  is H, OH, F, or Cl; and B and E are independently selected from CH and N, and optical and geometric isomers thereof; and nontoxic pharmaceutically acceptable acid addition salts, N-oxides, esters, quaternary ammonium salts and prodrugs thereof.

Especially preferred estrogen agonists/antagonists for the methods of the invention are:

cis-6-(4-fluoro-phenyl)-5-[4-(2-piperidin-1-yl-ethoxy)-phenyl]-5,6,7,8-tetrahydro-naphthalene-2-ol;

15 (-)-cis-6-phenyl-5-[4-(2-pyrrolidin-1-yl-ethoxy)-phenyl]-5,6,7,8-tetrahydro-naphthalene-2-ol;

cis-6-phenyl-5-[4-(2-pyrrolidin-1-yl-ethoxy)-phenyl]-5,6,7,8-tetrahydro-naphthalene-2-ol;

20 cis-1-[6'-pyrrolidinoethoxy-3'-pyridyl]-2-phenyl-6-hydroxy-1,2,3,4-tetrahydronaphthalene;

1-(4'-pyrrolidinoethoxyphenyl)-2-(4"-fluorophenyl)-6-hydroxy-1,2,3,4-tetrahydroisoquinoline;

-34-


cis-6-(4-hydroxyphenyl)-5-[4-(2-piperidin-1-yl-ethoxy)-phenyl]-5,6,7,8-tetrahydro-naphthalene-2-ol;

1-(4'-pyrrolidinoethoxyphenyl)-2-phenyl-6-hydroxy-1,2,3,4-tetrahydroisoquinoline and pharmaceutically acceptable salts thereof.

5 An especially preferred salt of (-)-cis-6-phenyl-5-[4-(2-pyrrolidin-1-yl-ethoxy)-phenyl]-5,6,7,8-tetrahydro-naphthalene-2-ol is the D-tartrate salt.

Other preferred estrogen agonists / antagonists are disclosed in U.S. Patent 5,047,431. The structure of these compounds are described by the formula designated herein as formula (II) below:

10



(II)

wherein

15 R<sup>1A</sup> and R<sup>2A</sup> may be the same or different and are either H, methyl, ethyl or a benzyl group; and optical or geometric isomers thereof; and pharmaceutically acceptable salts, N-oxides, esters, quaternary ammonium salts, and prodrugs thereof.

20 Additional preferred estrogen agonists / antagonists are the compounds disclosed in U.S. Patent No. 4,536,516; 4-hydroxy tamoxifen (i.e., tamoxifen wherein the 2-phenyl moiety has a hydroxy group at the 4 position) and other compounds as disclosed in U.S. Patent No. 4,623,660; raloxifene: (methanone, [6-hydroxy-2-(4-

-35-

hydroxyphenyl)benzo[b]thien-3-yl][4-[2-(1-piperidinyloxy)phenyl]-,hydrochloride) and other compounds as disclosed in U.S. Patent Numbers 4,418,068; 5,393,763; 5,457,117; 5,478,847 and 5,641,790; toremifene: (ethanamine, 2-[4-(4-chloro-1,2-diphenyl-1-butenyl)phenoxy]-N,N-dimethyl-, (Z)-, 2-hydroxy-1,2,3-


5 propanetricarboxylate (1:1) and other compounds as disclosed in U.S. Patent Numbers 4,696,949 and 4,996,225; centchroman: 1-[2-[[4-(methoxy-2,2, dimethyl-3-phenyl-chroman-4-yl)-phenoxy]-ethyl]-pyrrolidine and other compounds as disclosed in U.S. Patent No. 3,822,287; idoxifene: pyrrolidine, 1-[4-[1-(4-iodophenyl)-2-phenyl-1-butenyl]phenoxy]ethyl] and other compounds as disclosed in U.S. Patent

10 No. 4,839,155; 6-(4-hydroxy-phenyl)-5-[4-(2-piperidin-1-yl-ethoxy)-benzyl]-naphthalen-2-ol and other compounds as disclosed in U.S. Patent No. 5,484,795; and {4-[2-(2-aza-bicyclo[2.2.1]hept-2-yl)-ethoxy]-phenyl}-[6-hydroxy-2-(4-hydroxy-phenyl)-benzo[b]thiophen-3-yl]-methanone and other compounds as disclosed in published international patent application WO 95/10513. Other preferred compounds

15 include GW 5638 and GW 7604, the synthesis of which is described in Willson et al., J. Med. Chem., 1994; 37: 1550-1552.


Further preferred estrogen agonists / antagonists include EM-652 (as shown in the formula designated herein as formula (III)) and EM-800 (as shown in the formula designated herein as formula (IV)). The synthesis of EM-652 and EM-800

20 and the activity of various enantiomers is described in Gauthier et al., J. Med. Chem., 1997; 40: 2117-2122.



(III)



-36-



(IV)

Further preferred estrogen agonists / antagonists include TSE 424 and other  
5 compounds disclosed in U.S. Patent No. 5,998,402, U.S. Patent No. 5,985,910, U.S.  
Patent No. 5,780,497, U.S. Patent No. 5,880,137, and European Patent Application  
EP 0802183 A1 including the compounds described by the formulae designated  
herein as formulae V and VI, below:

-37-



wherein:

$R_{1B}$  is selected from H, OH or the  $C_1$ - $C_{12}$  esters (straight chain or branched) or  $C_1$ - $C_{12}$  (straight chain or branched or cyclic) alkyl ethers thereof, or halogens; or  
 25  $C_1$ - $C_4$  halogenated ethers including trifluoromethyl ether and trichloromethyl ether.


$R_{2B}$ ,  $R_{3B}$ ,  $R_{4B}$ ,  $R_{5B}$ , and  $R_{6B}$  are independently selected from H, OH or the  $C_1$ - $C_{12}$  esters (straight chain or branched) or  $C_1$ - $C_{12}$  alkyl ethers (straight chain or branched or cyclic) thereof, halogens, or  $C_1$ - $C_4$  halogenated ethers including trifluoromethyl ether and trichloromethyl ether, cyano,  $C_1$ - $C_6$  alkyl (straight chain or  
 30 branched), or trifluoromethyl;

$X_A$  is selected from H,  $C_1$ - $C_6$  alkyl, cyano, nitro, trifluoromethyl, and halogen;  
 s is 2 or 3;

-38-

$Y_A$  is selected from:

a) the moiety:



wherein  $R_{7B}$  and  $R_{8B}$  are independently selected from the group of H,  $C_1$ - $C_6$  alkyl, or phenyl optionally substituted by CN,  $C_1$ - $C_6$  alkyl (straight chain or branched),  $C_1$ - $C_6$  alkoxy (straight chain or branched), halogen, -OH, - $CF_3$ , or - $OCF_3$ ;

b) a five-membered saturated, unsaturated or partially unsaturated heterocycle containing up to two heteroatoms selected from the group consisting of -O-, -NH-, -N( $C_1$ - $C_4$  alkyl)-, -N=, and -S(O) $_u$ -, wherein u is an integer of from 0-2, optionally substituted with 1-3 substituents independently selected from the group consisting of hydrogen, hydroxyl, halo,  $C_1$ - $C_4$  alkyl, trihalomethyl,  $C_1$ - $C_4$  alkoxy, trihalomethoxy,  $C_1$ - $C_4$  acyloxy,  $C_1$ - $C_4$  alkylthio,  $C_1$ - $C_4$  alkylsulfinyl,  $C_1$ - $C_4$  alkylsulfonyl, hydroxy ( $C_1$ - $C_4$ )alkyl, -CO $_2$ H, -CN, -CONHR $_{1B}$ , -NH $_2$ ,  $C_1$ - $C_4$  alkylamino, di( $C_1$ - $C_4$ )alkylamino, -NHSO $_2$ R $_{1B}$ , -NHCOR $_{1B}$ , -NO $_2$ , and phenyl optionally substituted with 1-3 ( $C_1$ - $C_4$ )alkyl;

c) a six-membered saturated, unsaturated or partially unsaturated heterocycle containing up to two heteroatoms selected from the group consisting of -O-, -NH-, -N( $C_1$ - $C_4$  alkyl)-, -N=, and -S(O) $_u$ -, wherein u is an integer of from 0-2, optionally substituted with 1-3 substituents independently selected from the group consisting of hydrogen, hydroxyl, halo,  $C_1$ - $C_4$  alkyl, trihalomethyl,  $C_1$ - $C_4$  alkoxy, trihalomethoxy,  $C_1$ - $C_4$  acyloxy,  $C_1$ - $C_4$  alkylthio,  $C_1$ - $C_4$  alkylsulfinyl,  $C_1$ - $C_4$  alkylsulfonyl, hydroxy ( $C_1$ - $C_4$ )alkyl, -CO $_2$ H, -CN, -CONHR $_1$ , -NH $_2$ ,  $C_1$ - $C_4$  alkylamino, di( $C_1$ - $C_4$ )alkylamino, -NHSO $_2$ R $_{1B}$ , -NHCOR $_{1B}$ , -NO $_2$ , and phenyl optionally substituted with 1-3 ( $C_1$ - $C_4$ )alkyl;

d) a seven-membered saturated, unsaturated or partially unsaturated heterocycle containing up to two heteroatoms selected from the group consisting of -O-, -NH-, -N( $C_1$ - $C_4$  alkyl)-, -N=, and -S(O) $_u$ -, wherein u is an integer of from 0-2,

-39-

optionally substituted with 1-3 substituents independently selected from the group consisting of hydrogen, hydroxyl, halo, C<sub>1</sub>-C<sub>4</sub> alkyl, trihalomethyl, C<sub>1</sub>-C<sub>4</sub> alkoxy, trihalomethoxy, C<sub>1</sub>-C<sub>4</sub> acyloxy, C<sub>1</sub>-C<sub>4</sub> alkylthio, C<sub>1</sub>-C<sub>4</sub> alkylsulfinyl, C<sub>1</sub>-C<sub>4</sub> alkylsulfonyl, hydroxy (C<sub>1</sub>-C<sub>4</sub>)alkyl, -CO<sub>2</sub>H, -CN, -CONHR<sub>1B</sub>, -NH<sub>2</sub>, C<sub>1</sub>-C<sub>4</sub> alkylamino, di(C<sub>1</sub>-C<sub>4</sub>)alkylamino, -NHSO<sub>2</sub>R<sub>1B</sub>, -NHCOR<sub>1B</sub>, -NO<sub>2</sub>, and phenyl optionally substituted with 1-3 (C<sub>1</sub>-C<sub>4</sub>)alkyl; or

e) a bicyclic heterocycle containing from 6-12 carbon atoms either bridged or fused and containing up to two heteroatoms selected from the group consisting of -O-, -NH-, -N(C<sub>1</sub>-C<sub>4</sub> alkyl)-, and -S(O)<sub>u</sub>-, wherein u is an integer of from 0-2, optionally substituted with 1-3 substituents independently selected from the group consisting of hydrogen, hydroxyl, halo, C<sub>1</sub>-C<sub>4</sub> alkyl, trihalomethyl, C<sub>1</sub>-C<sub>4</sub> alkoxy, trihalomethoxy, C<sub>1</sub>-C<sub>4</sub> acyloxy, C<sub>1</sub>-C<sub>4</sub> alkylthio, C<sub>1</sub>-C<sub>4</sub> alkylsulfinyl, C<sub>1</sub>-C<sub>4</sub> alkylsulfonyl, hydroxy (C<sub>1</sub>-C<sub>4</sub>)alkyl, -CO<sub>2</sub>H-, -CN-, -CONHR<sub>1B</sub>-, -NH<sub>2</sub>-, -N=, C<sub>1</sub>-C<sub>4</sub> alkylamino, di(C<sub>1</sub>-C<sub>4</sub>)alkylamino, -NHSO<sub>2</sub>R<sub>1B</sub>, -NHCOR<sub>1B</sub>, -NO<sub>2</sub>, and phenyl optionally substituted with 1-3 (C<sub>1</sub>-C<sub>4</sub>) alkyl; and optical and geometric isomers thereof; and nontoxic pharmaceutically acceptable acid addition salts, N-oxides, esters, quaternary ammonium salts, and prodrugs thereof.


Preferred compounds of this invention are those having the general structures V or VI, above, wherein:

R<sub>1B</sub> is selected from H, OH or the C<sub>1</sub>-C<sub>12</sub> esters or alkyl ethers thereof, and halogen;

R<sub>2B</sub>, R<sub>3B</sub>, R<sub>4B</sub>, R<sub>5B</sub>, and R<sub>6B</sub> are independently selected from H, OH or the C<sub>1</sub>-C<sub>12</sub> esters or alkyl ethers thereof, halogen, cyano, C<sub>1</sub>-C<sub>6</sub> alkyl, or trihalomethyl, preferably trifluoromethyl, with the proviso that, when R<sub>1B</sub> is H, R<sub>2B</sub> is not OH;

X<sub>A</sub> is selected from H, C<sub>1</sub>-C<sub>6</sub> alkyl, cyano, nitro, trifluoromethyl, and halogen;

Y<sub>A</sub> is the moiety:






-40-

- $R_{7B}$  and  $R_{8B}$  are selected independently from H, C<sub>1</sub>-C<sub>6</sub> alkyl, or combined by  $-(CH_2)_w$ , wherein w is an integer of from 2 to 6, so as to form a ring, the ring being optionally substituted by up to three substituents selected from the group of hydrogen, hydroxyl, halo, C<sub>1</sub>-C<sub>4</sub> alkyl, trihalomethyl, C<sub>1</sub>-C<sub>4</sub> alkoxy, trihalomethoxy, C<sub>1</sub>-C<sub>4</sub> alkylthio, C<sub>1</sub>-C<sub>4</sub> alkylsulfinyl, C<sub>1</sub>-C<sub>4</sub> alkylsulfonyl, hydroxy (C<sub>1</sub>-C<sub>4</sub>)alkyl, -CO<sub>2</sub>H, -CN, -CONH(C<sub>1</sub>-C<sub>4</sub>alkyl), -NH<sub>2</sub>, C<sub>1</sub>-C<sub>4</sub> alkylamino, C<sub>1</sub>-C<sub>4</sub> dialkylamino, -NHSO<sub>2</sub>(C<sub>1</sub>-C<sub>4</sub>alkyl), -CO(C<sub>1</sub>-C<sub>4</sub>alkyl), and -NO<sub>2</sub>; and optical and geometric isomers thereof; and nontoxic pharmaceutically acceptable acid addition salts, N-oxides, esters, quaternary ammonium salts, and prodrugs thereof.
- The rings formed by a concatenated  $R_{7B}$  and  $R_{8B}$ , mentioned above, may include, but are not limited to, aziridine, azetidine, pyrrolidine, piperidine, hexamethylenamine or heptamethylenamine rings.

- Preferred compounds of structural formulas V and VI, above, are those wherein  $R_{1B}$  is OH;  $R_{2B}$  -  $R_{6B}$  are as defined above;  $X_A$  is selected from the group of Cl, NO<sub>2</sub>, CN, CF<sub>3</sub>, or CH<sub>3</sub>;  $Y_A$  is the moiety




- and  $R_{7B}$  and  $R_{8B}$  are concatenated together as  $-(CH_2)_t$ , wherein t is an integer of from 4 to 6, to form a ring optionally substituted by up to three substituents selected from the group of hydrogen, hydroxyl, halo, C<sub>1</sub>-C<sub>4</sub> alkyl, trihalomethyl, C<sub>1</sub>-C<sub>4</sub> alkoxy, trihalomethoxy, C<sub>1</sub>-C<sub>4</sub> alkylthio, C<sub>1</sub>-C<sub>4</sub> alkylsulfinyl, C<sub>1</sub>-C<sub>4</sub> alkylsulfonyl, hydroxy (C<sub>1</sub>-C<sub>4</sub>)alkyl, -CO<sub>2</sub>H, -CN, -CONH(C<sub>1</sub>-C<sub>4</sub>)alkyl, -NH<sub>2</sub>, C<sub>1</sub>-C<sub>4</sub> alkylamino, di(C<sub>1</sub>-C<sub>4</sub>)alkylamino, -NHSO<sub>2</sub>(C<sub>1</sub>-C<sub>4</sub>)alkyl, -NHCO(C<sub>1</sub>-C<sub>4</sub>)alkyl, and -NO<sub>2</sub>; and optical and geometric isomers thereof; and nontoxic pharmaceutically acceptable acid addition salts, N-oxides, esters, quaternary ammonium salts, and prodrugs thereof.

Another preferred compound is TSE-424 as described by the formula designated herein as formula (Va) below:

30

-41-



Another estrogen agonist/antagonist that can be used in the combination aspect of the present invention is arzoxifene, which is disclosed in U.S. patent no. 5,723,474.

A particularly preferred combination of an EP<sub>2</sub> selective receptor agonist and an estrogen agonist/antagonist is (3-(((4-tert-butyl-benzyl)-(pyridine-3-sulfonyl)-amino)-methyl)-phenoxy)-acetic acid and (-)-*cis*-6-phenyl-5-(4-(2-pyrrolidin-1-yl-ethoxy)-phenyl)-5,6,7,8-tetrahydro-naphthalene-2-ol. In a more preferred combination, the (3-(((4-tert-butyl-benzyl)-(pyridine-3-sulfonyl)-amino)-methyl)-phenoxy)-acetic acid is in the form of the sodium salt, and the (-)-*cis*-6-phenyl-5-(4-(2-pyrrolidin-1-yl-ethoxy)-phenyl)-5,6,7,8-tetrahydro-naphthalene-2-ol is in the form of the D-tartrate salt.

Preferred bisphosphonates include, tiludronic acid, alendronic acid, zoledronic acid, ibandronic acid, risedronic acid, etidronic acid, clodronic acid, and pamidronic acid and their pharmaceutically acceptable salts or prodrugs or salts of the prodrugs.

It will be recognized that prodrugs and pharmaceutically acceptable salts may be formed from the compounds of this invention. All of such prodrugs and pharmaceutically acceptable salts so formed are within the scope of this invention. Particularly preferred salt forms of the estrogen agonists/antagonists include, raloxifene hydrochloride, tamoxifen citrate, toremifene citrate, and lasofoxifene tartrate.

Those skilled in the art will recognize that anti-resorptive agents such as progestins, polyphosphonates, bisphosphonate(s), estrogen agonists/antagonists, estrogen, estrogen/progestin combinations, Premarin® (conjugated estrogens), estrone, estriol or 17 $\alpha$ - or 17 $\beta$ -ethynyl estradiol may be used in conjunction with the  
5 EP<sub>2</sub> selective receptor agonists in the present methods.

Exemplary progestins are available from commercial sources and include: algestone acetophenide, altrenogest, amadinone acetate, anagestone acetate, chlormadinone acetate, cingestol, clogestone acetate, clomegestone acetate, delmadinone acetate, desogestrel, dimethisterone, dydrogesterone, ethynerone,  
10 ethynodiol diacetate, etonogestrel, flurogestone acetate, gestaclone, gestodene, gestonorone caproate, gestrinone, haloprogestone, hydroxyprogesterone caproate, levonorgestrel, lynestrenol, medrogestone, medroxyprogesterone acetate, melengestrol acetate, methynodiol diacetate, norethindrone, norethindrone acetate, norethynodrel, norgestimate, norgestomet, norgestrel, oxogestone phenpropionate,  
15 progesterone, quingestanol acetate, quingestron, and tigestol.

Preferred progestins are medroxyprogesterone, norethindrone and norethynodrel.

Exemplary polyphosphonates include polyphosphonates of the type disclosed in U.S. Patent 3,683,080. Preferred polyphosphonates are geminal diphosphonates  
20 (also referred to as bis-phosphonates). Tiludronate disodium is an especially preferred polyphosphonate. Ibandronic acid is an especially preferred polyphosphonate. Alendronate is an especially preferred polyphosphonate. Zoledronic acid is an especially preferred polyphosphonate. Other preferred polyphosphonates are 6-amino-1-hydroxy-hexylidene-bisphosphonic acid and 1-  
25 hydroxy-3(methylpentylamino)-propylidene-bisphosphonic acid. The polyphosphonates may be administered in the form of the acid, or of a soluble alkali metal salt or alkaline earth metal salt. Hydrolyzable esters of the polyphosphonates are likewise included. Specific examples include ethane-1-hydroxy 1,1-diphosphonic acid, methane diphosphonic acid, pentane-1-hydroxy-1,1-diphosphonic acid,  
30 methane dichloro diphosphonic acid, methane hydroxy diphosphonic acid, ethane-1-amino-1,1-diphosphonic acid, ethane-2-amino-1,1-diphosphonic acid, propane-3-amino-1-hydroxy-1,1-diphosphonic acid, propane-N,N-dimethyl-3-amino-1-hydroxy-1,1-diphosphonic acid, propane-3,3-dimethyl-3-amino-1-hydroxy-1,1-diphosphonic

acid, phenyl amino methane diphosphonic acid, N,N-dimethylamino methane diphosphonic acid, N(2-hydroxyethyl) amino methane diphosphonic acid, butane-4-amino-1-hydroxy-1,1-diphosphonic acid, pentane-5-amino-1-hydroxy-1,1-diphosphonic acid, hexane-6-amino-1-hydroxy-1,1-diphosphonic acid and  
5 pharmaceutically acceptable esters and salts thereof.

Any prostaglandin may be used in combination with an EP<sub>2</sub> selective receptor agonist in the present methods. The term prostaglandin refers to compounds that are analogs of the natural prostaglandins PGD<sub>1</sub>, PGD<sub>2</sub>, PGE<sub>2</sub>, PGE<sub>1</sub> and PGF<sub>2</sub>. These compounds bind to the prostaglandin receptors. Such binding is readily determined  
10 by those skilled in the art of standard assays (e.g., An S. et al., Cloning and Expression of the EP<sub>2</sub> Subtype of Human Receptors for Prostaglandin E<sub>2</sub>, Biochemical and Biophysical Research Communications, 1993, 197(1):263-270).

Prostaglandins are alicyclic compounds related to the basic compound prostanoic acid. The carbon atoms of the basic prostaglandin are numbered  
15 sequentially from the carboxylic carbon atom through the cyclopentyl ring to the terminal carbon atom on the adjacent side chain. Normally the adjacent side chains are in the trans orientation. The presence of an oxo group at C-9 of the cyclopentyl moiety is indicative of a prostaglandin within the E class while PGE<sub>2</sub> contains a trans unsaturated double bond at the C<sub>13</sub>-C<sub>14</sub> and a cis double bond at the C<sub>5</sub>-C<sub>6</sub> position.

20 A variety of prostaglandins are described and referenced below. However, other prostaglandins will be known to those skilled in the art. Exemplary prostaglandins are disclosed in U.S. patents 4,171,331 and 3,927,197.

Norrdin et al., The Role of Prostaglandins in Bone In Vivo, Prostaglandins Leukotriene Essential Fatty Acids 41, 139-150, 1990 is a review of bone anabolic  
25 prostaglandins.

Any prostaglandin agonist/antagonist may be used as the second compound in certain aspects of this invention. The term prostaglandin agonist/antagonist refers to compounds which bind to prostaglandin receptors (e.g., An S. et al., Cloning and Expression of the EP<sub>2</sub> Subtype of Human Receptors for Prostaglandin E<sub>2</sub>,  
30 Biochemical and Biophysical Research Communications, 1993, 197(1):263-270) and mimic the action of prostaglandin *in vivo*. Such actions are readily determined by those skilled in the art of standard assays. Eriksen E.F. et al., Bone Histomorphometry, Raven Press, New York, 1994, pages 1-74; Grier S.J. et. al., The

-44-

Use of Dual-Energy X-Ray Absorptiometry In Animals, Inv. Radiol., 1996, 31(1):50-62; Wahner H.W. and Fogelman I., The Evaluation of Osteoporosis: Dual Energy X-Ray Absorptiometry in Clinical Practice., Martin Dunitz Ltd., London 1994, pages 1-296. A variety of these compounds are described and referenced below. However,  
5 other prostaglandin agonists/antagonists will be known to those skilled in the art. Exemplary prostaglandin agonists/antagonists are disclosed as follows: U.S. patent 3,932,389 discloses 2-descarboxy-2-(tetrazol-5-yl)-11-desoxy-15-substituted-omega-pentanoicprostaglandins; U.S. patent 4,018,892 discloses 16-aryl-13,14-dihydro-PGE<sub>2</sub> p-biphenyl esters; U.S. patent 4,219,483 and 4,132,847 discloses 2,3,6-substituted-4-  
10 pyrones; U.S. patent 4,000,309 and 3,982,016 discloses 16-aryl-13,14-dihydro-PGE<sub>2</sub> p-biphenyl esters; U.S. patent 4,621,100 discloses substituted cyclopentanes; and U.S. patent 5,216,183 discloses cyclopentanones.

Sodium fluoride may be used as the second compound in certain aspects of this invention. The term sodium fluoride refers to sodium fluoride in all its forms (e.g.,  
15 slow release sodium fluoride, sustained release sodium fluoride). Sustained release sodium fluoride is disclosed in U.S. patent 4,904,478. The activity of sodium fluoride is readily determined by those skilled in the art of biological protocols (e.g., see Eriksen E.F. et al., Bone Histomorphometry, Raven Press, New York, 1994, pages 1-74; Grier S.J. et. al., The Use of Dual-Energy X-Ray Absorptiometry In Animals, Inv.  
20 Radiol., 1996, 31(1):50-62; Wahner H.W. and Fogelman I., The Evaluation of Osteoporosis: Dual Energy X-Ray Absorptiometry in Clinical Practice., Martin Dunitz Ltd., London 1994, pages 1-296).

Any parathyroid hormone (PTH) may be used as the second compound in certain aspects of this invention. The term parathyroid hormone refers to parathyroid  
25 hormone, fragments or metabolites thereof and structural analogs thereof which can stimulate bone formation and increase bone mass. Also included are parathyroid hormone related peptides and active fragments and analogs of parathyroid related peptides (see PCT publication no. WO 94/01460). A variety of these compounds are described and referenced below. However, other parathyroid hormones will be known  
30 to those skilled in the art. Exemplary parathyroid hormones are disclosed in the following references.

"Human Parathyroid Peptide Treatment of Vertebral Osteoporosis", Osteoporosis Int., 3, (Supp 1):199-203.

"PTH 1-34 Treatment of Osteoporosis with Added Hormone Replacement Therapy: Biochemical, Kinetic and Histological Responses" Osteoporosis Int. 1:162-170.

Any growth hormone or growth hormone secretagogue may be used as the  
 5 second compound in certain aspects of this invention. The term growth hormone secretagogue refers to a compound which stimulates the release of growth hormone or mimics the action of growth hormone (e.g., increases bone formation leading to increased bone mass). Such actions are readily determined by those skilled in the art of standard assays well known to those of skill in the art. A variety of these  
 10 compounds are disclosed in the following published PCT patent applications: WO 95/14666; WO 95/13069; WO 94/19367; WO 94/13696; and WO 95/34311. However, other growth hormones or growth hormone secretagogues will be known to those skilled in the art.

In particular a preferred growth hormone secretagogue is N-[1(R)-[1,2-  
 15 Dihydro-1-methanesulfonylspiro[3H-indole-3,4'-piperidin]-1'-yl)carbonyl]-2-(phenylmethyloxy)ethyl]-2-amino-2-methylpropanamide:MK-677.

Other preferred growth hormone secretagogues include  
 2-amino-N-(2-(3a-(R)-benzyl-2-methyl-3-oxo-2,3,3a,4,6,7-hexahydro-  
 pyrazolo-[4,3-c]pyridin-5-yl)-1-(R)-benzyloxymethyl-2-oxo-ethyl)-isobutyramide or its  
 20 L-tartaric acid salt;

2-amino-N-(1-(R)-benzyloxymethyl-2-(3a-(R)-(4-fluoro-benzyl)-2-methyl-3-  
 oxo-2,3,3a,4,6,7-hexahydro-pyrazolo[4,3-c]pyridin-5-yl)-2-oxo-ethyl)isobutyramide;  
 2-amino-N-(2-(3a-(R)-benzyl-3-oxo-2,3,3a,4,6,7-hexahydro-pyrazolo[4,3-  
 c]pyridin-5-yl)-1-(R)benzyloxymethyl-2-oxo-ethyl)isobutyramide; and  
 25 2-amino-N-(1-(2,4-difluoro-benzyloxymethyl)-2-oxo-2-(3-oxo-3a-pyridin-2-  
 ylmethyl-2-(2,2,2-trifluoro-ethyl)-2,3,3a,4,6,7-hexahydro-pyrazolo[4,3-c]pyridin-5-yl)-  
 ethyl)-2-methyl-propionamide.

The term "treating", "treat" or "treatment" as used herein includes preventative (e.g., prophylactic), palliative and curative treatment.

30 By "pharmaceutically acceptable" it is meant the carrier, diluent, excipients, and/or salts or prodrugs must be compatible with the other ingredients of the formulation, and not deleterious to the recipient thereof.

The term "prodrug" means a compound that is transformed *in vivo* to yield a compound of the present invention. The transformation may occur by various mechanisms, such as through hydrolysis in blood. A discussion of the use of prodrugs is provided by T. Higuchi and W. Stella, "Pro-drugs as Novel Delivery Systems," Vol. 14 of the A.C.S. Symposium Series, and in Bioreversible Carriers in Drug Design, ed. Edward B. Roche, American Pharmaceutical Association and Pergamon Press, 1987.

For example, when a compound of the present invention contains a carboxylic acid functional group, a prodrug can comprise an ester formed by the replacement of the hydrogen atom of the acid group with a group such as (C<sub>1</sub>-C<sub>8</sub>)alkyl, (C<sub>2</sub>-C<sub>12</sub>)alkanoyloxymethyl, 1-(alkanoyloxy)ethyl having from 4 to 9 carbon atoms, 1-methyl-1-(alkanoyloxy)-ethyl having from 5 to 10 carbon atoms, alkoxycarbonyloxymethyl having from 3 to 6 carbon atoms, 1-(alkoxycarbonyloxy)ethyl having from 4 to 7 carbon atoms, 1-methyl-1-(alkoxycarbonyloxy)ethyl having from 5 to 8 carbon atoms, N-(alkoxycarbonyl)aminomethyl having from 3 to 9 carbon atoms, 1-(N-(alkoxycarbonyl)amino)ethyl having from 4 to 10 carbon atoms, 3-phthalidyl, 4-crotonolactonyl, gamma-butyrolacton-4-yl, di-N,N-(C<sub>1</sub>-C<sub>2</sub>)alkylamino(C<sub>2</sub>-C<sub>3</sub>)alkyl (such as β-dimethylaminoethyl), carbamoyl-(C<sub>1</sub>-C<sub>2</sub>)alkyl, N,N-di(C<sub>1</sub>-C<sub>2</sub>)alkylcarbamoyl-(C<sub>1</sub>-C<sub>2</sub>)alkyl and piperidino-, pyrrolidino- or morpholino(C<sub>2</sub>-C<sub>3</sub>)alkyl.

Similarly, when a compound of the present invention comprises an alcohol functional group, a prodrug can be formed by the replacement of the hydrogen atom of the alcohol group with a group such as (C<sub>1</sub>-C<sub>6</sub>)alkanoyloxymethyl, 1-((C<sub>1</sub>-C<sub>6</sub>)alkanoyloxy)ethyl, 1-methyl-1-((C<sub>1</sub>-C<sub>6</sub>)alkanoyloxy)ethyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonyloxymethyl, N-(C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonylaminomethyl, succinoyl, (C<sub>1</sub>-C<sub>6</sub>)alkanoyl, α-amino(C<sub>1</sub>-C<sub>4</sub>)alkanoyl, arylacyl and α-aminoacyl, or α-aminoacyl-α-aminoacyl, where each α-aminoacyl group is independently selected from the naturally occurring L-amino acids, P(O)(OH)<sub>2</sub>, -P(O)(O(C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub> or glycosyl (the radical resulting from the removal of a hydroxyl group of the hemiacetal form of a carbohydrate).

When a compound of the present invention comprises an amine functional group, a prodrug can be formed by the replacement of a hydrogen atom in the

-47-

amine group with a group such as  $R^X$ -carbonyl,  $R^XO$ -carbonyl,  $NR^XR^X$ -carbonyl where  $R^X$  and  $R^X$  are each independently (C<sub>1</sub>-C<sub>10</sub>)alkyl, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, benzyl, or  $R^X$ -carbonyl is a natural  $\alpha$ -aminoacyl or natural  $\alpha$ -aminoacyl-natural  $\alpha$ -aminoacyl, -C(OH)C(O)OY<sup>X</sup> wherein Y<sup>X</sup> is H, (C<sub>1</sub>-C<sub>6</sub>)alkyl or benzyl), -C(OY<sup>X0</sup>) Y<sup>X1</sup> wherein Y<sup>X0</sup> is (C<sub>1</sub>-C<sub>4</sub>) alkyl and Y<sup>X1</sup> is (C<sub>1</sub>-C<sub>6</sub>)alkyl, carboxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, amino(C<sub>1</sub>-C<sub>4</sub>)alkyl or  
 5 mono-N- or di-N,N-(C<sub>1</sub>-C<sub>6</sub>)alkylaminoalkyl, -C(Y<sup>X2</sup>) Y<sup>X3</sup> wherein Y<sup>X2</sup> is H or methyl and Y<sup>X3</sup> is mono-N- or di-N,N-(C<sub>1</sub>-C<sub>6</sub>)alkylamino, morpholino, piperidin-1-yl or pyrrolidin-1-yl.

The expression "pharmaceutically acceptable salt" refers to nontoxic anionic  
 10 salts containing anions such as (but not limited to) chloride, bromide, iodide, sulfate, bisulfate, phosphate, acetate, maleate, fumarate, oxalate, lactate, tartrate, citrate, gluconate, methanesulfonate and 4-toluene-sulfonate. The expression also refers to nontoxic cationic salts such as (but not limited to) sodium, potassium, calcium, magnesium, ammonium or protonated benzathine (N,N'-dibenzylethylenediamine),  
 15 choline, ethanolamine, diethanolamine, ethylenediamine, meglamine (N-methyl-glucamine), benethamine (N-benzylphenethylamine), piperazine or tromethamine (2-amino-2-hydroxymethyl-1,3-propanediol).

As used herein, the expressions "reaction inert solvent" and "inert solvent" refers to a solvent that does not interact with starting materials, reagents,  
 20 intermediates or products in a manner which adversely affects the yield of the desired product.

It will be recognized that the compounds of this invention can exist in radiolabelled form, i.e., said compounds may contain one or more atoms containing an atomic mass or mass number different from the atomic mass or mass number  
 25 ordinarily found in nature. Radioisotopes of hydrogen, carbon, phosphorous, fluorine and chlorine include <sup>3</sup>H, <sup>14</sup>C, <sup>32</sup>P, <sup>35</sup>S, <sup>18</sup>F and <sup>36</sup>Cl, respectively. Compounds of this invention which contain those radioisotopes and/or other radioisotopes of other atoms are within the scope of this invention. Tritiated, i.e., <sup>3</sup>H, and carbon-14, i.e., <sup>14</sup>C, radioisotopes are particularly preferred for their ease of preparation and detectability.  
 30 Radiolabelled compounds of this invention can generally be prepared of methods well known to those skilled in the art. Conveniently, such radiolabelled compounds can be prepared by carrying out the procedures disclosed in the above Schemes



and/or in the Examples and Preparations below by substituting a readily available radiolabelled reagent for a non-radiolabelled reagent.

It will be recognized by persons of ordinary skill in the art that some of the compounds of this invention have at least one asymmetric carbon atom and therefore  
5 are enantiomers or diastereomers. Diastereomeric mixtures can be separated into their individual diastereomers on the basis of their physicochemical differences by methods known *per se* as, for example, by chromatography and/or fractional crystallization. Enantiomers can be separated by converting the enantiomeric mixture into a diastereomeric mixture by reaction with an appropriate optically active  
10 compound (e.g., alcohol), separating the diastereomers and converting (e.g., hydrolyzing, including both chemical hydrolysis methods and microbial lipase hydrolysis methods, e.g., enzyme catalyzed hydrolysis) the individual diastereomers to the corresponding pure enantiomers. All such isomers, including diastereomers, enantiomers and mixtures thereof are considered as part of this invention. Also, some  
15 of the compounds of this invention are atropisomers (e.g., substituted biaryls) and are considered as part of this invention.

In addition, when the compounds of this invention, including the compounds of Formula I, the anti-resorptive agents, bone anabolic agents, prostaglandin agonists/antagonists, parathyroid hormones, growth hormones and growth hormone  
20 secretagogues, form hydrates or solvates, they are also within the scope of the invention.

Administration of the compounds of this invention can be via any method that delivers a compound of this invention systemically and/or locally. These methods include oral routes, parenteral, intraduodenal routes, etc. Generally, the compounds  
25 of this invention are administered orally, but parenteral administration (e.g., intravenous, intramuscular, transdermal, subcutaneous, rectal or intramedullary) may be utilized, for example, where oral administration is inappropriate for the target or where the patient is unable to ingest the drug.

The compounds of this invention may also be applied locally to a site in or on  
30 a patient in a suitable carrier or diluent, optionally in combination with one or more of the anabolic agents or anti-resorptive agents described above.

The amount and timing of compounds administered will, of course, be dependent on the subject being treated, on the severity of the affliction, on the

manner of administration and on the judgment of the prescribing physician. Thus, because of patient to patient variability, the dosages given herein are guidelines and the physician may titrate doses of the drug to achieve the treatment that the physician considers appropriate for the patient. In considering the degree of treatment desired, the physician must balance a variety of factors such as age of the patient, presence of preexisting disease, as well as presence of other diseases (e.g., cardiovascular disease).

In general an effective dosage for the anabolic agents used in this invention described above is in the range of 0.001 to 100 mg/kg/day, preferably 0.01 to 50 mg/kg/day.

The following paragraphs provide preferred dosage ranges for various anti-resorptive agents.

In general, an effective dosage for an anti-resorptive agent is about 0.001 mg/kg/day to about 20 mg/kg/day.

In general, an effective dosage for progestins is about 0.1 to 10 mg per day; the preferred dose is about 0.25 to 5 mg per day.

In general, an effective dosage for polyphosphonates is determined by its potency as a bone resorption inhibiting agent of standard assays.

Ranges for the daily administration of some polyphosphonates are about 0.001 mg/kg/day to about 20 mg/kg/day.

In general an effective dosage for the treatment of this invention, for example the bone resorption treatment of this invention, for the estrogen agonists/antagonists of this invention is in the range of 0.01 to 200 mg/kg/day, preferably 0.5 to 100 mg/kg/day.

In particular, an effective dosage for raloxifene is in the range of 0.1 to 100 mg/kg/day, preferably 0.1 to 10 mg/kg/day.

In particular, an effective dosage for tamoxifen is in the range of 0.1 to 100 mg/kg/day, preferably 0.1 to 5 mg/kg/day.

In particular, an effective dosage for 2-(4-methoxy-phenyl)-3-[4-(2-piperidin-1-yl-ethoxy)-phenoxy]- benzo[b]thiophen-6-ol is 0.001 to 1 mg/kg/day.

In particular, an effective dosage for

*cis*-6-(4-fluoro-phenyl)-5-(4-(2-piperidin-1-yl-ethoxy)-phenyl)-5,6,7,8-tetrahydro-naphthalene-2-ol;

(-)-*cis*-6-phenyl-5-(4-(2-pyrrolidin-1-yl-ethoxy)-phenyl)-5,6,7,8-tetrahydro-naphthalene-2-ol;

*cis*-6-phenyl-5-(4-(2-pyrrolidin-1-yl-ethoxy)-phenyl)-5,6,7,8-tetrahydro-naphthalene-2-ol;

5        *cis*-1-(6'-pyrrolidinoethoxy-3'-pyridyl)-2-phenyl-6-hydroxy-1,2,3,4-tetrahydronaphthalene;

1-(4'-pyrrolidinoethoxyphenyl)-2-(4"-fluorophenyl)-6-hydroxy-1,2,3,4-tetrahydroisoquinoline;

10        *cis*-6-(4-hydroxyphenyl)-5-(4-(2-piperidin-1-yl-ethoxy)-phenyl)-5,6,7,8-tetrahydro-naphthalene-2-ol; or

1-(4'-pyrrolidinoethoxyphenyl)-2-phenyl-6-hydroxy-1,2,3,4-

tetrahydroisoquinoline is in the range of 0.0001 to 100 mg/kg/day, preferably 0.001 to 10 mg/kg/day.

15        The compounds of the present invention are generally administered in the form of a pharmaceutical composition comprising at least one of the compounds of this invention together with a pharmaceutically acceptable vehicle or diluent. Thus, the compounds of this invention can be administered individually or together in any conventional oral, parenteral, rectal or transdermal dosage form.

20        For oral administration a pharmaceutical composition can take the form of solutions, suspensions, tablets, pills, capsules, powders, and the like. Tablets containing various excipients such as sodium citrate, calcium carbonate and calcium phosphate are employed along with various disintegrants such as starch and preferably potato or tapioca starch and certain complex silicates, together with binding agents such as polyvinylpyrrolidone, sucrose, gelatin and acacia.

25        Additionally, lubricating agents such as magnesium stearate, sodium lauryl sulfate and talc are often very useful for tableting purposes. Solid compositions of a similar type are also employed as fillers in soft and hard-filled gelatin capsules; preferred materials in this connection also include lactose or milk sugar as well as high molecular weight polyethylene glycols. When aqueous suspensions and/or elixirs are  
30        desired for oral administration, the compounds of this invention can be combined with various sweetening agents, flavoring agents, coloring agents, emulsifying agents and/or suspending agents, as well as such diluents as water, ethanol, propylene glycol, glycerin and various like combinations thereof.

For purposes of parenteral administration, solutions in sesame or peanut oil or in aqueous propylene glycol can be employed, as well as sterile aqueous solutions of the corresponding water-soluble salts. Such aqueous solutions may be suitably buffered, if necessary, and the liquid diluent first rendered isotonic with sufficient saline or glucose. These aqueous solutions are especially suitable for intravenous, intramuscular, subcutaneous and intraperitoneal injection purposes. In this connection, the sterile aqueous media employed are all readily obtainable by standard techniques well-known to those skilled in the art.

For purposes of transdermal (e.g., topical) administration, dilute sterile, aqueous or partially aqueous solutions (usually in about 0.1% to 5% concentration), otherwise similar to the above parenteral solutions, are prepared.

Methods of preparing various pharmaceutical compositions with a certain amount of active ingredient are known, or will be apparent in light of this disclosure, to those skilled in this art. For examples of methods of preparing pharmaceutical compositions, see Remington's Pharmaceutical Sciences, Mack Publishing Company, Easton, Pa., 19th Edition (1995).

Pharmaceutical compositions of the invention may contain 0.1%-95% of the compound(s) of this invention, preferably 1%-70%. In any event, the composition or formulation to be administered will contain a quantity of a compound(s) of this invention in an amount effective to treat the disease/condition of the subject being treated.

The present invention can also be administered using an injectable, flowable composition that provides sustained release at the local site of the injection by forming a biodegradable solid or gel depot, matrix or implant. An example of such an administration system is an EP<sub>2</sub> selective receptor agonist compound in a slow-release biodegradable polymer based delivery system.

The polymer based delivery system contains EP<sub>2</sub> selective receptor agonist compound, and optionally any additional therapeutically useful compounds, dissolved or dispersed in biodegradable, thermoplastic polymer solution or dispersion in an organic solvent. Upon injection of the flowable composition, the organic solvent diffuses away from the injection site, causing the polymer to precipitate or gel; thereby entrapping the compound in a sustained-release depot. The compound is subsequently released by diffusion from, and erosion of, the polymeric matrix. The

-52-

polymeric matrix slowly erodes by hydrolysis and eventually disappears from the site of administration. The molecular weight and concentration of the polymer can control the in vivo release of the compound as well as the degradation rate of the matrix.

- 5           The polymer based delivery system provides sustained release of an EP<sub>2</sub> selective receptor agonist compound in vivo for a sustained period of time with minimum or reduced burst in a patient in need thereof. A large burst of compound would result in poor local toleration due to local effects of the compound (e.g., irritation) and would minimize the amount of compound available for efficacy. The advantages this administration method is that it minimizes or reduces the initial burst, but still delivers compound at efficacious levels for a sustained period of time upon a single local injection.

- 10           The polymer system is prepared by contacting the flowable composition with a gelation medium to coagulate or gel the composition into a solid, microporous polymeric matrix or a gel polymeric matrix. The flowable composition contains a thermoplastic polymer or copolymer in combination with a suitable solvent. The polymers or copolymers, which form the body of the matrix, are substantially insoluble, preferably essentially completely insoluble, in water and body fluids. The insolubility of the matrix body enables it to function as a single site for the controlled release of the EP<sub>2</sub> selective receptor agonist compound. The polymers or copolymers also are biocompatible and biodegradable and/or bioerodible within the body of an animal, e.g., mammal. The biodegradation enables the patient to metabolize the polymer matrix so that it can be excreted by the patient without the need for further surgery to remove it. Because the flowable composition and polymer system are biocompatible, the insertion process and the presence of the polymer system within the body do not cause substantial tissue irritation or necrosis at the implant site. The composition of the present invention is administered as a flowable composition directly into body tissues.

- 25           Suitable thermoplastic polymers for incorporation into the solid matrix of the controlled release polymer system are solids, pharmaceutically compatible and biodegradable by cellular action and/or by the action of body fluids. Examples of appropriate thermoplastic polymers include polyesters of diols and dicarboxylic acids or of hydroxycarboxylic acids, such as polylactides, polyglycolides and

copolymers thereof. More preferably the polymer is the copolymer, poly-lactic-co-glycolic acid (abbreviated PLGH), which upon hydrolysis, produces lactic and glycolic acid. The burst of release of this copolymer can be minimized further by the addition of polyethylene glycol (PEG) to form the PEG end-capped PLGH.

5 Preferred materials for use in the present invention are the polylactides, polyglycolides and copolymers thereof. These polymers can be used to advantage in the polymer system in part because they show excellent biocompatibility. They produce little, if any, tissue irritation, inflammation, necrosis or toxicity. In the presence of water, these polymers produce lactic and glycolic acid, respectively,  
10 which are readily metabolized by the body. The polylactides can also incorporate glycolide monomer to enhance the resulting polymer's degradation. These polymers can also be used because they effectively control the rate of release of the EP<sub>2</sub> selective receptor agonist compound from the polymer system, and because they result in the local retention of the EP<sub>2</sub> selective receptor agonist compound at  
15 the site of the site of administration.

The solubility or miscibility of a thermoplastic polymer in the organic solvent of the composition will vary according to factors such as crystallinity, hydrophilicity, capacity for hydrogen bonding and molecular weight of the polymer. Consequently, the molecular weight and the concentration of the polymer in the solvent are  
20 adjusted to achieve desired miscibility, as well as a desired release rate for the incorporated EP<sub>2</sub> selective receptor agonist compound.

The flowable composition of thermoplastic polymer, solvent and the EP<sub>2</sub> selective receptor agonist compound is a stable flowable substance. A homogenous solution of the EP<sub>2</sub> selective receptor agonist compound in an organic solvent  
25 preferably results. The thermoplastic polymer is substantially soluble in the organic solvent. Upon placement of the flowable composition into the body, the solvent will dissipate and the polymer will solidify or gel to form the polymer system having the EP<sub>2</sub> selective receptor agonist compound within a solid or gel polymeric matrix.

It has been discovered that the molecular weight of the polymer used  
30 distinctly affects the rate of release of the EP<sub>2</sub> selective receptor agonist compound and the rate of degradation of the polymer from the site as long as the flowable composition has been used as an intermediate.

-54-

For certain preferred polymers for use in the present invention, the molecular weight of the polymer or copolymer is adjusted to be within a range of about 0.2 to about 0.4 inherent viscosity (I.V. in deciliters/g) for effective sustained release of the bone growth promoting compound. The typical rate of release of the incorporated bone growth promoting compound occurs at an I.V. of about 0.2 (about 8,000 to about 16,000 molecular weight) or about 0.3 (about 23,000 to about 45,000 molecular weight) but can vary depending on the particular components of the composition. For most systems, it is preferred to adjust the molecular weight of the polymer to about 0.2 I.V. for an effective sustained release of the EP<sub>2</sub> selective receptor agonist compound. The unit of measure for the molecular weight is daltons.

For a poly(DL-lactide) or a lactide-co-glycolide polymer system, the desired molecular weight range is about 0.2 to about 0.4 I.V., with an I.V. of about 0.2 being most preferred. The molecular weight of a polymer can be varied by any of a variety of methods. The choice of method is typically determined by the type of polymer composition. The preferred polymers for use are commercially available.

Highly preferred thermoplastic polymers for use in the present invention are the following: PLGH copolymer with 1:1 ratio of lactic and glycolic acid with an inherent viscosity of about 0.2 dl/g (commercially available from Boehringer Ingelheim as Copolymer RESOMER® RG 502 H) (about 12,000 molecular weight); PLGH copolymer with 1:1 ratio of lactic and glycolic acid with an inherent viscosity of about 0.3 dl/g (commercially available from Boehringer Ingelheim as Copolymer RESOMER® RG 503 H) (about 37,000 molecular weight); PLGH copolymer with 1:1 ratio of lactic and glycolic acid with an inherent viscosity of about 0.4 dl/g (commercially available from Boehringer Ingelheim as Copolymer RESOMER® RG 504 H) (about 47,000 molecular weight); and polyethylene glycol (PEG) end-capped PLGH copolymer with 1:1 ratio of lactic and glycolic acid with an inherent viscosity of about 0.79 dl/g (commercially available from Boehringer Ingelheim as PLG-PEG) (about 52,000 molecular weight).

By appropriate choice of the polymer molecular weight and viscosity, the rate and extent of release of the EP<sub>2</sub> selective receptor agonist compound from the polymer system can be varied from very fast to very slow. For example, according to the present invention, the release rate of (3-(((4-*tert*-butyl-benzyl)-(pyridine-3-

sulfonyl)-amino)-methyl)-phenoxy)-acetic acid, sodium salt, can be slowed to produce substantially complete release of the compound within about seven days. With the use of a greater viscosity of polymer according to the present invention, the period of time can be increased to about fourteen days. The desired release  
5 rate of the EP<sub>2</sub> selective receptor agonist compound will depend on several factors, such as the species of animal being treated as well as the specific condition being treated.

The concentration of the polymer in the system can also be varied to adjust the release rate of the incorporated EP<sub>2</sub> selective receptor agonist compound. It has  
10 been discovered that the more dilute the polymer concentration, the more readily the EP<sub>2</sub> selective receptor agonist compound will be released. This effect can be used in combination with other methods to more effectively control the release of the incorporated EP<sub>2</sub> selective receptor agonist compound as desired. For example, by adjusting the concentration of the polymer and EP<sub>2</sub> selective receptor  
15 agonist compound, if desired, a wide range of release rates can be obtained

The solvents used in the thermoplastic compositions of the present invention are preferably pharmaceutically acceptable, biocompatible and will dissipate into body fluid *in situ* such that they may be classed as having a solubility in water ranging from highly soluble to insoluble. Preferably, they cause relatively little, if  
20 any, tissue irritation or necrosis at the site of the injection and implantation. Preferably, the solvent may have at least a minimal degree of water solubility. When the organic solvent is water insoluble or is minimally soluble in water, the solvent will slowly disperse from the flowable polymeric composition. The result will be an implant that during the course of its life may contain a varying amount of  
25 residual solvent. Especially preferably, the organic solvent has a moderate to high degree of water solubility so that it will facilely disperse from the polymeric composition into the body fluids. Most preferably, the solvent disperses rapidly from the polymeric composition so as to quickly form a solid implant. Concomitant with the dispersion of solvent, the thermoplastic polymer coagulates or gels into the solid  
30 polymer system. Preferably, as the thermoplastic polymer coagulates, the solvent dispersion causes pore formation within the polymer system. As a result, the flowable composition containing thermoplastic polymer, solvent and EP<sub>2</sub> selective receptor agonist compound will form a porous solid polymer system. Also, when the



-56-

solvent is slightly water soluble or is water insoluble, the solvent dispersion may result in the formation of a solid porous implant, or if some solvent remains with the implant, the result may be formation of a gel implant having few or no pores.

Suitable solvents include those liquid organic compounds meeting the foregoing  
5 criteria. The preferred solvent for use in the present invention is N-methyl-2-pyrrolidone (NMP) due, at least in part, to its solvating ability and its biocompatibility.

The solvents for the thermoplastic polymer flowable compositions are chosen for compatibility and appropriate solubility of the polymer and solvent. Lower  
10 molecular weight thermoplastic polymers will normally dissolve more readily in the solvents than high molecular weight polymers. As a result, the concentration of a thermoplastic polymer dissolved in the various solvents differs depending upon type of polymer and its molecular weight. Conversely, the higher molecular weight thermoplastic polymers will tend to coagulate, gel or solidify faster than the very low  
15 molecular weight thermoplastic polymers. Moreover, the higher molecular weight polymers tend to give higher solution viscosities than the low molecular weight materials. Thus, for advantageous injection efficiency, in addition to advantageous release rate, the molecular weight and the concentration of the polymer in the solvent are controlled.

20 Upon formation of the polymer system from the flowable composition, the EP<sub>2</sub> selective receptor agonist compound becomes incorporated into the polymer matrix. After insertion of the flowable composition to form in situ the polymer system, the EP<sub>2</sub> selective receptor agonist compound will be released from the matrix into the adjacent tissues or fluids by diffusion and polymer degradation mechanisms.

25 Manipulation of these mechanisms also can influence the release of the EP<sub>2</sub> selective receptor agonist compound into the surroundings at a controlled rate. For example, the polymer matrix can be formulated to degrade after an effective and/or substantial amount of the EP<sub>2</sub> selective receptor agonist compound is released from the matrix. Thus, the release of the EP<sub>2</sub> selective receptor agonist compound from  
30 the matrix can be varied by, for example, the solubility of the EP<sub>2</sub> selective receptor agonist compound in water, the distribution of the bone growth promoting compound within the matrix, or the size, shape, porosity, solubility and biodegradability of the polymer matrix, among other factors. The release of the EP<sub>2</sub> selective receptor

agonist compound from the matrix is controlled relative to its inherent rate by varying the polymer molecular weight to provide a desired duration and rate of release.

For example, a preferred dosage form of the EP<sub>2</sub> selective receptor agonist compound, (3-(((4-*tert*-butyl-benzyl)-(pyridine-3-sulfonyl)-amino)-methyl)-phenoxy)-acetic acid, is a lyophile of the sodium salt to be reconstituted with a solution of PLGH in NMP before administration. The dosage form, consisting of the lyophilized compound in one syringe (syringe A) and a solution of PLGH in NMP in a second syringe (syringe B), is known as the A/B reconstitution system. The contents of both syringes are mixed together immediately prior to dose delivery at or near site. After reconstitution, the contents are transferred into a graduated dosing syringe for delivery. The administered dosage forms will be a solution and will result in the dispersion of the compound with PLGH in NMP at desired strengths of, for example, 5 and 50 mgA/ml (mgA/ml refers to the free acid equivalent of the sodium salt form of the compound). The dosage form is a parenteral (e.g., subcutaneous, intramuscular or intramedullary) sustained release injection for local administration. This compound in a slow-release polymer matrix (depot injection) is designed for administration at or near a site, and is not intended for intravenous administration. To provide adequate shelf-life stability for the dosage form, a two-syringe system (A/B), as described above, may be used, preferably with the sodium salt form of the compound. A uniphase formulation, preferably with the free acid form of the compound, is a preferred alternative formulation. Based on the compound and polymer stability, sterile filtration of the compound and irradiation of the polymer solution may be preferred for manufacturing a stable sterile product. In one embodiment, the dosage form can be manufactured and shipped as separate aluminum pouches containing syringes filled with the lyophile form of the compound in one pouch and the polymer solution in the other pouch. Delivery containers, systems and methods for the lyophilization of the bone growth promoting compounds of the present invention to prepare pharmaceutical compositions and kits are described in published International patent application WO 01/73363.

#### EXAMPLE A

To obtain dosage form at strengths of 5 and 50 mgA/ml, the following combinations A) and B) of lyophile and polymer syringe, respectively, were used:

-58-

A) 5 mgA/ml (upon reconstitution) of (3-(((4-*tert*-butyl-benzyl)-(pyridine-3-sulfonyl)-amino)-methyl)-phenoxy)-acetic acid, sodium salt formulation:

Drug Syringe A contained 4 mgA of the sodium salt lyophile in 1.25 ml male syringe without graduations; and

5 Vehicle Syringe B contained 0.8 ml 50% RG502H/50%NMP solution in 1.25 ml female syringe without graduations.

B) 50 mgA/ml (upon reconstitution) of (3-(((4-*tert*-butyl-benzyl)-(pyridine-3-sulfonyl)-amino)-methyl)-phenoxy)-acetic acid, sodium salt formulation:

Drug Syringe A contained 40 mgA of the sodium salt lyophile in 1.25 ml male (fat) B-D syringe without graduations; and

Vehicle Syringe B contained 0.8 ml 50% RG502H/50% NMP solution in 1.25 ml female (thin) syringe without graduations.

MgA refers to free acid equivalent of the sodium salt form of the compound;

The percentages used in these examples are based on the weight of the indicated ingredients;

RG502H is a PLGH copolymer with 1:1 ratio of lactic and glycolic acid with inherent viscosity of 0.2 dl/gm, which is commercially available such as from Boehringer Ingelheim as Copolymer RESOMER® RG 502 H.

#### EXAMPLE 1

20 50% RG502H/50% NMP with 5 mgA/ml of sodium salt of (3-(((4-*tert*-butyl-benzyl)-(pyridine-3-sulfonyl)-amino)-methyl)-phenoxy)-acetic acid, mixed A/B (polymer solution autoclaved, compound lyophilized)

#### EXAMPLE 2

25 50% RG502H/50% NMP with 10 mgA/ml of sodium salt of (3-(((4-*tert*-butyl-benzyl)-(pyridine-3-sulfonyl)-amino)-methyl)-phenoxy)-acetic acid, mixed A/B (polymer solution irradiated, compound lyophilized)

#### EXAMPLE 3

30 50% RG502H/50% NMP with 50 mgA/ml of sodium salt of (3-(((4-*tert*-butyl-benzyl)-(pyridine-3-sulfonyl)-amino)-methyl)-phenoxy)-acetic acid, mixed A/B (polymer solution irradiated, compound lyophilized)

## EXAMPLE 4

47% RG502H/3% PLG-PEG/50% NMP with 50 mgA/ml of sodium salt of (3-(((4-*tert*-butyl-benzyl)-(pyridine-3-sulfonyl)-amino)-methyl)-phenoxy)-acetic acid, uniphase

## 5 EXAMPLE 5

47% RG503H/3% PLG-PEG/50% NMP with 50 mgA/ml of sodium salt of (3-(((4-*tert*-butyl-benzyl)-(pyridine-3-sulfonyl)-amino)-methyl)-phenoxy)-acetic acid, uniphase

## EXAMPLE 6

10 45% RG504H/55% NMP with 50 mgA/ml of sodium salt of (3-(((4-*tert*-butyl-benzyl)-(pyridine-3-sulfonyl)-amino)-methyl)-phenoxy)-acetic acid, uniphase

## EXAMPLE 7

37% RG503H/63% NMP with 50 mgA/ml of sodium salt of (3-(((4-*tert*-butyl-benzyl)-(pyridine-3-sulfonyl)-amino)-methyl)-phenoxy)-acetic acid, mixed A/B  
15 (polymer solution autoclaved, compound lyophilized)

## EXAMPLE 8

37% RG503H/63% NMP with 50 mgA/ml of sodium salt of (3-(((4-*tert*-butyl-benzyl)-(pyridine-3-sulfonyl)-amino)-methyl)-phenoxy)-acetic acid, mixed A/B  
(polymer solution irradiated, compound lyophilized)

## 20 EXAMPLE 9

50% RG502H/50% NMP with 5 mgA/ml of (3-(((4-*tert*-butyl-benzyl)-(pyridine-3-sulfonyl)-amino)-methyl)-phenoxy)-acetic acid, uniphase.

Further exemplification of the polymer matrix delivery system described above can be found in U.S. provisional patent application No 60/337,255, filed November  
25 30, 2001.

In a preferred administration system, syringe A contains the lyophile of the sodium salt of (3-(((4-*tert*-butyl-benzyl)-(pyridine-3-sulfonyl)-amino)-methyl)-phenoxy)-acetic acid which is made to result in 4 mg per syringe or 40 mg per syringe. Syringe B contains Resomer 502H, 50:50 Poly(D,L lactide-co-glycolide),  
30 (50,50 PLGH) and N-methyl-2-pyrrolidone (NMP).

The dose using the above described A/B syringe system can vary widely and is determined by the disease being treated among other factors. A preferred dose range includes 0.5mgA up to about 100mgA. In syringes containing 50mgA/ml, preferred

-60-

doses include 0.1ml, 0.2ml, 0.6ml and 2ml. In syringes containing 5mgA/ml, preferred doses include 0.1ml, 0.2ml, 0.3ml, 0.6 and 2ml.

Other methods of administration of an EP<sub>2</sub> selective receptor agonist include local administration by injection to a particular site or delivery by a catheter to a site.

- 5 Additional examples can be found in U.S. provisional patent application number 60/335,156, filed November 30, 2001.

### GENERAL EXPERIMENTAL PROCEDURES

- 10 In general the compounds of this invention can be made by processes which include processes known in the chemical arts, particularly in light of the description contained herein. Certain processes for the manufacture of the compounds of this invention are provided as further features of the invention and are illustrated by the following reaction schemes. Other processes are described in the experimental section.

- 15 Some substituents (e.g., carboxyl) may best be prepared through conversion of another functional group (e.g., carboxyl substituents may be prepared through conversion of, e.g., hydroxyl or carboxaldehyde) at a point later in the synthetic sequence.

- 20 Compounds of Formula I wherein B is nitrogen may be prepared using methods described in SCHEMES 1-5. These methods include (a) sequential alkylation of a sulfonamide or amide with two appropriate alkylating agents, generally alkyl halides or alkyl sulfonates; (b) alkylation of a sulfonamide or amide with an alkyl halide or alkyl sulfonate; or (c) reductive amination of an aldehyde followed by reaction with an acylating agent such as an acyl chloride, a chloroformate, an isocyanate or a chlorocarbonyl amide; or a sulfonylating agent such as a sulfonyl chloride. When performing sequential alkylation, one of the alkylating agents will contain a Q-Z portion, where the Z portion is suitably protected if necessary, and the other alkylating agent will contain a K-M portion, where any functional groups requiring protection are suitably protected. The order of the alkylation, i.e., whether
- 25 the alkylating agent containing the Q-Z portion is added first or second, will depend upon the reactivity of the electrophilic side chain. When performing a reductive amination, the Q-Z portion may be attached to either the amine reagent or the aldehyde reagent depending upon the ease of preparation of the reagent and the
- 30

-61-

reactivity of the reagents in the reductive amination reaction. The reductive amination is followed by acylation or sulfonylation with an appropriate acylating agent or sulfonyl chloride and, if desired the product is hydrolysed. The starting materials, including amines, aldehydes and alkylating agents, are prepared using methods well known to those skilled in the art. Certain preferred methods for their preparation are described herein.


For example, compounds of Formula I wherein B is N are prepared of the methods set forth in SCHEMES 1 and 2 below. In general, the sequences involve sequential alkylation of an appropriate sulfonamide of Formula 1 or amide of Formula 1 with two appropriate alkyl halides or alkyl sulfonates. SCHEMES 1 and 2 differ only in the order of addition of the two alkylating agents. The alkylation order is typically chosen depending on the reactivity of the electrophilic side-chain. It is generally preferable to react the less reactive electrophilic side chain first. This reduces the amount of dialkylation which occurs in that first alkylation step, thereby resulting in a greater yield of monoalkylated material to be carried on to the next alkylation. In SCHEMES 1 and 2, one of the alkylating agents contains a carboxylic acid or a carboxylic acid isostere, suitably protected with an appropriate protecting group, if necessary. Further, in SCHEMES 1 and 2, the carboxylic acid precursor of Formula 3 is a carboxylic acid ester where R is a suitable carboxylic acid protecting group. Generally, the protecting group is either a straight chain lower alkyl, preferably methyl or ethyl, or a *tert*-butyl or phenyl group. Other acid isosteres can be employed by appropriately modifying SCHEMES 1 and 2 of methods well known to those skilled in the art (e.g., see SCHEME 6 which sets forth the preparation of a tetrazole). Typical alkylating agents are primary, secondary, benzylic or allylic halides and sulfonates and are preferably alkyl bromides or alkyl iodides.

The Formula 1 sulfonamide or amide is converted to its anion with a strong base such as sodium hydride, lithium diisopropylamide, lithium bis(trimethylsilyl)amide, potassium bis(trimethylsilyl)amide, potassium *tert*-butoxide, etc. in an aprotic solvent such as dimethylformamide, tetrahydrofuran or N,N-dimethylformamide/benzene at a temperature of about -78°C to about 100°C. The resulting anion is alkylated with an appropriate alkyl halide of Formula 2 or 3 or an appropriate alkyl sulfonate of Formula 2 or 3, wherein X' is the halide or sulfonate portion of the alkylating agent, at a temperature of about 0°C to about 100°C to yield

-62-


- the corresponding mono-alkylated compound of Formula 4 or 5. In some cases, varying amounts of a side-product resulting from dialkylation of the amide or sulfonamide are obtained and can be removed using chromatographic techniques, preferably by flash chromatography (W.C. Still, M. Kahn, A. Mitra, J. Org. Chem. 43, 2923, 1978). After the first alkylation is complete, the compound of Formula 4 or 5 is converted to an anion using a suitable base such as sodium hydride, lithium bis(trimethylsilyl)amide, lithium diisopropylamide, potassium bis(trimethylsilyl)amide, potassium tert-butoxide, or potassium carbonate in an aprotic solvent such as N,N-dimethylformamide, tetrahydrofuran,
- 10 N,N- dimethylformamide/benzene, or acetone at a temperature of about -78°C to about 100°C. Alkylation of the anion with an appropriate second alkyl halide of Formula 3 or 2 or alkyl sulfonate of Formula 3 or 2 provides the corresponding dialkylated compound of Formula 6. When R is methyl or ethyl, the ester of Formula 6 is hydrolyzed to the corresponding carboxylic acid of Formula I with a dilute
- 15 aqueous basic solution. This hydrolysis is preferably carried out using sodium or potassium hydroxide in aqueous methanol or ethanol, lithium hydroxide in aqueous alcoholic solvent or aqueous tetrahydrofuran at a temperature of about 0°C to about 80°C. Alternatively, the hydrolysis may be carried out by using methods well known to those skilled in the art, for example, methods described in "Protecting Groups in
- 20 Organic Synthesis," Second Edition, T.W. Greene and P.G.M. Wuts, John Wiley and Sons, Inc., 1991.

SCHEME 1



-63-

## SCHEME 2



Compounds of Formula I wherein B is N are also prepared from amines as set forth in SCHEMES 3-4. Generally, the appropriate amine starting materials of

5 Formulas 9 and 10 are commercially obtained or can be prepared using methods well known to those skilled in the art (see "The Chemistry of Amino, Nitroso and Nitro Compounds and their Derivatives," Ed. S. Patai, J. Wiley, New York, 1982). For example, the amine starting materials are prepared from the corresponding nitriles of

10 Formulas 7 or 8. Said nitriles are available from commercial sources or can be prepared using methods well known to those skilled in the art (see Rappaport, "The Chemistry of the Cyano Group," Interscience, New York, 1970 or Patai and Rappaport, "The Chemistry of Functional Groups," pt. 2, Wiley, New York, 1983). The nitrile of Formula 7 or 8 is reduced with a reducing agent such as borane-

15 tetrahydrofuran complex, borane-methyl sulfide complex or lithium aluminum hydride in an aprotic solvent such as tetrahydrofuran or diethyl ether at a temperature of about -78°C to about 60°C. Alternatively, the nitrile is hydrogenated under a hydrogen atmosphere typically at 0 to 50 psi in the presence of Raney nickel or a platinum or palladium catalyst in a protic solvent such as methanol or ethanol at a temperature of about 0°C to about 50°C. It may be desired to add an equivalent of an

20 acid, such as hydrogen chloride, to accomplish the reduction. The amine of Formula 9 or 10 thus obtained is converted to the sulfonamide of Formula 11 or 12 by sulfonylation with a sulfonyl chloride or said amine is converted to an amide of Formula 11 or 12 by acylation with an appropriate acyl chloride. Both the sulfonylation reactions and the acylation reactions are generally carried out in the



-64-


presence of a weak base such as triethylamine, pyridine, or 4-methylmorpholine in an aprotic solvent such as methylene chloride or diethyl ether at a temperature of about -20°C to about 50°C. Alternatively, coupling of amines of Formulas 9 or 10 with carboxylic acids are conveniently carried out in an inert solvent such as dichloromethane or N,N-dimethylformamide by a coupling reagent such as 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (EDC) or 1,3-dicyclohexylcarbodiimide (DCC) in the presence of 1-hydroxybenzotriazole hydrate (HOBT) to generate compounds of Formulas 11 or 12. Where the amine is present as the hydrochloride or other salt, it is preferable to add one equivalent of a suitable base such as triethylamine to the reaction mixture. Alternatively, the coupling can be effected with a coupling reagent such as benzotriazol-1-yloxy-tris(dimethylamino)phosphonium hexafluorophosphate (BOP) in an inert solvent such as methanol. Such coupling reactions are generally conducted at temperatures of about -30°C to about 80°C, preferably 0°C to about 25°C. For a discussion of other conditions used for coupling peptides see Houben-Weyl, Vol. XV, part II, E. Wunsch, Ed., George Thieme Verlag, 1974, Stuttgart. Alkylation and if desired, deprotection, of the Formula 11 or 12 compound as described in SCHEMES 1 and 2 affords the corresponding acid Formula 13 and 14 compound. The compounds of Formulas 11 and 12 are alkylated in a manner analogous to the alkylation of the compounds of Formulas 1, 4 and 5 of SCHEMES 1 and 2 hereinabove. The alkylated products are deprotected, if necessary, to afford the compounds of Formulas 13 and 14.

The amines of Formulas 9 and 10 are also prepared via reduction of an appropriate amide of Formulas 15 and 16. This reduction is achieved using reagents such as a borane-tetrahydrofuran complex, a borane-methyl sulfide complex, or diisobutylaluminum hydride in an aprotic solvent such as tetrahydrofuran or diethyl ether at a temperature of about -78°C to about 60°C.

The amines of Formulas 9 and 10 are also obtained from the corresponding nitro precursors by reduction of the nitro group using reducing reagents such as zinc/HCl, hydrogenation in the presence of Raney nickel, palladium, or platinum catalysts, and other reagents as described by P.N. Rylander in "Hydrogenation Methods," Academic Press, New York, 1985.


-65-

## SCHEME 3



-66-

## SCHEME 4



Amines and alkylating agents useful for the above syntheses are described and prepared as set forth in the section entitled PREPARATIONS below.

- 5 Alternatively, the compounds of Formula I wherein B is N are prepared by reductive amination of an aldehyde containing the appropriate suitably protected acidic functionality with an amine. This sequence is set forth in SCHEME 5. Alternatively, the amine may contain the appropriate suitably protected acidic functionality.


- 10 The reductive amination is typically carried out at a pH of between 6 and 8, using a reducing agent such as sodium cyanoborohydride or sodium triacetoxyborohydride. The reaction is normally performed in a protic solvent such as methanol or ethanol at temperatures of about -78°C to about 40°C. (e.g., see A. Abdel-Magid, C. Maryanoff, K. Carson, *Tetrahedron Lett.* **39**, 31, 5595-5598, 1990.)
- 15 The reductive amination reaction may also be carried out using titanium isopropoxide and sodium cyanoborohydride (R.J. Mattson et al, *J. Org. Chem.* 1990, 55, 2552-4)

-67-

or by preformation of the imine under dehydrating conditions followed by reduction. The resulting amine of Formulas 42 and 42A, is transformed to the desired amide or sulfonamide by coupling with an acid chloride, sulfonyl chloride, or carboxylic acid as set forth in SCHEMES 3 and 4. If desired, the amine intermediate of Formulas 42 or 42A may be converted to a urethane by treatment with a chloroformate or to a tetrasubstituted urea by treatment with a chlorocarbonyl amide. These reactions are performed in the presence of a weak base such as triethylamine, pyridine, or 4-methylmorpholine in an aprotic solvent such as methylene chloride or diethyl ether at a temperature of about -20°C to about 50°C. Conversion of the amine of Formulas 42 or 42A to a trisubstituted urea is accomplished by treatment with an isocyanate in an aprotic solvent such as methylene chloride or diethyl ether at temperatures ranging between -20°C and 50°C (for example, see SCHEME 5A). In cases where the amine is present as the hydrochloride salt, it is preferable to add an equivalent of a suitable base such as triethylamine to the reaction. If desired, hydrolysis of the resulting sulfonamide or amide provides the corresponding acid.


-68-

## SCHEME 5



-69-

## SCHEME 5A



Aldehydes useful in the above SCHEME 5 are described and prepared as set forth in the section entitled PREPARATIONS below.


- 5 Compounds of Formula I where B is N and Z is tetrazolyl are prepared as set forth in SCHEME 6. A sulfonamide or amide of Formula 4 is alkylated with the appropriate alkyl halide or sulfonate (wherein X' is halide or sulfonate), preferably a primary, secondary, benzylic, or allylic alkyl bromide, iodide, or sulfonate, which contains a nitrile to provide a nitrile of Formula 59. This alkylation is achieved by
- 10 treatment of the sulfonamide or amide of Formula 59 with a base such as sodium hydride, lithium bis(trimethylsilyl)amide, potassium bis(trimethylsilyl)amide, potassium tert-butoxide, or potassium carbonate in an aprotic solvent such as dimethylformamide, dimethylformamide/benzene, or acetone followed by reaction of the resulting anion with a suitable alkylating agent. Alkylation occurs at a temperature
- 15 of about -78°C to about 100°C. A preferred method for converting the resulting nitrile of Formula 59 to the tetrazole of Formula 60 is treatment of the alkylated nitrile with dibutyltin oxide and trimethylsilylazide, in refluxing toluene (S.J. Wittenberger and

-70-

B.G. Donner, J. Org. Chem. 1993, 58, 4139-4141, 1993). For a review of alternative preparations of tetrazoles see R.N. Butler, Tetrazoles, In Comprehensive Heterocyclic Chemistry; Potts, K.T. Ed.; Pergamon Press: Oxford, 1984, Vol. 5, pp 791-838.

5

## SCHEME 6




Alternatively, certain compounds of Formula I wherein B is N are prepared as set forth in SCHEME 7. Thus, esters of Formula 46 are prepared using the procedures described above in SCHEMES 1 and 2. Subsequent Heck coupling of this intermediate to an arylhalide (preferably an aryl bromide or aryl iodide), an aryl triflate, or a ring system which contains a vinyl bromide, iodide, or triflate is accomplished with a palladium catalyst, such as palladium acetate or tetrakis(triphenylphosphine)palladium(0) in the presence of a trialkylamine, such as triethylamine. In some cases, an additive such as a triarylphosphine or triarylsarsene may be added to the reaction. The reaction is typically performed in an aprotic solvent such as dimethylformamide or acetonitrile at a temperature of about 0°C to about 150°C (see R.F. Heck in Comp. Org. Syn., Vol. 4, Ch. 4.3, p. 833 or Daves and Hallberg, Chem. Rev. 1989, 89, 1433). If desired, the compound of Formula 47 can

-71-

- be hydrolyzed to the corresponding acid. Alternatively, the compound of Formula 47 can be hydrogenated and, if desired, further hydrolyzed to the corresponding acid of Formula 49. Hydrogenation is preferably achieved under a hydrogen atmosphere typically at 0 to 50 psi in the presence of a palladium or platinum catalyst in an alcoholic solvent such as ethanol or methanol at a temperature of about 0°C to about 50°C. In cases where M represents a partially saturated ring system, hydrogenation will generate a fully saturated ring system.

## SCHEME 7



10


- Alternatively, certain compounds of Formula I wherein B is N are prepared as described in SCHEME 8. Compounds of Formula 51 are prepared as described in SCHEMES 1 and 2 by alkylation of compounds of Formula 5 with an electrophile of Formula 2 which contains the appropriate functionality on the M ring. At least one of the substituents on the M ring must be suitable for subsequent conversion to an aldehyde. For example, electrophiles of Formula 2 containing a protected alcohol on the M ring may be alkylated and then deprotected and oxidized to the aldehyde,



-72-

using reagents well known to those skilled in the art, to generate compounds of Formula 51. An alternative method is to alkylate with an electrophile of Formula 2 where M contains a vinyl group. After alkylation, oxidative cleavage of the double bond provides the desired aldehyde of Formula 51. The oxidative cleavage is accomplished by transforming the double bond to the 1,2-diol with catalytic osmium tetroxide and N-methylmorpholine followed by oxidative cleavage to the aldehyde using sodium periodate. Alternatively, oxidative cleavage via ozonolysis followed by reduction using reagents such as methyl sulfide, triphenylphosphine, zinc/acetic acid, or thiourea, generates the desired aldehyde of Formula 51. Addition of LMetal where LMetal is any organometallic reagent such as an organolithium or a Grignard reagent in an aprotic solvent such as diethyl ether or tetrahydrofuran at a temperature of about -78°C to about 80°C, followed by hydrolysis of the ester as described above, provides the desired compound of Formula 50.

## SCHEME 8




Alternatively, certain compounds of Formula I wherein B is N are prepared as described in SCHEME 9. The appropriate sulfonamide or amide of Formula 5 is alkylated using the conditions described in SCHEMES 1 and 2. The alkylating agent is an electrophile which contains an aromatic bromide or iodide or a ring system which contains a vinyl bromide or iodide ( $\text{Ar}^1$ ) to provide compounds of Formula 53. Suzuki-type coupling of the compound of Formula 53 thus obtained with an aryl boronic acid ( $\text{Ar}^2$ ) provides Formula 53a compounds. For a review of the Suzuki reaction see A.R. Martin and Y. Yang in *Acta Chem. Scand.* 1993, 47, 221. The

-73-

- coupling reaction is achieved using about two equivalents of a base, such as sodium carbonate, potassium carbonate, sodium hydroxide, thallium hydroxide or potassium phosphate, in the presence of a palladium catalyst, such as tetrakis(triphenylphosphine)palladium(0), palladium acetate, palladium chloride,
- 5 tris(dibenzylideneacetone)dipalladium(0) or [1,4-bis(diphenylphosphine)butane]palladium(0). The reaction may be run in an aqueous alcoholic solvent such as methanol or ethanol; or in other aqueous solvents such as aqueous tetrahydrofuran, aqueous acetone, aqueous glycol dimethyl ether, or aqueous benzene at temperatures ranging from about 0°C to about 120°C. When
- 10 Ar<sup>1</sup> is a partially saturated ring, reduction of the ring to provide a saturated ring system may, if desired, be performed at this point. This transformation is achieved by hydrogenating the partially saturated ring in the presence of a catalyst such as palladium or platinum in an alcoholic solvent (ethanol or methanol) and/or ethyl acetate. Ester hydrolysis of compounds of Formulas 53 or 53a, if desired, provides
- 15 the corresponding acid. The resulting acids may contain functional groups on either of the ring systems (Ar<sup>1</sup> or Ar<sup>2</sup>) which can be modified using methods well known to those skilled in the art. Examples of such modifications are shown in SCHEME 10.


SCHEME 9



-74-

Compounds of Formula 54 which contain an aldehyde functional group are prepared using methods described in SCHEMES 8 and 9. Of SCHEME 10, treatment of a compound of Formula 54 with an appropriate organometallic reagent (LMetal), such as an organolithium or Grignard reagent, in an aprotic solvent such as diethyl ether or tetrahydrofuran at a temperature of about -78°C to about 80°C, followed by hydrolysis of the ester, provides compounds of Formula 56. Alternatively, reduction of the aldehyde followed by hydrolysis provides Formula 55 compounds.

SCHEME 10



10

Alternatively, certain compounds of Formula I wherein B is N are prepared as described in SCHEME 11. The starting alcohols of Formula 58 are prepared of methods well known to persons skilled in the art, for example, by using methods described in SCHEMES 1 and 2. It will be recognized by a person of ordinary skill in the art that protecting groups may be required in the synthesis of certain of these alcohols. Intermediate 58 is coupled with a variety of aryl alcohols (M is as defined above) using Mitsunobu coupling conditions (for a review see O. Mitsunobu, Synthesis, 1, 1981). Typically the coupling is achieved by addition of a coupling agent


15

-75-

such as triphenylphosphine and diethyl azodicarboxylate or diisopropyl azodicarboxylate in an inert solvent such as methylene chloride or tetrahydrofuran at a temperature of about 0°C to about 80°C. If desired, subsequent hydrolysis yields the corresponding acid.

5


SCHEME 11



Alternatively, certain compounds of Formula I wherein B is N are prepared as described in SCHEME 12. A compound of Formula 102 is added to a compound of Formula 105 (wherein X is as defined above for the compound of Formula I) in the presence of a Lewis acid such as titanium tetrachloride or a mineral acid such as hydrochloric acid. If desired the ester of Formula 106 can be converted to the corresponding acid by hydrolysis or deprotection.

15

SCHEME 12



Alternatively, certain compounds of Formula I wherein B is N are prepared as described in SCHEME 13. Chloromethyl compounds of Formula 104 are treated with the appropriate substituted aromatic ring system, M, such as 4-ethoxybenzene or thiophene in the presence of a Lewis acid such as titanium tetrachloride or a mineral acid such as hydrochloric acid in an aprotic solvent such as chloroform at a

20


-76-

temperature of about 0°C to about 80°C to yield compounds of Formula 107 which may subsequently be hydrolyzed or deprotected as described above to yield the corresponding carboxylic acids. Alternatively, chloromethyl compounds of Formula 104 can be treated with a Lewis acid such as titanium tetrachloride and an

5 appropriately substituted vinyl silane in an aprotic solvent such as methylene chloride at a temperature of about -50°C to about 50°C to give compounds of Formula 108. If desired, the compounds of Formula 108 may subsequently be hydrolyzed or deprotected as described above to yield the corresponding acid. If desired, reduction of the double bond can be accomplished using conditions described in SCHEME 7.

10

SCHEME 13




Alternatively, certain compounds of Formula I wherein B is N are prepared as

15 described in SCHEME 14. Chloromethyl compounds of Formula 104 are treated with a Lewis acid such as titanium tetrachloride and an appropriately substituted allyl silane in an aprotic solvent such as chloroform at a temperature of about 0°C to about 80°C to give compounds of Formula 109 which may subsequently be hydrolyzed or deprotected as described above to afford the corresponding carboxylic acids.


-77-

## SCHEME 14



- 5 Alternatively, certain compounds of Formula I wherein B is N are prepared as described in SCHEME 15. Chloromethyl compounds of Formula 104 are treated with a sulfinic acid of Formula 111 in the presence of a base such as triethylamine in an aprotic solvent such as chloroform at a temperature of about -30°C to about 50°C to give compounds of Formula 112 which may subsequently be hydrolyzed or
- 10 deprotected as described above to yield the corresponding acid.

## SCHEME 15




- Formula I compounds wherein B is C(H) and Q, G, M and K are as described above in the Summary of the Invention can be prepared of SCHEME 16. Formula 113 beta-ketoesters are alkylated sequentially with Formula 114 compounds to form
- 15

-78-

- Formula 115 compounds followed by alkylation with Formula 116 compounds to give Formula 117 compounds (J. Med. Chem. 26, 1993, p335-41). Alkylations can be carried out in a suitable solvent such as DMF, THF, ether, or benzene using an appropriate base such as sodium hydride, LDA, or potassium carbonate at a temperature of about -78°C to about 80°C. The resulting Formula 117 disubstituted keto esters are hydrolyzed and decarboxylated to give the corresponding Formula 118 compound by using an aqueous base such as sodium hydroxide to hydrolyze the ester, followed by an acidic quench such as with aqueous hydrochloric acid to effect decarboxylation.

10

SCHEME 16




-79-

- Alternatively, Formula I compounds wherein B is C(H) and G, Q, M and K are as described above in the Summary of the Invention may be prepared of SCHEME 17. Sequential alkylation of a malonate derivative of Formula 119 provides the Formula 121 dialkylated compound. Deprotection of the ester group by treatment with
- 5 a strong acid such as TFA or HCl in ethanol at a temperature of about -20°C to about 50°C leads to the Formula 122 decarboxylated product. Conversion of the acid to an acid chloride using thionyl chloride or oxalyl chloride in an aprotic solvent at a temperature of about -78°C to about 50°C or to a Weinreb amide using methoxymethyl amine in the presence of a suitable coupling agent such as DCC or
- 10 DEC in an aprotic solvent at a temperature of about -30°C to about 50°C provides Formula 123 compounds. Formula 123 compounds are suitable substrates for addition of various organometallic species, e.g., Grignard reagents and organo-cadmium reagents which, after hydrolysis of the terminal ester, provide the keto-acid compounds of Formula 118.
- 15 Alternatively, Formula 118 compounds can be prepared using methods described previously in Schemes 7-11 where one or both of the side chains are further functionalized after attachment to the alkanoyl fragment.



-80-

SCHEME 17




## **PREPARATIONS**

### **Amines, Amides and Sulfonamides**


- Certain amides or sulfonamides described by Formulas 21, 22, and 23
- 5 wherein W and Z are as described above in the Summary of the Invention and X and M are aromatic or saturated ring systems are prepared as set forth in SCHEME 18. Alkynyl amides or sulfonamides of Formulas 25, 26 and 27 are prepared by coupling an alkynyl sulfonamide or alkynyl amide of Formula 24 to an aromatic or vinyl halide, preferably an aromatic or vinyl bromide or iodide wherein W and Z are as defined
- 10 above and where X and M represent an aromatic ring or a partially saturated ring system. The coupling is typically accomplished in the presence of copper iodide, a palladium catalyst, such as palladium chloride, bis(triphenylphosphine)palladium dichloride, or tetrakis(triphenylphosphine)palladium(0), and an amine such as triethylamine, diisopropylamine, or butylamine in an aprotic solvent such as
- 15 acetonitrile at a temperature of about 0°C to about 100°C. The alkynes of Formulas 25, 26 and 27 are converted to the corresponding alkanes of Formulas 21, 22 or 23 via hydrogenation in the presence of a palladium or platinum catalyst in a solvent such as methanol, ethanol, and/or ethyl acetate at a temperature of about 0°C to about 50°C. In the case where M represents a partially saturated ring system,
- 20 hydrogenation will convert M to a fully saturated ring system. Alternatively, the alkynes are converted to cis-alkenes using the Lindlar catalyst (Pd-CaCO<sub>3</sub>-PbO) or other suitable catalyst. Alkylation and deprotection as described in SCHEMES 1 and 2 affords the corresponding compounds of Formula I.

-82-

SCHEME 18




SCHEME 19



-83-

- Compounds of Formula 33 are prepared from a suitable amine of Formula 32 (e.g., methoxyarylalkylamine). Amines of Formula 32 are commercially available or are prepared by methods well known to those skilled in the art (for example, see
- 5 SCHEME 4). Amines of Formula 32 are converted to sulfonamides or amides of Formula 31 using methods, for example, described in SCHEME 3 and 4. The resulting aromatic methyl ether of Formula 31 is deprotected with reagents such as boron tribromide, pyridinium hydrochloride, hydrogen bromide/acetic acid, or other reagents as described in *Protecting Groups in Organic Synthesis, Second Edition,*
- 10 T.W. Greene and P.G.M. Wuts, John Wiley and Sons, Inc., 1991. Alkylation with a bromoalkylester using a mild base such as potassium carbonate in an aprotic solvent such as dimethylformamide or acetone at a temperature of about 0°C to about 100°C generates an amide or sulfonamide of Formula 33.

## SCHEME 20



## ALKYLATING AGENTS

Numerous methods exist for the synthesis of the desired alkylating agents used in the above procedures and are known to those skilled in the art (see "The


-84-

Chemistry of the Carbon-Halogen Bond," Ed. S. Patai, J. Wiley, New York, 1973 and "The Chemistry of Halides, Pseudo-Halides, and Azides," Eds. S. Patai and Z. Rappaport, J. Wiley, New York, 1983). Some examples are shown in SCHEMES 20-24. As shown in SCHEME 20, tolyl or allylic substrates can be converted via

5 halogenation to benzylic or allylic bromides wherein M, X, W and Z are as described above in the Summary of the Invention. This reaction is typically performed with N-bromosuccinimide (NBS) in the presence of a radical initiator such as 2,2'-azobisisobutyronitrile (AIBN) or a peroxide, preferably benzoyl peroxide. Alternatively,

10 the reaction can be initiated with light. The reaction is performed in an inert solvent such as carbon tetrachloride or chloroform at a temperature of about 50°C to about 100°C.

SCHEME 21




15 SCHEME 21 sets forth the synthesis of alkylating agents useful for preparing compounds of Formula I where M represents a biaryl or aryl cyclic group. Suzuki-type coupling of an aryl iodide or bromide or a ring system containing a vinyl bromide or iodide ( $\text{Ar}^2$ ) with a methylaryl boronic acid ( $\text{Ar}^1$ ) using the conditions described in

20 SCHEME 9 provides compounds of Formula 34. Where a vinyl bromide or iodide is used, compounds of Formula 34 can be reduced to generate a fully saturated ring. The reduction is accomplished by hydrogenation in the presence of palladium or platinum catalysts typically in protic solvents such as methanol or ethanol; or in tetrahydrofuran or ethyl acetate. Halogenation of the methyl group using reagents and conditions as described in SCHEME 20 provides alkylating agents of Formula

25 35.

-85-


SCHEME 22



Another common method for accessing alkyl halides is by halogenation of an alcohol or an alcohol derivative. Alcohols are obtained from commercial sources or can be prepared using methods well known to those skilled in the art. For example, SCHEME 22 sets forth the reduction of a carboxylic acid or ester to the corresponding alcohol using reagents such as, but not limited to, sodium borohydride, lithium aluminum hydride, borane-tetrahydrofuran complex, borane-methyl sulfide complex, etc. The corresponding alkyl chlorides are typically prepared from the alcohols with reagents such as hydrogen chloride, thionyl chloride, phosphorous pentachloride, phosphorous oxychloride, or triphenylphosphine/carbon tetrachloride. For the preparation of alkyl bromides, the alcohol is commonly treated with reagents such as hydrogen bromide, phosphorous tribromide, triphenylphosphine/bromine, or carbonyldiimidazole/allyl bromide (Kamijo, T., Harada, H., Iizuka, K. Chem. Pharm. Bull. 1983, 38, 4189). To access alkyl iodides, an appropriate alcohol is typically reacted with reagents such as triphenylphosphine/iodine/imidazole or hydrogen iodide. Alternatively, alkyl chlorides can be converted to the more reactive alkyl bromides or alkyl iodides by reaction with an inorganic salt such as sodium bromide, lithium bromide, sodium iodide, or potassium iodide in solvents such as acetone or methyl ethyl ketone. Alkyl sulfonates can also be used as electrophiles or can be converted to alkyl halides. Alkyl sulfonates are prepared from the corresponding alcohol using a mild base such as triethylamine or pyridine and a sulfonyl chloride in an inert solvent such as methylene chloride or diethyl ether. If desired, conversion to the halide is accomplished by treating the alkyl sulfonate with an inorganic halide (sodium iodide, sodium bromide, potassium iodide, potassium bromide, lithium chloride, lithium bromide, etc) or a tetrabutylammonium halide.

-86-


SCHEME 23



- Cinnamic acids or esters are commonly available from commercial sources and can be converted to alkylating agents of Formulas 37 or 38 as follows (see
- 5 SCHEME 23). The cinnamic acid or ester derivatives are reduced by hydrogenation in the presence of palladium or platinum catalysts typically in protic solvents (e.g., methanol or ethanol), tetrahydrofuran, or ethyl acetate. Reduction and conversion to the alkyl halide or sulfonate as described in SCHEME 22 provides the compounds of Formula 38. Where appropriate, the cinnamic acids or esters are converted directly to
- 10 the alcohols of Formula 39 by treating those cinnamic acids or esters with reagents such as lithium aluminum hydride in an inert solvent such as tetrahydrofuran and diethyl ether. Alternatively, the cinnamic acid or ester can be reduced to an allylic alcohol of Formula 40 using reagents such as lithium aluminum hydride/aluminum chloride, diisobutylaluminum hydride or lithium borohydride. Conversion to the allylic halide or
- 15 sulfonate as described in SCHEME 22 provides the compounds of Formula 37.

-87-

## SCHEME 24




- 5           The preparation of alkylating agents of Formula 41 wherein W and M are as described above in the Summary of the Invention is set forth in SCHEME 24. Compounds of Formula 42 can be alkylated with a variety of bases. The choice of base is dependent on the nature of W and M. Some preferred bases include, but are not limited to, sodium hydroxide, sodium hydride, lithium diisopropylamide, lithium
- 10 bis(trimethylsilyl)amide, potassium bis(trimethylsilyl)amide and potassium tert-butoxide. Treatment of the resulting anion with one of a variety of dialkylhalides generates the desired alkylating agents of Formula 41. For the preparation of compounds where W is an oxygen and M is an aromatic ring, it is preferred to form the alkoxide anion with sodium hydroxide followed by addition of a dihaloalkane, e.g.
- 15 dibromoalkane. The reaction is normally performed in water at about 75°C to about 125°C.



-88-

SCHEME 25



Aldehydes useful for the method described in SCHEME 5 are available from commercial sources or can be prepared from available intermediates using methods well known to those skilled in the art (for a general reference see "The Chemistry of the Carbonyl Group," Ed. S. Patai, Interscience, New York (1966-70)). SCHEME 25 demonstrates an exemplary method used to prepare Formula 43 hydroxy aldehydes where M in SCHEME 5 contains a hydroxy substituted alkyl group. Treatment of a dialdehyde, where one of the aldehydes is protected as an acetal of Formula 44 wherein the OR groups are conventional substituents used in an acetal protecting group, with an organometallic reagent (LMetal), preferably an organolithium or Grignard reagent, in an inert solvent such as tetrahydrofuran or diethyl ether, provides compounds of Formula 45. Subsequent acetal hydrolysis under mildly acidic conditions, e.g., dilute hydrogen chloride, Amberlyst-15<sup>®</sup> resin, silica gel, or other reagents as described in "Protecting Groups in Organic Synthesis," Second Edition, T.W. Greene and P.G.M. Wuts, John Wiley and Sons, Inc., 1991 provides the desired hydroxy aldehydes of Formula 43.

Aldehydes useful for the method described in SCHEME 5 may be prepared using the methods described in SCHEMES 26-28. For example, as shown in SCHEME 26, an aryl boronic acid which contains an aldehyde can be coupled to an aryl bromide, aryl iodide, or a ring system which contains a vinyl bromide or iodide


-89-

using the Suzuki protocol described for SCHEME 9 to provide aldehydes of Formula 60.

SCHEME 27 describes the preparation of aldehydes of Formula 62 which contain a suitably protected acid moiety and can be used for the preparation of compounds of Formula 42A described in SCHEME 5. Selective reduction of nitriles (see SCHEMES 3-4 for preparations) of Formula 61 provides aldehydes of Formula 62. A preferred method for this reduction involves heating the nitrile with aluminum-nickel (Raney) alloy in the presence of an acid such as formic acid. Aldehydes of Formula 64 useful for the preparation of compounds of Formula 42 (SCHEME 5) may be prepared from starting nitriles of Formula 63 by treatment with a variety of reducing agents such as diisobutylaluminum hydride, tin (II) chloride/hydrogen chloride, or lithium triethoxyalanate.


A method for the preparation of proprionaldehydes of Formula 65 is described in SCHEME 28 and follows the procedures described by Kang (J. Org. Chem. 1996, 61, 2604) and by Jeffery (J. Chem. Soc. Chem. Commun. 1984, 19, 1287). An aryl iodide or bromide is coupled to allyl alcohol in the presence of a suitable palladium catalyst, preferably palladium acetate. The reaction is performed in a suitable polar, aprotic solvent, preferably dimethylformamide, with addition of a base, such as sodium bicarbonate, and an ammonium salt, such as tetrabutylammonium chloride and provides proprionaldehydes of Formula 65.

## SCHEME 26




-90-

## SCHEME 27



## SCHEME 28




## CHLOROMETHYL INTERMEDIATES


Intermediate chloromethyl compounds can be prepared as described in SCHEMES 29 and 30. In general, the appropriate Formula 66 or 68 sulfonamide or carboxamide is treated with a formaldehyde equivalent such as paraformaldehyde in an inert organic solvent such as methylene chloride or chloroform with a suitable catalyst such as HCl, zinc chloride or trimethylsilyl chloride at temperatures ranging from about 0°C to about 60°C to give the Formula 67 and 69 chloromethyl derivatives, respectively.

-91-

SCHEME 29



SCHEME 30




5

SCHEME 31 sets forth the synthesis of biaryl aldehydes of Formula 60. Fluorobenzonitriles of Formula 70 are heated with a nucleophilic group such as a pyrrazole or imidazole in a suitable solvent, preferably DMF to effect displacement of the fluoro group yielding intermediates of Formula 71. The desired biaryl aldehydes of Formula 60 are obtained by reduction of the nitrile via hydrogenation with Raney

10

-92-

alloy in formic acid, or by reduction with a hydride source such as diisobutylaluminum hydride.



5

The examples and synthetic procedures set forth in this application are intended to illustrate particular embodiments of the invention and are not intended to limit the scope of the specification or claim in any manner. All documents cited in this application are hereby incorporated by reference.

10

Example 17-((4-Butyl-benzyl)-(pyridine-3-sulfonyl)-amino)-heptanoic acidStep A: Reductive Amination

7-(4-Butyl-benzylamino)-heptanoic acid methyl ester. A solution of 7-amino-  
5 heptanoic acid methyl ester hydrochloride, prepared of Preparation 1, (1.12 g, 5.9  
mmol), 4-butyl-benzaldehyde (0.915 g, 5.65 mmol) and triethylamine (0.83 mL, 5.98  
mmol) in 20 mL MeOH was stirred at room temperature for 3 hours. After cooling to  
0°C, NaBH<sub>4</sub> (0.342 g, 9.04 mmol) was added and the reaction was stirred for 15  
10 minutes at room temperature. The mixture was quenched with 1:1 NaHCO<sub>3</sub>:H<sub>2</sub>O and  
the MeOH was removed *in vacuo*. The resulting residue was diluted with CH<sub>2</sub>Cl<sub>2</sub> and  
the organic solution was washed with water and brine, dried over MgSO<sub>4</sub>, filtered and  
concentrated *in vacuo* to afford the title compound of Step A (1.4 g). <sup>1</sup>H NMR (400  
MHz, CDCl<sub>3</sub>) δ 7.08-7.38 (m, 4H), 3.62 (s, 2H), 3.29 (s, 3H), 2.52-2.66 (m, 4H), 2.25  
(t, 2H), 1.53-1.63 (m, 6H), 1.25-1.40 (m, 6H), 0.85 (t, 3H); MS 306 (M+1).

Step B: Amide formation

7-((4-Butyl-benzyl)-(pyridine-3-sulfonyl)-amino)-heptanoic acid methyl ester. A  
solution of 7-(4-butyl-benzylamino)-heptanoic acid methyl ester prepared of Example  
1, Step A (0.10 g, 0.33 mmol), N,N-diisopropylethylamine (0.85 g 0.66 mmol) and  
pyridine-3-sulfonyl chloride hydrochloride, prepared of Preparation 2, (0.070 g, 0.33  
20 mmol) in 3 mL CH<sub>2</sub>Cl<sub>2</sub> was stirred at room temperature overnight. The mixture was  
diluted with CH<sub>2</sub>Cl<sub>2</sub> and the organic solution was washed with water and brine, dried  
over MgSO<sub>4</sub>, filtered and concentrated *in vacuo*. The product was purified by flash  
chromatography on silica gel (10% EtOAc/hexanes to 30% EtOAc/hexanes) to afford  
the title compound of Step B. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 9.01 (s, 1H), 8.75 (d,  
25 1H), 8.04 (d, 1H), 7.41 (dd, 1H), 7.23 (m, 4H), 4.30 (s, 2H), 3.62 (s, 3H), 3.08 (t, 2H),  
2.55 (t, 2H), 2.19 (t, 2H), 1.10-1.58 (m, 12H), 0.87 (t, 3H); MS 447 (M+1).

Step C: Ester Hydrolysis

7-((4-Butyl-benzyl)-(pyridine-3-sulfonyl)-amino)-heptanoic acid. A solution of 7-((4-  
butyl-benzyl)-(pyridine-3-sulfonyl)-amino)-heptanoic acid methyl ester prepared of  
30 Example 1, Step B (0.040 g, 0.158 mmol), in 2 mL MeOH and 0.5 mL 2N NaOH was  
stirred at room temperature overnight. The mixture was quenched with 2N HCl and  
was diluted with CH<sub>2</sub>Cl<sub>2</sub>. The organic layer was washed with 1N HCl and water, dried  
over MgSO<sub>4</sub>, filtered and concentrated *in vacuo*. The product was purified by flash

-94-

chromatography on silica gel (2% MeOH/CH<sub>2</sub>Cl<sub>2</sub> to 5% MeOH/CH<sub>2</sub>Cl<sub>2</sub>) to afford the title compound (42 mg). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 9.09 (s, 1H), 8.77 (d, 1H), 8.08 (d, 1H), 7.48 (dd, 1H), 7.09 (m, 4H), 4.32 (s, 2H), 3.12 (s, 2H), 2.55 (t, 2H), 2.25 (t, 2H), 1.12-1.58 (m, 12H), 0.88 (t, 3H); MS 431 (M-1).

5

Examples 1a-1an

Examples 1a-1an were prepared from the appropriate starting materials in a manner analogous to the method of Example 1, with variations in reaction time, temperature, and reagents as noted.

Example 1a

10

7-(Benzenesulfonyl-(4-butyl-benzyl)-amino)-heptanoic acid

<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.83 (d, 2H), 7.51-7.59 (m, 3H), 7.11 (m, 4H), 4.28 (s, 2H), 3.07 (t, 2H), 2.57 (t, 2H), 2.24 (t, 2H), 1.51-1.59 (m, 2H), 1.44-1.49 (m, 2H), 1.27-1.35 (m, 4H), 1.08-1.15 (m, 4H), 0.91 (t, 3H); MS 430 (M-1).

Example 1b

15

(3-(((1-Methyl-1H-indol-3-ylmethyl)-(pyridine-3-sulfonyl)-amino)-methyl)-phenyl)-acetic acid

<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 8.93 (s, 1H), 8.66 (s, 1H), 7.96 (d, 1H), 7.39 (d, 1H), 7.01-7.37 (m, 9H), 6.77 (s, 1H), 4.56 (s, 2H), 4.41 (s, 2H), 3.66 (s, 3H), 3.52 (s, 2H); MS 448 (M-1).

20

Example 1c(3-(((5-Phenyl-furan-2-ylmethyl)-(pyridine-3-sulfonyl)-amino)-methyl)-phenyl)-acetic acid

<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 8.02 (d, 1H), 7.22-7.34 (m, 12H), 6.42 (d, 1H), 6.17 (d, 1H), 4.45 (s, 2H), 4.40 (s, 2H), 3.60 (s, 2H); MS 461 (M-1).

25

Example 1d(3-(((5-Benzyl-pyridin-2-ylmethyl)-(pyridine-3-sulfonyl)-amino)-methyl)-phenyl)-acetic acid

Step A: Reaction time of 3.5 h at room temperature. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 8.97 (s, 1H), 8.71 (d, 1H), 8.15 (s, 1H), 7.98 (d, 1H), 7.44 (d, 1H), 7.04-7.34 (m, 10H), 4.54 (s, 2H), 4.43 (s, 2H), 3.87 (s, 2H), 3.50 (s, 2H); MS 486 (M-1).

30

Example 1e3-(((4-Phenethylsulfanyl-benzyl)-(pyridine-3-sulfonyl)-amino)-methyl)-phenyl)-acetic acid

-95-

Step A: Reaction time of 4 h at room temperature.  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  8.00 (d, 1H), 7.50 (bs, 1H), 6.90-7.38 (m, 15H), 4.31 (s, 4H), 3.49 (s, 2H), 3.11 (t, 2H), 2.87 (t, 2H); MS 531 (M-1).

Example 1f

5    (3-(((3-Hydroxy-4-propoxy-benzyl)-(pyridine-3-sulfonyl)-amino)-methyl)-phenyl)-acetic acid

Step A: Reaction time of 3.5 h at room temperature.  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  8.95 (s, 1H), 8.72 (s, 1H), 7.98 (d, 1H), 7.37 (m, 1H), 7.13-7.23 (m, 2H), 6.94-7.00 (m, 2H), 6.55-6.68 (m, 3H), 4.55 (s, 2H), 4.31 (s, 2H), 3.95 (t, 2H), 3.52 (s, 2H), 1.78 (m, 10    2H), 0.99 (t, 3H).

Example 1g

(3-(((4-Pentyl-benzyl)-(pyridine-3-sulfonyl)-amino)-methyl)-phenyl)-acetic acid

Step A: Reaction time of 3.5 h at room temperature.  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  8.98 (s, 1H), 8.74 (s, 1H), 8.00 (d, 1H), 7.39 (m, 1H), 7.14-7.26 (m, 2H), 6.95-7.05 (m, 15    6H), 4.35 (s, 4H), 3.54 (s, 2H), 2.54 (t, 2H), 1.56 (m, 2H), 1.29 (m, 4H), 0.88 (t, 3H); MS 465 (M-1).

Example 1h

(3-(((4-Methylsulfamoyl-benzyl)-(pyridine-3-sulfonyl)-amino)-methyl)-phenyl)-acetic acid

20    Step A: Reaction time of 3.5 h at room temperature.  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  9.06 (s, 1H), 8.85 (s, 1H), 8.16 (d, 1H), 7.53-7.64 (m, 3H), 6.91-7.26 (m, 6H), 4.39 (s, 2H), 4.35 (s, 2H), 3.50 (s, 2H), 2.63 (s, 3H); MS 488 (M-1).

Example 1i

(3-(((4-Isopropoxy-benzyl)-(pyridine-3-sulfonyl)-amino)-methyl)-phenyl)-acetic acid

25    Step A: Reaction time of 3.5 h at room temperature.  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  8.97 (s, 1H), 8.74 (s, 1H), 8.03 (m, 1H), 7.42 (m, 1H), 6.94-7.25 (m, 6H), 6.72 (m, 2H), 4.48 (m, 1H), 4.32 (m, 4H), 3.52 (s, 2H), 1.29 (t, 6H); MS 453 (M-1).

Example 1j

(3-(((4-Chloro-thiophen-2-ylmethyl)-(pyridine-3-sulfonyl)-amino)-methyl)-phenyl)-acetic acid

30

Step A: Reaction time of 3.5 h at room temperature.  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  9.01 (s, 1H), 8.79 (s, 1H), 8.07 (d, 1H), 7.45 (m, 1H), 7.20-7.29 (m, 2H), 7.12 (d, 1H), 7.10 (s, 1H), 7.07 (s, 1H), 4.46 (s, 2H), 4.42 (s, 2H), 3.60 (s, 2H); MS 435 (M-1).



-96-

Example 1k(3-(((4-Butyl-benzyl)-(4-nitro-benzenesulfonyl)-amino)-methyl)-phenyl)-acetic acid

<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 8.23 (m, 2H), 7.85 (m, 2H), 7.15-7.25 (m, 2H), 6.95-7.02 (m, 6H), 4.32 (m, 4H), 3.53 (s, 2H), 2.52 (m, 2H), 1.51 (m, 2H), 1.30 (m, 2H), 0.89 (t, 3H); MS 495 (M-1).

Example 1l(3-(((4-Butyl-benzyl)-(4-cyano-benzenesulfonyl)-amino)-methyl)-phenyl)-acetic acid

<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 8.21 (d, 1H), 7.67-7.84 (m, 3H), 6.89-7.24 (m, 8H), 4.46 (s, 1H), 4.38 (s, 1H), 4.32 (m, 2H), 3.54 (s, 1H), 3.38 (s, 1H), 2.55 (m, 2H), 1.58 (m, 2H), 1.33 (m, 2H), 1.29 (s, 1H), 0.89 (t, 3H); MS 475 (M-1).

Example 1m(3-(((4-Butyl-benzyl)-(3-fluoro-benzenesulfonyl)-amino)-methyl)-phenyl)-acetic acid

<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.58 (m, 1H), 7.45 (m, 1H), 6.92-7.24 (m, 10H), 4.29 (m, 4H), 3.52 (d, 2H), 2.52 (d, 2H), 1.52 (m, 2H), 1.29 (m, 2H), 0.90 (m, 3H); MS 468 (M-1).

Example 1n(3-(((4-Butyl-benzyl)-(5-pyridin-2-yl-thiophene-3-sulfonyl)-amino)-methyl)-phenyl)-acetic acid

Step B: N,N-diisopropylethylamine was replaced by triethylamine. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.81 (m, 2H), 7.17-7.27 (m, 6H), 6.94-7.16 (m, 6H), 4.29 (d, 4H), 3.55 (s, 2H), 2.54 (m, 2H), 1.54 (m, 2H), 1.31 (m, 2H), 0.91 (t, 3H); MS 533 (M-1).

Example 1o(3-(((4-Butyl-benzyl)-(toluene-4-sulfonyl)-amino)-methyl)-phenyl)-acetic acid

Step B: N,N-diisopropylethylamine was replaced by triethylamine. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.71 (d, 2H), 7.24-7.29 (m, 2H), 7.11-7.19 (m, 2H), 6.87-7.01 (m, 2H), 4.26 (d, 4H), 3.52 (s, 2H), 2.55 (m, 2H), 2.43 (s, 3H), 1.54 (m, 2H), 1.32 (m, 2H), 0.91 (t, 3H); MS 464 (M-1).

Example 1p(3-(((2,3-Dihydro-benzo[1,4]dioxin-6-ylmethyl)-(pyridine-3-sulfonyl)-amino)-methyl)-phenyl)-acetic acid

<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 8.98 (s, 1H), 8.76 (s, 1H), 8.02 (d, 1H), 7.40 (m, 1H), 7.14-7.26 (m, 2H), 7.02 (d, 1H), 6.96 (s, 1H), 6.72 (d, 1H), 6.59 (m, 2H), 4.35 (s, 2H), 4.25 (s, 2H), 4.20 (s, 4H), 3.55 (s, 2H); MS 453 (M-1).

-97-

Example 1q(3-(((Benzofuran-2-ylmethyl-(pyridine-3-sulfonyl)-amino)-methyl)-phenyl)-acetic acid

<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 9.05 (s, 1H), 8.66 (s, 1H), 8.04 (d, 1H), 7.11-7.42 (m, 9H), 6.44 (s, 1H), 4.45 (s, 1H), 4.39 (s, 1H), 3.59 (s, 1H); MS 435 (M-1).

5

Example 1r(3-(((4-Butyl-benzyl)-(1-methyl-1H-imidazole-4-sulfonyl)-amino)-methyl)-phenyl)-acetic acid

<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.58 (s, 1H), 7.28 (s, 1H), 6.99-7.26 (m, 8H), 4.33 (d, 4H), 3.65 (s, 3H), 3.52 (s, 2H), 2.54 (t, 2H), 1.54 (m, 2H), 1.32 (m, 2H), 0.91 (t, 3H);

10 MS 454 (M-1).

Example 1s(3-(((4-Imidazol-1-yl-benzyl)-(pyridine-3-sulfonyl)-amino)-methyl)-phenyl)-acetic acid

<sup>1</sup>H NMR (400 MHz, CD<sub>3</sub>OD) δ 9.45 (m, 1H), 9.44 (s, 1H), 9.03 (m, 1H), 8.91 (d, 1H), 8.19 (t, 1H), 8.04 (m, 1H), 7.77 (s, 1H), 7.61 (d, 2H), 7.53 (d, 2H), 7.11 (m, 4H), 4.70

15 (s, 2H), 4.51 (s, 2H), 3.33 (s, 2H); MS 461 (M-1).

Example 1t(3-(((Pyridine-3-sulfonyl)-(4-pyrimidin-2-yl-benzyl)-amino)-methyl)-phenyl)-acetic acid

<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 9.10 (s, 1H), 8.80 (m, 3H), 8.14 (d, 1H), 8.02 (d, 2H), 7.47 (m, 1H), 7.06-7.25 (m, 6H), 6.83 (s, 1H), 4.40 (s, 2H), 4.33 (s, 2H), 3.41 (s, 2H);

20 MS 473 (M-1).

Example 1u(3-(((Pyridine-3-sulfonyl)-(4-thiazol-2-yl-benzyl)-amino)-methyl)-phenyl)-acetic acid

<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 9.11 (s, 1H), 8.85 (s, 1H), 8.15 (d, 1H), 7.87 (s, 2H), 7.63 (d, 2H), 7.51 (m, 1H), 7.37 (s, 1H), 7.07-7.27 (m, 6H), 6.83 (s, 1H), 4.37 (s, 2H),

25 4.33 (s, 2H), 3.41 (s, 2H).

Example 1v(3-(((1-Methyl-1H-imidazole-4-sulfonyl)-(4-thiazol-2-yl-benzyl)-amino)-methyl)-phenyl)-acetic acid

Step B: N,N-diisopropylethylamine was replaced by triethylamine. <sup>1</sup>H NMR (400 MHz,

30 CDCl<sub>3</sub>) δ 7.85 (s, 1H), 7.59 (m, 2H), 7.47 (s, 1H), 7.34 (s, 1H), 7.07-7.25 (m, 6H),

6.88 (s, 1H), 4.46 (s, 2H), 4.38 (s, 2H), 3.77 (s, 3H), 3.40 (s, 2H); MS 483 (M-1).

-98-

Example 1w(3-(((4-Dimethylamino-benzyl)-(pyridine-3-sulfonyl)-amino)-methyl)-phenyl)-acetic acid

Step A: Reaction time of 4 h at room temperature. Step B: N,N-diisopropylethylamine was replaced by triethylamine. <sup>1</sup>H NMR (400 MHz, CD<sub>3</sub>OD) δ 8.09 (d, 1H), 7.09-7.16 (m, 2H), 6.93-6.99 (m, 7H), 6.65 (d, 2H), 5.36 (s, 2H), 4.32 (s, 2H), 4.27 (s, 2H), 2.89 (s, 6H); MS 438 (M-1).

Example 1x(3-(((4-Cyclohexyl-benzyl)-(pyridine-3-sulfonyl)-amino)-methyl)-phenoxy)-acetic acid

Step B: N,N-diisopropylethylamine was replaced by triethylamine. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 8.95 (s, 1H), 8.73 (d, 1H), 8.00 (d, 1H), 7.39 (m, 1H), 7.17 (t, 1H), 7.13 (d, 2H), 7.08 (d, 2H), 6.81 (d, 1H), 6.73 (d, 1H), 6.61 (s, 1H), 4.54 (s, 2H), 4.34 (s, 4H), 2.43 (m, 1H), 1.81 (d, 4H), 1.37 (t, 4H), 1.23 (m, 1H); MS 495 (M+1), 493 (M-1).

Example 1y

(3-(((2-(3,5-Dichloro-phenoxy)-ethyl)-(pyridine-3-sulfonyl)-amino)-methyl)-phenoxy)-acetic acid

Step B: N,N-diisopropylethylamine was replaced by triethylamine. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 9.07 (s, 1H), 8.78 (d, 1H), 8.12 (d, 1H), 7.47 (m, 1H), 7.25 (m, 1H), 6.82-6.91 (m, 4H), 6.53 (s, 2H), 4.61 (s, 2H), 4.47 (s, 2H), 3.91 (t, 2H), 3.54 (t, 2H); MS 511 (M+1), 509 (M-1).

Example 1z(3-(((4-Dimethylamino-benzyl)-(pyridine-3-sulfonyl)-amino)-methyl)-phenoxy)-acetic acid

Step B: N,N-diisopropylethylamine was replaced by triethylamine. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 8.91 (s, 1H), 8.79 (m, 1H), 8.04 (d, 1H), 7.43 (m, 1H), 7.16 (t, 1H), 6.94 (d, 2H), 6.81 (d, 2H), 6.64 (d, 2H), 6.49 (s, 1H), 4.51 (s, 2H), 4.28 (s, 4H), 2.91 (s, 6H); MS 456 (M+1), 454 (M-1).

Example 1aa(3-(((4-tert-Butyl-benzyl)-(pyridine-3-sulfonyl)-amino)-methyl)-phenoxy)-acetic acid

Step B: N,N-diisopropylethylamine was replaced by triethylamine. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 8.95 (s, 1H), 8.74 (s, 1H), 7.99 (d, 1H), 7.39 (m, 1H), 7.25 (m, 2H), 7.15 (t, 1H), 7.04 (d, 2H), 6.81 (d, 1H), 6.72 (d, 1H), 6.62 (s, 1H), 4.55 (s, 2H), 4.35 (s, 4H), 1.27 (s, 9H); MS 469 (M+1), 467 (M-1).

-99-

Example 1ab(3-(((3-(3-Chloro-phenyl)-propyl)-(pyridine-3-sulfonyl)-amino)-methyl)-phenoxy)-acetic acid

Step B: N,N-diisopropylethylamine was replaced by triethylamine. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 8.98 (s, 1H), 8.77 (d, 1H), 8.07 (d, 1H), 7.48 (m, 1H), 7.21 (m, 2H), 6.91 (s, 1H), 6.86 (m, 3H), 6.78 (s, 1H), 4.61 (s, 2H), 4.31 (s, 2H), 3.15 (t, 2H), 2.43 (t, 2H), 1.68 (m, 2H); MS 475 (M+1), 473 (M-1).

Example 1ac

10 (3-(((4-tert-Butyl-benzyl)-(1-methyl-1H-imidazole-4-sulfonyl)-amino)-methyl)-phenoxy)-acetic acid

Step B: N,N-diisopropylethylamine was replaced by triethylamine. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.66 (s, 1H), 7.08-7.31 (m, 6H), 6.70-6.78 (m, 3H), 4.54 (s, 2H), 4.35 (s, 4H), 3.68 (s, 3H), 1.27 (s, 9H); MS 469.9 (M-1).

Example 1ad

15 (3-(((4-Cyclohexyl-benzyl)-(pyridine-3-sulfonyl)-amino)-methyl)-phenyl)-acetic acid

Step B: N,N-diisopropylethylamine was replaced by triethylamine. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 8.98 (bs, 1H), 8.75 (bs, 1H), 7.98 (d, 1H), 7.39 (bs, 1H), 6.97-7.25 (m, 8H), 4.36 (d, 4H), 3.54 (s, 2H), 2.44 (s, 1H), 1.72-1.82 (m, 4H), 1.24-1.36 (m, 5H); MS 476.9 (M-1).

20

Example 1ae

(3-(((1-Methyl-1H-imidazole-4-sulfonyl)-(4-phenoxy-benzyl)-amino)-methyl)-phenyl)-acetic acid

Step B: N,N-diisopropylethylamine was replaced by triethylamine. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.52 (s, 1H), 7.06-7.37 (m, 10H), 6.94 (d, 2H), 6.83 (d, 2H), 4.38 (s, 4H), 3.71 (s, 3H), 1.72-1.82 (m, 4H), 3.56 (s, 2H); MS 490 (M-1).

25

Example 1af

(3-(((4-Phenoxy-benzyl)-(pyridine-3-sulfonyl)-amino)-methyl)-phenyl)-acetic acid

<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 9.00 (bs, 1H), 8.76 (bs, 1H), 8.04 (d, 1H), 7.41 (t, 1H), 7.35 (m, 1H), 6.86-7.32 (m, 10H), 6.84 (d, 2H), 4.37 (d, 4H), 3.54 (s, 2H); MS 487 (M-1).

30

Example 1ag

(3-(((4-(2-Oxo-pyrrolidin-1-yl)-benzyl)-(pyridine-3-sulfonyl)-amino)-methyl)-phenyl)-acetic acid

-100-

Step B: N,N-diisopropylethylamine was replaced by triethylamine.  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  9.06 (bs, 1H), 8.80 (bs, 1H), 8.14 (m, 1H), 7.47 (m, 1H), 6.96-7.26 (m, 7H), 4.28 (m, 4H), 3.78 (m, 2H), 3.35 (m, 2H), 2.59 (m, 2H), 2.11 (m, 2H); MS 478 (M-1).

Example 1ah

5     (3-((Benzo[1,3]dioxol-5-ylmethyl-(pyridine-3-sulfonyl)-amino)-methyl)-phenyl)-acetic acid

$^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  8.98 (s, 1H), 8.76 (s, 1H), 8.04 (d, 1H), 7.41 (m, 1H), 7.14-7.20 (m, 2H), 7.00 (d, 1H), 6.94 (s, 1H), 6.64 (t, 2H), 6.55 (d, 1H), 4.34 (s, 2H), 4.26 (s, 2H), 3.54 (s, 2H); MS 439 (M-1).

10                     Example 1ai

(3-(((1-Methyl-1H-imidazole-4-sulfonyl)-(4-pyrimidin-5-yl-benzyl)-amino)-methyl)-phenyl)-acetic acid

Step B: N,N-diisopropylethylamine was replaced by triethylamine.  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  9.18 (s, 1H), 8.91 (s, 2H), 7.05-7.54 (m, 11H), 4.49 (s, 2H), 4.40 (s, 2H), 3.75 (s, 3H), 3.55 (s, 2H); MS 476 (M-1).

Example 1aj

(3-(((Pyridine-3-sulfonyl)-(4-pyrimidin-5-yl-benzyl)-amino)-methyl)-phenyl)-acetic acid

$^1\text{H}$  NMR (400 MHz,  $\text{CD}_3\text{OD}$ )  $\delta$  9.17 (s, 1H), 9.01 (s, 1H), 8.77 (s, 1H), 7.57 (m, 4H), 7.45 (d, 2H), 7.05-7.16 (m, 5H), 4.48 (s, 2H), 4.43 (s, 2H), 3.45 (s, 2H).

20                     Example 1ak

(3-(((4-Pyrazin-2-yl-benzyl)-(pyridine-3-sulfonyl)-amino)-methyl)-phenyl)-acetic acid

$^1\text{H}$  NMR (400 MHz,  $\text{DMSO}-d_6$ )  $\delta$  9.18 (s, 1H), 9.02 (s, 1H), 8.83 (d, 1H), 8.68 (s, 1H), 8.57 (s, 1H), 8.25 (d, 1H), 7.96 (d, 2H), 7.60 (m, 1H), 7.26 (d, 2H), 7.15 (m, 2H), 7.05 (m, 2H), 4.42 (s, 2H), 4.41 (s, 2H).

25                     Example 1al

(3-(((1-Methyl-1H-imidazole-4-sulfonyl)-(4-pyrimidin-2-yl-benzyl)-amino)-methyl)-phenyl)-acetic acid

$^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  8.78 (d, 2H), 7.94 (d, 2H), 7.54 (s, 1H), 7.44 (s, 1H), 7.22-7.03 (m, 6H), 6.87 (s, 1H), 4.45 (s, 2H), 4.39 (s, 2H), 3.73 (s, 3H), 3.38 (s, 2H);

30     MS 476 (M-1).

Example 1am

(3-(((4-Butyl-benzyl)-phenylmethanesulfonyl-amino)-methyl)-phenyl)-acetic acid

-101-

<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.31-6.96 (m, 13H), 4.13 (s, 2H), 4.05 (s, 2H), 4.03 (s, 2H), 3.62 (s, 2H), 2.60 (t, 2H), 1.58 (m, 2H), 1.33 (m, 2H), 0.91 (t, 3H); MS 464 (M-1).

Example 1an

5-(3-((Pyridine-3-sulfonyl)-(4-thiazol-2-yl-benzyl)-amino)-propyl)-thiophene-2-carboxylic acid

Step A: Triethylamine was replaced with N,N-diisopropylethylamine. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 9.18 (d, 1H), 8.82 (d, 1H), 8.05 (d, 1H), 7.73-7.20 (m, 8H), 6.60 (d, 1H), 4.35 (s, 2H), 3.22 (t, 2H), 2.70 (t, 2H), 1.85-1.70 (m, 2H).

Example 2

10 (3-(((2-(3-Chloro-phenoxy)-ethyl)-(pyridine-3-sulfonyl)-amino)-methyl)-phenyl)-acetic acid

Step A: Alkylation

(3-(((2-(3-Chloro-phenoxy)-ethyl)-(pyridine-3-sulfonyl)-amino)-methyl)-phenyl)-acetic acid methyl ester. To a solution of sodium hydride (60% in mineral oil, 0.016 g, 0.3996 mmol) in 2 mL DMF was added (3-((pyridine-3-sulfonylamino)-methyl)-phenyl)-acetic acid methyl ester (from Preparation 14, 0.096 g, 0.333 mmol) at 0°C and the reaction was stirred at room temperature for 30 minutes. After cooling to 0°C, 1-(2-bromo-ethoxy)-3-chloro-benzene (from Preparation 29, 0.094 g, 0.399 mmol) was added and the reaction was stirred at room temperature overnight. The DMF was removed *in vacuo*. The residue was diluted with EtOAc and the organic solution was washed with water and brine, dried over MgSO<sub>4</sub>, filtered and concentrated *in vacuo*. The product was purified by flash chromatography on silica gel (0.5% MeOH/CH<sub>2</sub>Cl<sub>2</sub> to 2% MeOH/CH<sub>2</sub>Cl<sub>2</sub>) to afford the title compound of Step A (0.025 g). MS 475 (M+1).

Step B: Ester Hydrolysis

(3-(((2-(3-Chloro-phenoxy)-ethyl)-(pyridine-3-sulfonyl)-amino)-methyl)-phenyl)-acetic acid. A solution of the compound of Example 2, Step A (0.025 g, 0.053 mmol), in 2 mL MeOH and 0.5 mL 2N NaOH was stirred at room temperature overnight. The mixture was quenched with 2N HCl and was diluted with CH<sub>2</sub>Cl<sub>2</sub>. The organic layer was washed with 1N HCl and water, dried over MgSO<sub>4</sub>, filtered, and concentrated *in vacuo*. The product was purified by flash chromatography on silica gel (2% MeOH/CH<sub>2</sub>Cl<sub>2</sub> to 5% MeOH/CH<sub>2</sub>Cl<sub>2</sub>) to afford the title compound (20 mg). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 9.05 (s, 1H), 8.77 (d, 1H), 8.11 (d, 1H), 7.43 (m, 1H), 7.08-7.27

-102-

(m, 5H), 6.89 (d, 1H), 6.62 (s, 1H), 6.55 (d, 1H), 4.51 (s, 2H), 3.95 (t, 2H), 3.59 (s, 4H); MS 495 (M-2).

#### Examples 2a-2c

Examples 2a-2c were prepared from the appropriate starting materials in a manner  
5 analogous to the method of Example 2.

#### Example 2a

Trans-(3-(((3-(3,5-Dichloro-phenyl)-allyl)-(pyridine-3-sulfonyl)-amino)-methyl)-  
phenyl)-acetic acid

<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 9.08 (bs, 1H), 8.81 (bs, 1H), 8.11 (d, 1H), 7.48 (bs, 1H),  
10 7.12-7.28 (m, 4H), 6.98 (s, 2H), 6.19 (d, 1H), 5.86 (m, 1H), 4.38 (s, 2H), 3.93 (d, 2H),  
3.58 (s, 2H).

#### Example 2b

(3-(((2-(3,5-Dichloro-phenoxy)-ethyl)-(pyridine-3-sulfonyl)-amino)-methyl)-phenyl)-  
acetic acid

15 <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 8.96 (bs, 1H), 8.70 (bs, 1H), 8.04 (d, 1H), 7.41 (m, 1H),  
7.24-7.09 (m, 4H), 6.86 (s, 1H), 6.47 (s, 2H), 4.44 (s, 2H), 3.86 (m, 2H), 3.49 (s, 2H),  
3.31 (m, 2H).

#### Example 2c

(3-(((4-(1-Hydroxy-hexyl)-benzyl)-(pyridine-3-sulfonyl)-amino)-  
20 methyl)-phenyl)-acetic acid

<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 8.91 (bs, 1H), 8.72 (bs, 1H), 8.03 (d, 1H), 7.40 (bs, 1H),  
7.16-6.99 (m, 7H), 6.81 (s, 1H), 4.57 (t, 1H), 4.29 (s, 4H), 3.43 (m, 2H), 1.70 (m, 1H),  
1.61 (m, 1H), 1.32-1.16 (m, 8H), 0.82 (t, 3H).

#### Example 3

25 5-(3-((2-Benzylsulfanyl-ethyl)-(pyridine-3-sulfonyl)-amino)-propyl)-thiophene-2-  
carboxylic acid

#### Step A: Reductive Amination

5-(3-(2-Benzylsulfanyl-ethylamino)-propyl)-thiophene-2-carboxylic acid tert-butyl  
ester. Step A was performed in a manner analogous to the method of Step A of  
30 Example 1.

#### Step B: Amide Formation

5-(3-((2-Benzylsulfanyl-ethyl)-(pyridine-3-sulfonyl)-amino)-propyl)-thiophene-2-  
carboxylic acid tert-butyl ester. Step B was performed in a manner analogous to the

-103-

method of Step B of Example 1, except triethylamine was used in place of N,N-diisopropylethylamine.

Step C: Ester Hydrolysis

5 5-(3-((2-Benzylsulfanyl-ethyl)-(pyridine-3-sulfonyl)-amino)-propyl)-thiophene-2-carboxylic acid•TFA. A solution of 5-(3-((2-benzylsulfanyl-ethyl)-(pyridine-3-sulfonyl)-amino)-propyl)-thiophene-2-carboxylic acid tert-butyl ester prepared of Example 3, Step B (0.038 g) in 1 mL CH<sub>2</sub>Cl<sub>2</sub> was cooled to 0°C and 1 mL TFA was added. The mixture was warmed to room temperature and was stirred for 1 h. The CH<sub>2</sub>Cl<sub>2</sub> and TFA were removed by evaporation, azeotroping with added CH<sub>2</sub>Cl<sub>2</sub> to yield the title  
10 compound (46.3 mg). MS 475 (M-1).

Examples 3a-3i were prepared from the appropriate starting materials in a manner analogous to the method of Example 3 with variations thereto noted.

Example 3a

15 5-(3-((2-(3-Chloro-phenylsulfanyl)-ethyl)-(pyridine-3-sulfonyl)-amino)-propyl)-thiophene-2-carboxylic acid  
<sup>1</sup>H NMR (400 MHz, CD<sub>3</sub>OD) δ 8.93 (s, 1H), 8.78 (d, 1H), 8.21 (d, 1H), 7.64 (m, 1H), 7.57 (s, 1H), 7.35 (s, 1H), 7.19-7.28 (m, 3H), 6.87 (s, 1H), 3.16-3.35 (m, 6H), 2.87 (t, 2H), 1.89 (t, 2H); MS 497,499 (M+).

Example 3b

20 (3-(((Pyridine-3-sulfonyl)-(4-thiazol-2-yl-benzyl)-amino)-methyl)-phenoxy)-acetic acid•2TFA  
<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 9.40 (bs, 1H), 8.98 (s, 1H), 8.84 (s, 1H), 8.28 (m, 1H), 8.10 (s, 1H), 7.78 (m, 2H), 7.68 (m, 1H), 7.51 (s, 1H), 7.24 (m, 3H), 7.12 (t, 1H), 6.77 (m, 1H), 6.48 (s, 1H), 4.53 (s, 2H), 4.45 (s, 2H), 4.34 (s, 2H); MS 494 (M-1).

25 Example 3c

(3-(((Pyridine-3-sulfonyl)-(4-pyrimidin-2-yl-benzyl)-amino)-methyl)-phenoxy)-acetic acid•2HCl

The TFA salt was converted to the HCl salt by stirring in 2 equivalents 1N HCl followed by removal of water and drying *in vacuo*. <sup>1</sup>H NMR (400 MHz, CD<sub>3</sub>OD) δ 9.00  
30 (d, 2H), 8.78 (d, 1H), 8.25 (d, 2H), 8.08 (t, 1H), 7.60 (t, 1H), 7.42 (m, 3H), 7.11 (m, 1H), 6.81 (d, 1H), 6.72 (m, 3H), 4.65 (s, 2H), 4.60 (s, 2H), 4.49 (s, 2H).



-104-

Example 3d

(3-(((1-Methyl-1H-imidazole-4-sulfonyl)-(4-thiazol-2-yl-benzyl)-amino)-methyl)-phenoxy)-acetic acid•2TFA

<sup>1</sup>H NMR (400 MHz, CD<sub>3</sub>OD) δ 7.93 (s, 1H), 7.85 (d, 1H), 7.76 (d, 2H), 7.70 (s, 1H),  
5 7.60 (d, 1H), 7.26 (d, 2H), 7.09 (t, 1H), 6.75 (d, 2H), 6.68 (s, 1H), 4.51 (s, 2H), 4.41 (s, 2H), 4.35 (s, 2H), 3.76 (s, 3H); MS 498 (M<sup>+</sup>).

Example 3e

(3-(((Pyridine-3-sulfonyl)-(4-pyridin-2-yl-benzyl)-amino)-methyl)-phenoxy)-acetic acid  
•HCl

10 No triethylamine was used in Step A. The TFA salt was converted to the HCl salt by stirring in 2 equivalents 1N HCl followed by removal of water and drying *in vacuo*. MS 490 (M+1), 488 (M-1).

Example 3f

(3-(((1-Methyl-1H-imidazole-4-sulfonyl)-(4-pyridin-2-yl-benzyl)-amino)-methyl)-phenoxy)-acetic acid•HCl

15 No triethylamine was used in Step A. The TFA salt was converted to the HCl salt by stirring in 2 equivalents 1N HCl followed by removal of water and drying *in vacuo*. MS 493 (M+1), 491 (M-1).

Example 3g

20 (3-(((Pyridine-3-sulfonyl)-(4-pyridin-3-yl-benzyl)-amino)-methyl)-phenoxy)-acetic acid•HCl

No triethylamine was used in Step A. The TFA salt was converted to the HCl salt by stirring in 2 equivalents 1N HCl followed by removal of water and drying *in vacuo*. MS 490 (M+1), 488 (M-1).

25 Example 3h

(3-(((1-Methyl-1H-imidazole-4-sulfonyl)-(4-pyridin-3-yl-benzyl)-amino)-methyl)-phenoxy)-acetic acid•HCl

No triethylamine was used in Step A. The TFA salt was converted to the HCl salt by stirring in 2 equivalents 1N HCl followed by removal of water and drying *in vacuo*. MS  
30 493 (M+1), 491 (M-1).

Example 3i

(3-(((Pyridine-3-sulfonyl)-(4-pyridin-4-yl-benzyl)-amino)-methyl)-phenoxy)-acetic acid  
•HCl

-105-

No triethylamine was used in Step A. The TFA salt was converted to the HCl salt by stirring in 2 equivalents 1N HCl followed by removal of water and drying *in vacuo*. MS 490 (M+1), 488 (M-1).

#### Example 4

5     5-(3-((3-(3-Chloro-phenyl)-propyl)-(pyridine-3-sulfonyl)-amino)-propyl)-thiophene-2-carboxylic acid

#### Step A: Sulfonamide formation

5-(3-((3-(3-Chloro-phenyl)-propyl)-(pyridine-3-sulfonyl)-amino)-propyl)-thiophene-2-carboxylic acid methyl ester. A solution of 5-(3-(3-(3-chloro-phenyl)-propylamino)-propyl)-thiophene-2-carboxylic acid methyl ester (from Preparation 8, 0.0855 g, 0.243 mmol), triethylamine (0.0541 g 0.534 mmol), and pyridine-3-sulfonyl chloride hydrochloride (from Preparation 2, 0.0572 g, 0.267 mmol) in 10 mL CH<sub>2</sub>Cl<sub>2</sub> combined at 0°C was stirred at room temperature overnight. The organic solution was washed with water, saturated NaHCO<sub>3</sub> and brine, dried over MgSO<sub>4</sub>, filtered and concentrated *in vacuo* to afford the title compound of Step A as an oil. MS 494 (M+1).

#### Step B: Ester Hydrolysis

5-(3-((3-(3-Chloro-phenyl)-propyl)-(pyridine-3-sulfonyl)-amino)-propyl)-thiophene-2-carboxylic acid. A solution of 5-(3-((3-(3-chloro-phenyl)-propyl)-(pyridine-3-sulfonyl)-amino)-propyl)-thiophene-2-carboxylic acid methyl ester prepared of Example 4, Step B (0.119 g, 0.241 mmol), in 5 mL EtOH and 0.72 mL 1N NaOH was stirred at room temperature overnight. The reaction mixture was adjusted to pH 6.2 and the layers were separated. The organic solution was washed with water, dried over MgSO<sub>4</sub>, filtered and concentrated *in vacuo* to afford the title compound (16 mg). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 8.00 (d, 1H, J=8), 7.70 (d, 1H, J=4), 7.30-7.60 (m, 6H), 6.75 (d, 1H, J=4), 3.20 (m, 4H), 2.95 (t, 2H, J=7), 2.60 (t, 2H, J=7), 1.70-2.00 (m, 4H); MS 478 (M+1), 476 (M-1).

#### Examples 4a-4h

Examples 4a-4h were prepared from the appropriate starting in a manner analogous to the method of Example 4.

30

#### Example 4a

5-(3-((3-(3-Chloro-phenyl)-propyl)-(4-methoxy-benzenesulfonyl)-amino)-propyl)-thiophene-2-carboxylic acid

-106-

$^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.70 (d, 1H,  $J=7$ ), 7.00-7.40 (m, 8H), 6.80 (d, 1H,  $J=4$ ), 3.89 (s, 3H), 3.10 (m, 4H), 2.95 (t, 2H,  $J=7$ ), 2.50 (t, 2H,  $J=7$ ), 1.70-2.00 (m, 2H); MS 508 (M+1), 506 (M-1).

Example 4b

5 5-(3-((Benzo[1,2,5]thiadiazole-4-sulfonyl)-(3-(3-chloro-phenyl)-propyl)-amino)-propyl)-thiophene-2-carboxylic acid

$^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.00-7.70 (m, 8H), 6.70 (d, 1H,  $J=4$ ), 3.05 (m, 4H), 2.90 (t, 2H,  $J=7$ ), 2.54 (t, 2H,  $J=7$ ), 1.72-1.92 (m, 2H); MS 536 (M+), 535 (M-1).

Example 4c

10 5-(3-(Benzenesulfonyl-(3-(3-chloro-phenyl)-propyl)-amino)-propyl)-thiophene-2-carboxylic acid

$^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  6.70-7.92 (m, 11H), 3.26 (m, 4H), 3.05 (m, 4H), 2.73 (m, 2H), 2.50 (m, 2H), 1.70 (m, 2H); MS 578(M+1), 576 (M-1).

Example 4d

15 5-(3-((3-(3-Chloro-phenyl)-propyl)-phenylmethanesulfonyl-amino)-propyl)-thiophene-2-carboxylic acid

$^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.50 (d, 1H,  $J=4$ ), 7.00-7.40 (m, 9H), 6.85 (d, 1H,  $J=4$ ), 3.00 (m, 4H), 2.60 (m, 2H), 2.40 (m, 2H), 1.60-1.80 (m, 2H); MS 490 (M-1).

Example 4e

20 5-(3-((3-(3-Chloro-phenyl)-propyl)-(pyridine-3-sulfonyl)-amino)-propyl)-furan-2-carboxylic acid

$^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  9.00 (m, 1H), 8.70 (m, 1H), 8.00 (d, 1H,  $J=6$ ), 7.50 (m, 1H), 6.80-7.04 (m, 6H), 3.20 (m, 4H), 2.78 (m, 2H), 2.50 (m, 2H), 1.62-2.00 (m, 4H); MS 463 (M+1), 461 (M-1).

25

Example 4f

5-(3-((3-(3-Chloro-phenyl)-propyl)-(naphthalene-2-sulfonyl)-amino)-propyl)-thiophene-2-carboxylic acid

$^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  8.40 (d, 1H,  $J=2$ ), 7.00-8.00 (m, 11H), 6.80 (d, 1H,  $J=4$ ), 3.20 (m, 4H), 2.82 (t, 2H,  $J=7$ ), 2.60 (t, 2H,  $J=7$ ), 1.80-2.00 (m, 2H); MS 528.9 (M+1).

30

Example 4g

5-(3-((3-(3-Chloro-phenyl)-propyl)-(naphthalene-1-sulfonyl)-amino)-propyl)-thiophene-2-carboxylic acid

-107-

<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 8.60 (d, 1H, J=5), 6.95-8.22 (m, 11H), 6.70 (d, 1H, J=4), 3.20 (m, 4H), 2.40 (t, 2H, J=7), 1.72-1.95 (m, 4H); MS 528.9 (M+1).

#### Example 4h

5 5-(3-((2-Acetylamino-4-methyl-thiazole-5-sulfonyl)-(3-(3-chloro-phenyl)-propyl)-amino)-propyl)-thiophene-2-carboxylic acid

<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.61 (d, 1H, J=4), 7.00-7.30 (m, 4H), 3.60 (d, 1H, J=3.8), 2.80 (t, 2H, J=7.0), 2.60 (t, 2H, J=6.8), 2.40 (s, 3H), 2.30 (s, 3H), 1.70-2.00 (m, 4H); MS 556 (M+1), 554 (M-1).

#### Example 5

10 5-(3-((3-(3-Chloro-phenyl)-propyl)-(pyridine-3-carbonyl)-amino)-propyl)-thiophene-2-carboxylic acid

##### Step A: Amide formation

5-(3-((3-(3-Chloro-phenyl)-propyl)-(pyridine-3-carbonyl)-amino)-propyl)-thiophene-2-carboxylic acid methyl ester. A solution of 5-(3-(3-(3-chloro-phenyl)-propylamino)-propyl)-thiophene-2-carboxylic acid methyl ester (from Preparation 8, 0.075 g, 0.213 mmol), DCC (0.0483 g 0.234 mmol) and nicotinic acid (0.0289 g, 0.234 mmol) in 10 mL CH<sub>2</sub>Cl<sub>2</sub> was stirred at room temperature overnight. The mixture was filtered and the filtrate was concentrated *in vacuo*. The residue was dissolved in 15 mL EtOAc and the insolubles were removed via filtration. The organic solution was washed with water followed by brine, dried over MgSO<sub>4</sub>, filtered, and concentrated *in vacuo* to afford the title compound of Step A as an oil (113 mg). MS 457 (M+).

##### Step B: Ester Hydrolysis

Step B was performed in a manner analogous to the method of Step B of Example 4. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 8.60 (d, 1H, J=8), 6.80-7.70 (m, 8H), 6.60 (d, 1H, J=4), 3.25 (m, 4H), 2.80 (m, 2H), 2.45 (m, 2H), 1.60-2.05 (m, 4H); MS 443 (M+1), 441 (M-1).

#### Examples 5a-5b

Examples 5a-5b were prepared from the appropriate starting in a manner analogous to the method of Example 5.

30 Example 5a

5-(3-((3-(3-Chloro-phenyl)-propyl)-(pyridin-2-yl-acetyl)-amino)-propyl)-thiophene-2-carboxylic acid

-108-

<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 8.60 (m, 1H), 7.00-7.80 (m, 8H), 6.60 (m, 1H), 4.00 (s, 2H), 3.32 (m, 4H), 2.72 (m, 2H), 2.50 (m, 2H), 1.70-2.00 (m, 4H); MS 457 (M+1), 455 (M-1).

Example 5b

5     5-(3-((3-(3-Chloro-phenyl)-propyl)-(pyridin-3-yl-acetyl)-amino)-propyl)-thiophene-2-carboxylic acid

<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.60-7.80 (m, 2H), 7.00-7.50 (m, 7H), 6.70 (d, 1H, J=4), 3.60 (s, 2H), 3.10-3.40 (m, 4H), 2.80 (m, 2H), 2.60 (m, 2H), 1.70-2.00 (m, 4H); MS 457 (M+1), 455 (M-1).

10

Example 6

5-(3-((2-Chloro-benzenesulfonyl)-(3-(3-chloro-phenyl)-propyl)-amino)-propyl)-thiophene-2-carboxylic acid

Step A: Amide formation

15     5-(3-((2-Chloro-benzenesulfonyl)-(3-(3-chloro-phenyl)-propyl)-amino)-propyl)-thiophene-2-carboxylic acid tert-butyl ester. A stock solution of 5-(3-(3-(3-chloro-phenyl)-propylamino)-propyl)-thiophene-2-carboxylic acid tert-butyl ester (from Preparation 9, 0.10 g, 0.254 mmol) in 10 mL CH<sub>2</sub>Cl<sub>2</sub> was prepared and 1 mL of the solution (0.010 g, 0.0254 mmol) was added to a 1 dram vial. To this was added triethylamine (0.78 mL, 0.056 mmol) and 2-chloro-benzenesulfonyl chloride (0.0059 g, 0.028 mmol). The reaction was stirred overnight at room temperature and was diluted with 2 mL CH<sub>2</sub>Cl<sub>2</sub>. The organic solution was washed with 3 mL of 5.5% aqueous HCl solution (2X) and 3 mL saturated bicarbonate solution (2X). The organic layer was dried with MgSO<sub>4</sub> and was concentrated to yield the title compound of Step A (10 mg).

25

Step B: Ester Hydrolysis

30     5-(3-((2-Chloro-benzenesulfonyl)-(3-(3-chloro-phenyl)-propyl)-amino)-propyl)-thiophene-2-carboxylic acid. A solution of 5-(3-((2-chloro-benzenesulfonyl)-(3-(3-chloro-phenyl)-propyl)-amino)-propyl)-thiophene-2-carboxylic acid tert-butyl ester prepared of Example 6, Step A (0.010 g, 0.010 mmol) in 4N HCl in 1,4 dioxane (3 mL) and the reaction was stirred overnight at room temperature. HCl (g) was bubbled in until reaction was determined to be complete by thin layer chromatography. The reaction mixture was concentrated *in vacuo*. The resulting organic residue was azeotroped with CCl<sub>4</sub> to produce a powder (5 mg). <sup>1</sup>H NMR (400

-109-

MHz, CDCl<sub>3</sub>) δ 8.00 (d, 1H, J=4), 7.00-7.72 (m, 8H), 6.75 (d, 1H, J=4), 3.20-3.40 (m, 4H), 2.81 (m, 2H), 2.52 (m, 2H), 1.90 (m, 2H), 1.80 (m, 2H), 1.20 (m, 2H); MS 509.9 (M-1).

#### Examples 6a-6j

- 5 Examples 6a-6j were prepared from the appropriate starting material in a manner analogous to the method of Example 6.

#### Example 6a

5-(3-((3-(3-Chloro-phenyl)-propyl)-(2,5-dimethyl-benzenesulfonyl)-amino)-propyl)-thiophene-2 carboxylic acid

- 10 <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.70 (d, 1H, J=7), 7.00-7.40 (m, 7H), 6.80 (d, 1H, J=4), 3.32 (m, 4H), 2.50 (s, 3H), 2.36 (s, 3H), 1.84 (m, 2H), 1.75 (m, 2H), 1.22 (m, 2H); MS 506.1 (M+1), 504.1 (M-1).

#### Example 6b

5-(3-((3-(3-Chloro-phenyl)-propyl)-(2,4-dioxo-1,2,3,4-tetrahydro-quinazoline-6-sulfonyl)-amino)-propyl)-thiophene-2-carboxylic acid

- 15 <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 6.80-7.92 (m, 9H), 3.20 (m, 4H), 2.80 (m, 2H), 1.75-2.00 (m, 4H), 1.20 (m, 2H); MS 594.0 (M-1+Cl).

#### Example 6c

5-(3-((4-(2-Carboxy-benzoylamino)-butane-1-sulfonyl)-(3-(3-chloro-phenyl)-propyl)-amino)-propyl)-thiophene-2-carboxylic acid

- 20 <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.70 (d, 1H, J=6), 7.62 (d, 1H, J=4), 7.55 (d, 1H, J=8), 7.45-7.20 (m, 6H), 6.80-6.90 (m, 10H), 3.22 (m, 4H), 2.70 (m, 2H), 2.60 (m, 2H), 1.80-2.00 (m, 4H), 1.22 (m, 2H); MS 620.1 (M-1).

#### Example 6d

5-(3-((3-(3-Chloro-phenyl)-propyl)-(4-(3,5-dioxo-4,5-dihydro-3H-[1,2,4]triazin-2-yl)-benzenesulfonyl)-amino)-propyl)-thiophene-2-carboxylic acid

- 25 <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.60-7.92 (m, 4H), 6.80 (m, 7H), 3.22 (m, 4H), 2.80 (m, 2H), 2.60 (m, 2H), 1.82 (m, 2H), 1.22 (m, 2H); MS 587.1 (M-1).

#### Example 6e

5-(3-((3-(3-Chloro-phenyl)-propyl)-(2-methoxycarbonyl-benzenesulfonyl)-amino)-propyl)-thiophene-2-carboxylic acid

- 30

-110-

<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.75 (d, 1H, J=4), 7.00-7.70 (m, 8H), 6.85 (d, 1H, J=4), 3.90 (s, 3H), 3.31 (m, 4H), 2.70 (m, 2H), 2.50 (m, 2H), 1.82-2.00 (m, 4H), 1.20 (m, 2H); MS 534.1 (M-1).

Example 6f

5     5-(3-((4-Bromo-benzenesulfonyl)-(3-(3-chloro-phenyl)-propyl)-amino)-propyl)-thiophene-2-carboxylic acid

<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.75 (d, 1H, J=4), 7.00-7.70 (m, 8H), 6.80 (d, 1H, J=4), 3.10 (m, 4H), 2.86 (m, 2H), 2.55 (m, 2H), 1.90 (m, 2H), 1.80 (m, 2H); MS 557.9 (M+1), 555.9 (M-1).

10                   Example 6g

5-(3-((3-(3-Chloro-phenyl)-propyl)-(4-(1,1-dimethyl-propyl)-benzenesulfonyl)-amino)-propyl)-thiophene-2-carboxylic acid

<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.95 (d, 1H, J=4), 7.00-7.80 (m, 8H), 6.80 (d, 1H, J=4), 3.20 (m, 4H), 2.80 (m, 2H), 2.50 (m, 2H), 1.30 (s, 3H), 1.70-1.90 (m, 4H), 1.55 (m, 2H), 0.60 (t, 3H, J=7); MS 548 (M+1).

Example 6h

5-(3-((3-(3-Chloro-phenyl)-propyl)-(3,5-dimethyl-isoxazole-4-sulfonyl)-amino)-propyl)-thiophene-2-carboxylic acid

20     <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 6.95-7.40 (m, 4H), 6.80 (d, 1H, J=8), 6.75 (d, 1H, J=8), 2.91 (m, 2H), 2.60 (s, 3H), 2.40 (m, 2H), 2.20 (s, 3H), 1.72-1.92 (m, 4H), 1.20 (m, 2H); MS 495 (M-1).

Example 6i

5-(3-((3-(3-Chloro-phenyl)-propyl)-(2,5-dimethoxy-benzenesulfonyl)-amino)-propyl)-thiophene-2-carboxylic acid

25     <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.70 (d, 1H, J=4), 7.00-7.50 (m, 7H), 6.80 (d, 1H, J=4), 4.00 (s, 3H), 3.80 (s, 3H), 3.25 (m, 4H), 2.85 (m, 2H), 2.52 (m, 2H), 1.70-2.00 (m, 2H); MS 538.1 (M+1), 536.1 (M-1).

Example 6j

30     5-(3-((3-(3-Chloro-phenyl)-propyl)-(2-fluoro-benzenesulfonyl)-amino)-propyl)-thiophene-2-carboxylic acid

<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.00-8.00 (m, 9H), 6.80 (d, 1H, J=7.2), 3.30 (m, 4H), 2.85 (m, 2H), 2.55 (m, 2H), 1.70-2.00 (m, 4H), 1.20 (m, 2H); MS 494.1 (M-1).

-111-

Example 75-(3-(1-(3-(3-Chloro-phenyl)-propyl)-3-ethyl-ureido)-propyl)-thiophene-2-carboxylic acidStep A: Isocyanate addition

- 5 5-(3-(1-(3-(3-Chloro-phenyl)-propyl)-3-ethyl-ureido)-propyl)-thiophene-2-carboxylic acid tert-butyl ester. A stock solution of 5-(3-(3-(3-chloro-phenyl)-propylamino)-propyl)-thiophene-2-carboxylic acid tert-butyl ester (from Preparation 9, 0.10 g, 0.254 mmol) in 10 mL CH<sub>2</sub>Cl<sub>2</sub> was prepared and 1 mL (0.010 g, 0.0254 mmol) was added to a 1 dram vial. Triethylamine (0.7 mL, 0.051 mmol) and ethyl isocyanate (0.004 g, 10 0.051 mmol) were added and the mixture was stirred overnight at room temperature. The solution was diluted with 2 mL CH<sub>2</sub>Cl<sub>2</sub>. The organic solution was washed with 3 mL of 5.5% aqueous HCl solution (2X) followed by 3 mL saturated bicarbonate solution (2X). The organic layer was dried with MgSO<sub>4</sub> and was concentrated to yield the title compound of Step A (10 mg).

15 Step B: Ester Hydrolysis

Step B was performed in a manner analogous to the method of Step B of Example 6. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.70 (d, 1H, J=4), 7.00-7.40 (m, 4H), 6.80 (d, 1H, J=4), 3.20 (m, 6H), 2.80 (m, 2H), 2.60 (m, 2H), 1.80-2.00 (m, 4H), 1.05 (t, 3H, J=7); MS 409.1 (M+1), 407.1 (M-1).

20 Examples 7a-7j

Examples 7a-7j were prepared from the appropriate starting materials in a manner analogous to the method of Example 7.

Example 7a

25 5-(3-(1-(3-(3-Chloro-phenyl)-propyl)-3-isopropyl-ureido)-propyl)-thiophene-2-carboxylic acid

<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.70 (d, 1H, J=4), 7.00-7.40 (m, 4H), 6.80 (d, 1H, J=4), 3.20 (m, 4H), 2.85 (m, 2H), 2.60 (m, 2H), 1.75-2.00 (m, 4H), 1.05 (d, 6H, J=7); MS 423.1 (M+1), 421.1 (M-1).

Example 7b

30 5-(3-(1-(3-(3-Chloro-phenyl)-propyl)-3-phenyl-ureido)-propyl)-thiophene-2-carboxylic acid



-112-

$^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.75 (d, 1H,  $J=7$ ), 7.00-7.50 (m, 9H), 6.80 (d, 1H,  $J=4$ ), 3.20 (m, 4H), 2.90 (m, 2H), 2.60 (m, 2H), 1.80-2.00 (m, 4H); MS 457.1 (M+1), 455.2 (M-1).

Example 7c

5      5-(3-(1-(3-(3-Chloro-phenyl)-propyl)-3-(3,4-dichloro-phenyl)-ureido)-propyl)-thiophene-2-carboxylic acid

$^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  6.80-7.60 (m, 9H), 3.20 (m, 4H), 2.90 (m, 2H), 2.60 (m, 2H), 1.86-2.00 (m, 4H); MS 527.0 (M+1), 525.0 (M-1).

Example 7d

10      5-(3-(1-(3-(3-Chloro-phenyl)-propyl)-3-propyl-ureido)-propyl)-thiophene-2-carboxylic acid

$^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.70 (d, 1H,  $J=4$ ), 7.00-7.30 (m, 4H), 6.80 (d, 1H,  $J=4$ ), 3.20-3.30 (m, 5H), 2.95 (t, 2H,  $J=7$ ), 2.60 (t, 2H,  $J=7$ ), 1.70-2.00 (m, 4H), 0.95 (t, 3H,  $J=7$ ); MS 423 (M+1), 421 (M-1).

15      Example 7e

5-(3-(3-(4-Chloro-phenyl)-1-(3-(3-chloro-phenyl)-propyl)-ureido)-propyl)-thiophene-2-carboxylic acid

$^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.70 (d, 1H,  $J=4$ ), 7.00-7.30 (m, 8H), 6.80 (d, 1H,  $J=4$ ), 3.22 (m, 4H), 2.90 (m, 2H), 2.65 (m, 2H), 1.69-2.02 (m, 4H); MS 491 (M+1), 489 (M-1).

20

Example 7f

5-(3-(1-(3-(3-Chloro-phenyl)-propyl)-3-(2,3-dichloro-phenyl)-ureido)-propyl)-thiophene-2-carboxylic acid

$^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.70 (bs, 1H), 7.00-7.30 (m, 7H), 6.80 (bs, 1H), 3.20 (m, 4H), 2.80 (m, 2H), 2.60 (m, 2H), 1.75-2.00 (m, 4H); MS 527 (M+1), 525.1 (M-1).

25

Example 7g

5-(3-(1-(3-(3-Chloro-phenyl)-propyl)-3-(3,5-dichloro-phenyl)-ureido)-propyl)-thiophene-2-carboxylic acid

$^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.70 (d, 1H,  $J=4$ ), 7.00-7.30 (m, 7H), 6.80 (d, 1H,  $J=4$ ), 3.20 (m, 4H), 2.80 (m, 2H), 2.60 (m, 2H), 1.70-2.00 (m, 4H); MS 527.1 (M+1), 525.1 (M-1).

30

-113-

Example 7h5-(3-(1-(3-(3-Chloro-phenyl)-propyl)-3-(2,6-difluoro-phenyl)-ureido)-propyl)-thiophene-2-carboxylic acid

5  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.70 (d, 1H,  $J=4$ ), 7.00-7.30 (m, 7H), 6.80 (d, 1H,  $J=4$ ), 3.20 (m, 4H), 2.86 (m, 2H), 2.65 (m, 2H), 1.73-1.95 (m, 4H); MS 493.1 (M+1), 491.1 (M-1).

Example 7i5-(3-(1-(3-(3-Chloro-phenyl)-propyl)-3-(4-fluoro-phenyl)-ureido)-propyl)-thiophene-2-carboxylic acid

10  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.70 (bs, 1H), 7.00-7.60 (m, 8H), 6.80 (bs, 1H), 3.30 (m, 4H), 2.90 (m, 2H), 2.60 (m, 2H), 1.80-2.00 (m, 4H); MS 475.1 (M+1), 473.1 (M-1).

Example 7j5-(3-(3-Butyl-1-(3-(3-chloro-phenyl)-propyl)-ureido)-propyl)-thiophene-2-carboxylic acid

15  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.70 (bs, 1H), 7.00-7.20 (m, 4H), 6.80 (bs, 1H), 3.20 (m, 6H), 2.90 (m, 2H), 2.60 (m, 2H), 1.70-2.00 (m, 4H), 0.95 (t, 3H,  $J=6.8$ ); MS 437.2 (M+1), 435.2 (M-1).

Example 85-(3-(1-(3-(3-Chloro-phenyl)-propyl)-(pyrrolidine-1-carbonyl)-amino)-propyl)-thiophene-2-carboxylic acidStep A: Amide formation

5-(3-(1-(3-(3-Chloro-phenyl)-propyl)-3-ethyl-ureido)-propyl)-thiophene-2-carboxylic acid tert-butyl ester. A stock solution of 5-(3-(3-(3-chloro-phenyl)-propylamino)-propyl)-thiophene-2-carboxylic acid tert-butyl ester (from Preparation 9, 0.10 g, 0.254 mmol) in 10 mL  $\text{CH}_2\text{Cl}_2$  was prepared and 1 mL (0.010 g, 0.0254 mmol) was added to a 1 dram vial. Triethylamine (0.7 mL, 0.051 mmol) and ethyl isocyanate (0.004 g, 0.051 mmol) were added and the reaction was stirred overnight at room temperature. The reaction was diluted with 2 mL  $\text{CH}_2\text{Cl}_2$  and the organic solution was washed with 3 mL of 5.5% aqueous HCl solution (2X) followed by 3 mL saturated bicarbonate solution (2X). The organic layer was dried with  $\text{MgSO}_4$  and concentrated to yield the title compound of Step A (10 mg).

-114-

Step B: Ester Hydrolysis

Step B was performed in a manner analogous to the method of Step B of Example 6.

<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.70 (d, 1H, J=4), 7.00-7.40 (m, 4H), 6.80 (d, 1H, J=4), 3.20 (m, 8H), 2.80 (m, 2H), 2.60 (m, 2H), 1.70-2.00 (m, 8H), 1.20 (m, 4H); MS 435.1 (M+1), 433.2 (M-1).

Examples 8a-8c

Examples 8a-8c were prepared from the appropriate starting material in a manner analogous to the method of Example 8.

Example 8a

10 5-(3-((3-(3-Chloro-phenyl)-propyl)-(morpholine-4-carbonyl)-amino)-propyl)-thiophene-2-carboxylic acid

<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.65 (d, 1H, J=4), 7.00-7.40 (m, 4H), 6.80 (d, 1H, J=4), 3.60 (m, 4H), 3.00-3.20 (m, 8H), 2.80 (m, 2H), 2.60 (m, 2H), 1.70-2.00 (m, 4H); MS 451.1 (M+1), 449.2 (M-1).

15 Example 8b

5-(3-((3-(3-Chloro-phenyl)-propyl)-isopropoxycarbonyl-amino)-propyl)-thiophene-2-carboxylic acid

<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 6.70 (d, 1H, J=4), 7.00-7.30 (m, 4H), 6.80 (d, 1H, J=4), 3.20 (m, 4H), 2.80 (t, 2H, J=6.7), 2.60 (t, 2H, J=6.7), 1.80-2.00 (m, 4H), 1.01 (d, 6H); MS 424 (M+1), 422 (M-1).

Example 8c

5-(3-((3-(3-Chloro-phenyl)-propyl)-propoxycarbonyl-amino)-propyl)-thiophene-2-carboxylic acid

<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.70 (bs, 1H), 7.00-7.30 (m, 4H), 6.80 (bs, 1H), 4.00 (t, 2H, J=6.8), 3.30 (m, 4H), 2.80 (m, 2H), 2.60 (m, 2H), 1.40-2.00 (m, 6H), 0.90 (t, 3H, J=7); MS 424 (M+1), 422.2 (M-1).

Example 9

(3-(((4-Butyl-benzyl)-(pyridine-3-sulfonyl)-amino)-methyl)-phenyl)-acetic acid

Step A: Reductive Amination

30 (3-((4-Butyl-benzylamino)-methyl)-phenyl)-acetic acid methyl ester. A solution of 4-butyl-benzylamine (from Preparation 15, 0.918 g, 6 mmol) in MeOH was added to 4N HCl in dioxane (0.75 mL, 3 mmol) followed by addition of (3-formyl-phenyl)-acetic acid methyl ester (from Preparation 13, 0.534 g, 3.0 mmol). NaCNBH<sub>3</sub> (0.194 mL, 3

-115-

mmol) was added and the reaction was stirred at room temperature overnight. The mixture was diluted with EtOAc and 2N NaOH was added. The organic solution was dried over MgSO<sub>4</sub>, filtered, and concentrated *in vacuo*. The product was purified via flash chromatography (50% hexanes, 50% EtOAc, 0.1% Et<sub>3</sub>N) to afford the title

5 compound of Step A. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.08-7.38 (m, 8H), 3.75 (s, 2H), 3.73 (s, 2H), 3.70 (s, 3H), 3.62 (s, 2H), 2.61 (t, 2H), 1.58 (m, 2H), 1.37 (m, 2H), 0.92 (t, 3H); MS 326 (M+1).

#### Step B: Amide Formation

(3-(((4-Butyl-benzyl)-(pyridine-3-sulfonyl)-amino)-methyl)-phenyl)-acetic acid methyl

10 ester. Step B was performed in a manner analogous to the method of Step B of Example 1 to provide the title compound.

#### Step C: Ester Hydrolysis

(3-(((4-Butyl-benzyl)-(pyridine-3-sulfonyl)-amino)-methyl)-phenyl)-acetic acid. Step C was performed in a manner analogous to the method of Step C of Example 1 to

15 provide the title compound. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 8.99 (bs, 1H), 8.74 (bs, 1H), 7.99 (d, 1H), 7.36 (bs, 1H), 7.20-7.25 (m, 2H), 6.95-7.19 (m, 6H), 4.33 (s, 4H), 3.3-3.54 (s, 2H), 2.54 (m, 2H), 1.54 (m, 2H), 1.32 (m, 2H), 0.91 (t, 3H).

#### Examples 9a-9d

Examples 9a-9d were prepared from the appropriate starting materials in a manner  
20 analogous to the method of Example 9.

#### Example 9a

(3-((Benzenesulfonyl-(4-butyl-benzyl)-amino)-methyl)-phenyl)-acetic acid

<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.83 (d, 2H), 7.46-7.58 (m, 3H), 7.24 (s, 1H), 7.14 (m, 2H), 6.86-6.98 (m, 5H), 4.29 (d, 4H), 3.51 (s, 2H), 2.52 (t, 2H), 1.53 (m, 2H), 1.30 (m, 25 2H), 0.90 (t, 2H); MS 450 (M-1).

#### Example 9b

(3-(((4-Butyl-benzyl)-(thiophene-2-sulfonyl)-amino)-methyl)-phenyl)-acetic acid

<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.53 (m, 2H), 7.16 (m, 2H), 6.89-7.14 (m, 7H), 4.27 (d, 4H), 3.52 (s, 2H), 2.49 (t, 2H), 1.51 (m, 2H), 1.29 (m, 2H), 0.88 (t, 2H); MS 456 (M- 30 1).

#### Example 9c

(3-(((4-Acetylamino-benzenesulfonyl)-(4-butyl-benzyl)-amino)-methyl)-phenyl)-acetic acid

-116-

<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.69 (m, 2H), 7.49 (d, 2H), 7.06-7.23 (m, 6H), 6.91 (d, 1H), 6.68 (s, 1H), 4.30 (d, 4H), 3.44 (s, 2H), 2.54 (t, 2H), 2.17 (s, 3H), 1.54 (m, 2H), 1.29 (m, 2H), 0.89 (t, 2H); MS 507 (M-1).

Example 9d

5     (3-(((Benzo[1,2,5]oxadiazole-4-sulfonyl)-(4-butyl-benzyl)-amino)-methyl)-phenyl)-acetic acid

<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.94 (d, 1H), 7.88 (d, 2H), 7.36 (t, 1H), 7.07 (s, 2H), 6.90-6.96 (m, 6H), 5.3 (d, 4H), 3.46 (s, 2H), 2.46 (t, 2H), 1.47 (m, 2H), 1.26 (m, 2H), 0.88 (t, 2H); MS 4.92 (M-1).

10

Example 10

(3-(((1-Methyl-1H-imidazole-4-sulfonyl)-(4-pyrimidin-2-yl-benzyl)-amino)-methyl)-phenoxy)-acetic acid•HCl

Step A: Reductive Amination

(3-((4-Pyrimidin-2-yl-benzylamino)-methyl)-phenoxy)-acetic acid t-butyl ester. Step A

15     was performed in a manner analogous to the method of Step A of Example 1.

Step B: Amide Formation

(3-(((1-Methyl-1H-imidazole-4-sulfonyl)-(4-pyrimidin-2-yl-benzyl)-amino)-methyl)-phenoxy)-acetic acid tert-butyl ester. Step B was performed in a manner analogous to the method of Step B of Example 1 using triethylamine in place of N,N-

20     diisopropylethylamine as base.

Step C: Ester Hydrolysis

(3-(((1-Methyl-1H-imidazole-4-sulfonyl)-(4-pyrimidin-2-yl-benzyl)-amino)-methyl)-phenoxy)-acetic acid•HCl. A solution of (3-(((1-methyl-1H-imidazole-4-sulfonyl)-(4-pyrimidin-2-yl-benzyl)-amino)-methyl)-phenoxy)-acetic acid tert-butyl ester prepared of Example 10, Step B (0.094 g, 0.17 mmol) in 1N HCl in diethyl ether was stirred for 20 minutes as a precipitate formed. To the mixture was added 1 mL water and 1 mL dioxane and the reaction was stirred for 3 hours. The solvent was removed *in vacuo*, azeotroping with ethanol to yield the title compound as a solid (54 mg). <sup>1</sup>H NMR (400 MHz, CD<sub>3</sub>OD) δ 9.09 (m, 2H), 8.95 (bs, 1H), 8.24 (d, 2H), 8.04 (s, 1H), 7.71 (s, 1H), 7.44 (d, 2H), 7.13 (m, 1H), 6.82 (d, 1H), 6.76 (d, 1H), 6.69 (s, 1H), 4.61 (s, 2H), 4.54 (s, 2H), 4.46 (s, 2H), 3.92 (s, 3H); MS 494 (M+1).

25

30

-117-

Preparation 1

7-Amino-heptanoic acid methyl ester hydrochloride. A solution of 7-amino-heptanoic acid (3.0 g, 21.0 mmol), in 25 mL MeOH and 2.4 mL concentrated HCl was heated at reflux for 4 hours and was stirred at room temperature for 60 h. The mixture was  
5 concentrated *in vacuo* to afford the title compound (3.3 g). <sup>1</sup>H NMR (400 MHz, CD<sub>3</sub>OD) δ 3.62 (s, 3H), 2.89 (m, 2H), 2.31 (t, 2H), 1.62 (m, 4H), 1.37 (m, 4H).

Preparation 2

Pyridine-3-sulfonyl chloride hydrochloride. The title compound was prepared using the method described by Karaman, R. and coworkers J. Am. Chem. Soc. 114, 12,  
10 1992, 4889-4898.

Preparation 3

3-(3-Chloro-phenyl)-propionaldehyde. A solution of 1-chloro-3-iodobenzene (9.63 g, 40.38 mmol), allyl alcohol (5.86 g, 100.96 mmol), sodium bicarbonate (8.48 g, 100.96 mmol), tetrabutylammonium chloride (11.22 g, 40.38 mmol), and Pd(OAc)<sub>2</sub> (317 mg, 1.413 mmol) in 25 mL DMF was stirred at 50°C for 16 h. The mixture was cooled to  
15 room temperature, diluted with water, and the aqueous solution was washed with EtOAc. The organic solution was washed with water followed by brine, dried over MgSO<sub>4</sub>, filtered and concentrated *in vacuo*. The product was purified via flash chromatography on silica gel (9:1 hexanes:EtOAc) to afford the title compound as an  
20 oil (5.04 g).

Preparation 45-(3-Oxo-propyl)-thiophene-2-carboxylic acid tert-butyl esterStep A: Ester Formation

5-Bromo-thiophene-2-carboxylic acid tert-butyl ester. To a solution of anhydrous  
25 MgSO<sub>4</sub> (11.60 g, 96.4 mmol) in 100 mL CH<sub>2</sub>Cl<sub>2</sub> was added concentrated H<sub>2</sub>SO<sub>4</sub> (1.45 mL, 24.1 mmol) and the mixture was stirred for 15 minutes followed by addition of 5-bromo-thiophene-2-carboxylic acid (5.0 g, 24.1 mmol). After stirring for 1 minute, tert-butanol (11.6 g, 20 mmol) was added and the reaction was stirred at room  
30 temperature for 16 h. The reaction was quenched with saturated NaHCO<sub>3</sub>. The layers were separated, the aqueous layer was extracted with CH<sub>2</sub>Cl<sub>2</sub>, and the combined organic layers were dried over MgSO<sub>4</sub>. The organic solution was concentrated to give a clear oil which was purified via medium pressure

-118-

chromatography (3% EtOAc in hexanes) to afford the title compound of Step A (4.97 g). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.45 (d, 1H), 7.02 (d, 1H), 1.54 (s, 9H).

Step B: Aldehyde formation

- 5-(3-Oxo-propyl)-thiophene-2-carboxylic acid tert-butyl ester. To a solution of 5-bromo-thiophene-2-carboxylic acid tert-butyl ester prepared of the method of Preparation 4, Step A (0.50 g, 1.89 mmol) in 5 mL DMF was added allyl alcohol (0.51 mL, 7.57 mmol) followed by NaHCO<sub>3</sub> (0.397 g, 4.72 mmol), tetrabutylammonium chloride (0.525 g, 1.89 mmol), and palladium acetate (0.021 g, 0.094 mmol). The reaction was placed in an oil bath heated to 65°C and was heated to 90°C for 2 h.
- 10 The mixture was diluted with EtOAc and 25 mL water and the solids were removed by filtration through Celite®. The layers were separated, and the organic solution was washed with water (4x), dried over MgSO<sub>4</sub> and concentrated to a dark yellow oil which was purified via medium pressure chromatography (7:1 hexanes:EtOAc) to afford the title compound (0.190 g). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 9.80 (s, 1H), 7.51 (d, 1H), 6.78 (d, 1H), 3.14 (t, 2H), 2.86 (t, 2H), 1.54 (s, 9H).
- 15

Preparation 5

5-(3-Amino-propyl)-thiophene-2-carboxylic acid methyl ester

Step A

- 5-(3-tert-Butoxycarbonylamino-prop-1-ynyl)-thiophene-2-carboxylic acid methyl ester.
- 20 A mixture of prop-2-ynyl-carbamic acid tert-butyl ester (from Preparation 41, 1.67 g, 0.011 mmol), 5-bromo-thiophene-2-carboxylic acid methyl ester (2.50 g, 0.011 mmol), tetrakis(triphenylphosphine(0) palladium (0.622 g, 0.0538 mmol), CuI (0.102 g, 0.538 mmol) and triethylamine (1.57 mL, 0.011 mmol) in 50 mL acetonitrile was heated at reflux for 16 h. The reaction was cooled to room temperature, diluted with 75 mL
- 25 EtOAc, washed with 5.5% HCl, water and brine, dried over MgSO<sub>4</sub>, filtered and concentrated *in vacuo* to an oil. The product was purified via flash chromatography (9:1 to 4:1 hexanes:EtOAc) to afford the title compound of Step A as an oil (2.06 g). MS 313 (M+18).

Step B

- 30 5-(3-tert-Butoxycarbonylamino-propyl)-thiophene-2-carboxylic acid methyl ester. A solution of 5-(3-tert-butoxycarbonylamino-prop-1-ynyl)-thiophene-2-carboxylic acid methyl ester prepared of Preparation 5, Step A (2.06 g) and 10% Pd/C (1.03 g) in 50 mL MeOH was hydrogenated on a Parr shaker at 50 psi H<sub>2</sub> for 16 h. The reaction

-119-

was filtered through Celite® with the aid of MeOH and the filtrate was concentrated *in vacuo* to afford the title compound of Step B as a solid (1.93 g). MS 317 (M+18).

#### Step C

5-(3-Amino-propyl)-thiophene-2-carboxylic acid methyl ester. A solution of 5-(3-tert-butoxycarbonylamino-propyl)-thiophene-2-carboxylic acid methyl ester prepared of Preparation 5, Step B (0.118 g, 0.5 mmol) in 50 mL MeOH was cooled to 0°C and was saturated with HCl (g). The reaction was stirred at room temperature for 90 minutes. The solution was concentrated to a solid which was partitioned between EtOAc and saturated NaHCO<sub>3</sub>. The layers were separated, and the combined organic layers were washed with brine, dried over MgSO<sub>4</sub>, filtered and concentrated *in vacuo* to afford the title compound as an oil (399 mg). MS 200 (M+1).

#### Preparation 6

5-(3-Amino-propyl)-furan-2-carboxylic acid methyl ester hydrochloride salt. The compound of Preparation 6 was prepared from the appropriate starting materials in a manner analogous to the method of Preparation 5 with the following exceptions: (1) the hydrogenation performed in Step B was carried out for 5.5 h; and (2) in Step C, the reaction was stirred for 16 h at room temperature and was concentrated *in vacuo* to provide the title compound as the hydrochloride salt.

#### Preparation 7

5-(3-Amino-propyl)-thiophene-2-carboxylic acid tert-butyl ester

#### Step A

Prop-2-ynyl-carbamic acid benzyl ester. To a solution of propargylamine (6.4 g, 71.2 mmol) in pyridine (100 mL) was added benzylchloroformate (13.37 g, 78.2 mmol) in 100 mL CH<sub>2</sub>Cl<sub>2</sub> over 0.5 h. The reaction was stirred for 16 h and the volatiles were removed *in vacuo*. The residue was dissolved in EtOAc and the organic solution was washed with water (2x). The organic solution was washed with dilute aqueous HCl followed by saturated NaHCO<sub>3</sub>. The organic solution was dried over MgSO<sub>4</sub>, filtered, and concentrated *in vacuo* to provide the title compound of Step A (4.43 g).

#### Step B

5-(3-Benzyloxycarbonylamino-prop-1-ynyl)-thiophene-2-carboxylic acid tert-butyl ester. The title compound of Step B was prepared from the appropriate starting material in a manner analogous to the method used in Step A of Preparation 5.



-120-

Step C

5-(3-Amino-propyl)-thiophene-2-carboxylic acid tert-butyl ester. To a solution of 5-(3-benzyloxycarbonylamino-prop-1-ynyl)-thiophene-2-carboxylic acid tert-butyl ester prepared of Preparation 7, Step B (1.0 g, 2.69 mmol) in 15 mL MeOH and 2.69 mL  
5 1N HCl (aq) was added Pd(OH)<sub>2</sub>. The mixture was hydrogenated on a Parr shaker at 45 psi H<sub>2</sub> for 16 h. The mixture was filtered through Celite<sup>®</sup>, the catalyst was replaced, and the reaction was shaken for another 6 h. The mixture was filtered through Celite<sup>®</sup> and concentrated *in vacuo*. The residue was chased with CCl<sub>4</sub> and was triturated with Et<sub>2</sub>O. The product was isolated as a solid (360 mg).

10

Preparation 8

5-(3-(3-(3-Chloro-phenyl)-propylamino)-propyl)-thiophene-2-carboxylic acid methyl ester. A solution of 5-(3-amino-propyl)-thiophene-2-carboxylic acid methyl ester (from Preparation 5, Step C, 0.118 g, 0.5 mmol) and N,N-diisopropylethylamine (0.071 g, 0.55 mmol) in 10 mL MeOH was stirred at room temperature for 30 minutes and 3-(3-chloro-phenyl)-propionaldehyde (from Preparation 3, 0.093 g, 0.55 mmol) was added.  
15 The mixture was stirred for 90 minutes. The reaction was cooled to 0°C, NaBH<sub>4</sub> (30.3 mg, 0.801 mmol) was added and the mixture was stirred for 30 minutes. The reaction was quenched with 1:1 NaHCO<sub>3</sub>:H<sub>2</sub>O and was washed with CH<sub>2</sub>Cl<sub>2</sub>. The CH<sub>2</sub>Cl<sub>2</sub> extracts were washed with brine, dried over MgSO<sub>4</sub>, filtered, and  
20 concentrated *in vacuo* to afford the title compound as an oil (171 mg). MS 352 (M+1).

Preparations 9-10

The compounds of Preparations 9 and 10 were prepared from the appropriate starting materials in a manner analogous to the method of Preparation 8.

Preparation 9

25 5-(3-(3-(3-Chloro-phenyl)-propylamino)-propyl)-thiophene-2-carboxylic acid tert-butyl ester

Preparation 10

5-(3-(3-(3-Chloro-phenyl)-propylamino)-propyl)-furan-2-carboxylic acid methyl ester  
MS 336 (M+1).

30

Preparation 11

(3-Formyl-phenoxy)-acetic acid methyl ester. A mixture of (3-formyl-phenoxy)-acetic acid (3.6 g, 20.0 mmol), potassium carbonate (3.30 g, 23.9 mmol) and methyl iodide (1.86 g, 30.0 mmol) in 25 mL DMF was heated to 110°C for 2 hours and was stirred

-121-

at room temperature for 16 h. The mixture was diluted with water and the aqueous solution was extracted with EtOAc. The organic solution was washed with water, dried over  $\text{MgSO}_4$ , filtered and concentrated *in vacuo*. The product was purified via silica gel chromatography (4:1 hexanes: EtOAc) to afford the title compound as a pale yellow oil (3.4 g).  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  9.94 (s, 1H), 7.48 (m, 2H), 7.33 (s, 1H), 7.23 (m, 1H), 4.68 (s, 2H), 3.79 (s, 3H).

#### Preparation 12

##### 3-(3-Chloro-phenyl)-propylamine

#### STEP A

10 3-(3-Chloro-phenyl)-acrylamide. A solution of 3-(3-chloro-phenyl)-acrylic acid (Aldrich, 15.0 g, 82.15 mmol) in 50 mL thionyl chloride was heated at reflux for 30 minutes. The excess thionyl chloride was removed via distillation at atmospheric pressure. The residue was azeotroped with benzene *in vacuo* to give 17.288 g of an orange oil. The oil was dissolved in 25 mL  $\text{CH}_2\text{Cl}_2$  and the solution was added slowly to liquid  $\text{NH}_3$  (20 mL, 80.07 mmol) in  $\text{CHCl}_3$  (50 mL) at  $-78^\circ\text{C}$ . The resulting suspension was warmed to room temperature and was concentrated *in vacuo* to afford the title compound of Step A as a gray solid (19.38 g).  $^1\text{H}$  NMR (400 MHz,  $\text{CD}_3\text{OD}$ )  $\delta$  7.57 (s, 1H), 7.45 (m, 2H), 7.36 (m, 1H), 6.64 (d, 1H); MS 182 (M+1), 180 (M-1).

20

#### STEP B

3-(3-Chloro-phenyl)-propylamine. A 1.0 M solution of  $\text{LiAlH}_4$  in THF (6.0 mL) was added dropwise to a suspension of 3-(3-chloro-phenyl)-acrylamide prepared of Preparation 12, Step A (1.0 g, 5.51 mmol) in 30 mL THF at  $0^\circ\text{C}$ . The reaction was warmed to room temperature and was stirred for 5 h. An additional 4 mL of 1 M  $\text{LiAlH}_4$  was added and the reaction was stirred for 18 h. An additional 2 mL of 1 M  $\text{LiAlH}_4$  was added and the reaction was stirred for 24 h. The reaction mixture was quenched by dropwise addition of water. The mixture was concentrated *in vacuo* to remove THF and was diluted with water. The aqueous solution was extracted with EtOAc. The organic solution was washed with water, dried over  $\text{MgSO}_4$ , filtered and concentrated *in vacuo*. The residue was dissolved in  $\text{CHCl}_3$  and the organic solution was washed with 1M HCl. The aqueous solution was basified to pH 11 with 1M NaOH and the product was extracted into  $\text{CHCl}_3$ . The organic solution was dried over  $\text{MgSO}_4$ , filtered and concentrated *in vacuo* to afford the title compound as a

-122-

yellow oil (0.134 g).  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.20-7.22 (m, 3H), 7.16 (m, 1H), 2.74 (t, 2H), 2.61 (t, 2H), 1.74 (m, 2H); MS 170 (M+1).

### Preparation 13

#### (3-Formyl-phenyl)-acetic acid methyl ester

5

#### Step A

(3-Cyano-phenyl)-acetic acid methyl ester. Nitrogen was bubbled through a mixture of (3-bromo-phenyl)-acetic acid methyl ester (22.85 g, 99.78 mmol),  $\text{Zn}(\text{CN})_2$  (7.25 g, 61.75 mmol), and DMF (100 mL) for about 5 minutes followed by addition of tetrakis(triphenylphosphine(0) palladium (4.60 g, 3.98 mmol). The mixture was heated for 3 h at  $80^\circ\text{C}$  and was cooled to room temperature. Aqueous 2N  $\text{NH}_4\text{OH}$  was added and the product was extracted into EtOAc (3x). The organic solution was washed with 2N  $\text{NH}_4\text{OH}$  (2x) followed by brine (2x). The organic solution was dried ( $\text{MgSO}_4$ ), filtered, and concentrated *in vacuo*. Purification by flash chromatography (6:1 hexanes:EtOAc) provided the title compound of Step A as an oil (15.19 g).  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.57-7.41 (m, 4H), 3.706 (s, 3H), 3.703 (s, 2H).

15

#### Step B

(3-Formyl-phenyl)-acetic acid methyl ester. A mixture of (3-cyano-phenyl)-acetic acid methyl ester prepared of Preparation 13, Step A (1.56 g, 8.91 mmol), aluminum-nickel alloy (1.63 g) and 75% formic acid (25 mL) was heated at reflux for 1.75 h. The mixture was cooled to room temperature and the solids were removed by filtration through Celite<sup>®</sup> with the aid of boiling EtOH. Water was added, and the aqueous solution was washed with  $\text{CH}_2\text{Cl}_2$  (3x). Aqueous saturated  $\text{NaHCO}_3$  was carefully added to the organic solution until the pH was about 8-9. The organic solution was washed with brine, dried over  $\text{MgSO}_4$ , and concentrated. Purification by flash chromatography (5:1 hexanes:EtOAc) provided the title compound as a clear and colorless oil (870 mg).  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  9.98 (s, 1H), 7.77 (m, 2H), 7.55-7.46 (m, 2H), 3.68 (s, 5H).

20

25

### Preparation 14

(3-((Pyridine-3-sulfonylamino)-methyl)-phenyl)-acetic acid methyl ester. To a solution of (3-aminomethyl-phenyl)-acetic acid methyl ester hydrochloride (from Preparation 18, 0.56 g) and diisopropylamine (2.2 mL) in 10 mL dichloromethane was added pyridine-3-sulfonyl chloride (from Preparation 2, 0.601 g, 2.83 mmol) and the reaction was stirred at room temperature for 16 h. Aqueous 1N HCl was added and the

30

-123-

solution was washed with  $\text{CH}_2\text{Cl}_2$ . The organic solution was washed with saturated  $\text{NaHCO}_3$ , dried over  $\text{MgSO}_4$ , filtered and concentrated *in vacuo* to afford the title compound. Purification via flash chromatography on silica gel (2:1 hexanes:EtOAc) afforded the title compound as a white solid.  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  8.91 (s, 1H), 8.71 (d, 1H), 8.04 (d, 1H), 7.37 (m, 1H), 7.05-7.24 (m, 4H), 5.87 (bs, 1H), 4.14 (s, 2H), 3.62 (s, 3H), 3.52 (s, 2H).

#### Preparation 15

##### Method A

4-Butylbenzylamine. A solution of 4-butylbenzonitrile (3.63 g, 22.8 mmol) in THF (10 mL) was placed in a three-neck round bottom flask equipped with a Vigreux column and short-path distillation head. The solution was heated to reflux and  $\text{BH}_3$ -methyl sulfide complex (2.0 M in THF, 15 mL, 30 mmol) was added dropwise over 15 minutes. Methyl sulfide was distilled off from the reaction mixture over 1 h and the solution was cooled to room temperature. Aqueous HCl (6N, 25 mL) was added slowly via an addition funnel and the mixture was heated at reflux for 30 minutes. The reaction was cooled to  $0^\circ\text{C}$  and NaOH (7.0 g) was added portionwise. The aqueous solution was washed with EtOAc (3x) and the organic solution was dried ( $\text{MgSO}_4$ ), filtered, and concentrated to provide the title compound of Method A (4.01 g).  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.34 (m, 2H), 7.24 (m, 2H), 4.04 (s, 2H), 2.62 (t, 2H), 1.58 (m, 2H), 1.34 (m, 2H), 0.92 (t, 3H).

##### Method B

4-Butylbenzylamine hydrochloride. A solution of 4-butylbenzonitrile (30.09 g) in EtOH (380 mL) and HCl (4N in dioxane, 50 mL, 200 mmol) was hydrogenated at 50 psi on a Parr shaker in the presence of 10% palladium on carbon (6.09 g). The catalyst was removed via filtration through Celite<sup>®</sup> and the solution was concentrated *in vacuo*. The residue was suspended in  $\text{Et}_2\text{O}$  and filtered to provide 4-butylbenzylamine hydrochloride as an off-white solid (32.47 g).  $^1\text{H}$  NMR (400 MHz,  $\text{CD}_3\text{OD}$ )  $\delta$  7.33 (d, 2H), 7.22 (d, 2H), 4.04 (s, 2H), 2.60 (t, 2H), 1.56 (m, 2H), 1.31 (m, 2H), 0.89 (t, 3H).

Using the appropriate starting materials, the compounds of Preparations 16-18 were prepared in a manner analogous to the method of Preparation 15.

PREPARATION 16

2-(3,5-Dichloro-phenoxy)-ethylamine. The title compound was prepared following Method A of Preparation 15.

PREPARATION 17

- 5    2-(3-Chloro-phenoxy)-ethylamine. The title compound was prepared following Method A of Preparation 15.

PREPARATION 18

- 10    (3-Aminomethyl-phenyl)-acetic acid methyl ester hydrochloride. The title compound was prepared from (3-cyano-phenyl)-acetic acid methyl ester (from Preparation 13, Step A) using the procedure described for Preparation 15, Method B except the hydrogenation was performed in MeOH. The catalyst was removed via filtration and the organic solution was concentrated *in vacuo*. The resulting solid was stirred in EtOAc and filtered to provide the title compound as a white solid. <sup>1</sup>H NMR (400 MHz, CD<sub>3</sub>OD)  $\delta$  7.42-7.32 (m, 4H), 4.09 (s, 2H), 3.69 (s, 2H), 3.67 (s, 3H); MS 180 (M+1).

15

Preparation 19*trans*-1-(3-Bromo-propenyl)-3,5-dichloro-benzeneSTEP A

- 20    1-(3,5-Dichloro-phenyl)-prop-2-en-1-ol. A solution of 3,5-dichlorobenzaldehyde (7.5 g, 43 mmol) in THF (75 mL) was cooled to 0°C and vinylmagnesium bromide (1M in THF, 48 mL, 48 mmol) was added dropwise. The reaction was warmed to room temperature and was stirred for 16 h. Aqueous HCl (1N) and EtOAc were added. The aqueous solution was washed with EtOAc and the organic solution was dried (MgSO<sub>4</sub>), filtered, and concentrated. The residue was used in the next step without further purification.

25

STEP B

- 30    The residue prepared in Step A was dissolved in Et<sub>2</sub>O and HBr gas was slowly bubbled into the solution for about 15 minutes. The reaction was stirred at room temperature for 24 h and water and EtOAc were added. The aqueous solution was extracted with EtOAc and the organic solution was dried (MgSO<sub>4</sub>), filtered, and concentrated. Purification by flash chromatography (hexanes) provided the title compound of Preparation 19 (6.91 g). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  7.24 (s, 3H), 6.53 (d, 1H), 6.40 (m, 1H), 4.10 (m, 2H).

-125-

Preparation 20(3-Aminomethyl-phenoxy)-acetic acid tert-butyl esterStep A

- (3-Formyl-phenoxy)-acetic acid tert-butyl ester. To a solution of 3-hydroxybenzaldehyde (5.00 g, 40.9 mmol) in DMF (40 mL) was added 1M potassium tert-butoxide in tert-butanol (40.9 mL, 40.9 mmol). The reaction was stirred for 2 minutes and tert-butyl bromoacetate (6.61 mL, 40.9 mmol) was added. The reaction was stirred for 1 hour and was quenched with 200 mL water. The product was extracted into EtOAc and the organic solution was washed with water, dried over
- 10 MgSO<sub>4</sub>, filtered, and concentrated *in vacuo*. Purification via flash chromatography on silica gel (9:1 hexanes:EtOAc) afforded the title compound of Step A as a clear oil (3.53 g). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 9.94 (s, 1H), 7.48 (m, 2H), 7.32 (s, 1H), 7.21 (m, 1H), 4.56 (s, 2H), 1.45 (s, 9H).

Step B

- 15 (3-(Hydroxyimino-methyl)-phenoxy)-acetic acid tert-butyl ester. To a solution of (3-formyl-phenoxy)-acetic acid tert-butyl ester prepared of Preparation 20, Step A (2.05 g, 8.68 mmol) in MeOH (30 mL) was added NH<sub>2</sub>OH•HCl (0.66 g, 9.54 mmol) and pyridine (3.5 mL, 43.4 mmol) and the reaction was stirred for 2 hours. The MeOH was removed *in vacuo* and the residue was diluted with EtOAc and 1N HCl. The layers
- 20 were separated and the aqueous solution was washed with EtOAc. The combined organic layers were dried over MgSO<sub>4</sub>, filtered and concentrated *in vacuo* to afford the title compound of Step B (1.99 g). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 8.07 (s, 1H), 7.23-7.28 (m, 2H), 7.12 (m, 1H), 6.93 (d, 1H), 4.51 (s, 2H), 1.46 (s, 9H).

Step C

- 25 (3-Aminomethyl-phenoxy)-acetic acid tert-butyl ester. To a solution of (3-(hydroxyimino-methyl)-phenoxy)-acetic acid tert-butyl ester prepared of Preparation 20, Step B (2.25 g, 5.96 mmol) in EtOH (10 mL) was added Raney Nickel (about 1 g, washed with water followed by EtOH) in 100 mL EtOH. Additional EtOH (90 mL) was required for the transfer. Ammonium hydroxide (10 mL) was added and the mixture
- 30 was shaken under 45 psi of H<sub>2</sub> for 4 hours. The catalyst was removed via filtration through Celite® and the solution was concentrated to a clear oil. Purification via flash chromatography on silica gel (96.5/3.5/0.1 to 9/1/0.1 CH<sub>2</sub>Cl<sub>2</sub>/MeOH/NH<sub>4</sub>OH) afforded the title compound as a yellow oil. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.23 (m, 1H), 6.92

-126-

(m, 2H), 6.72 (d, 1H), 4.50 (s, 2H), 3.82 (s, 2H), 1.96 (m, 2H), 1.46 (s, 9H); MS 238 (M+1).

#### Preparation 21

##### 4-Pyrimidin-2-yl-benzaldehyde

- 5 A solution of 2-bromopyrimidine (1.00 g, 6.3 mmol) and tetrakis(triphenylphosphine) palladium (0.218 g, 0.189 mmol) in ethylene glycol dimethyl ether (30 mL) was stirred at room temperature for 10 minutes. A solution of 4-formylbenzene boronic acid (1.14 g, 7.61 mmol) and sodium bicarbonate (1.58 g, 18.9 mmol) in 15 mL water was added and the reaction was heated at reflux for 16 h. The mixture was diluted with
- 10 water and CH<sub>2</sub>Cl<sub>2</sub>. The layers were separated, and the aqueous solution was washed with CH<sub>2</sub>Cl<sub>2</sub>. The combined organic layers were dried over MgSO<sub>4</sub>, filtered, and concentrated *in vacuo*. The residue was purified via flash chromatography on silica gel (10% to 30% hexanes in EtOAc) to afford the title compound (0.979 g). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 10.11 (s, 1H), 8.83 (s, 2H), 8.82 (s, 1H), 7.98 (s, 2H), 7.23
- 15 (s, 2H).

#### Preparations 22-27

Preparations 22-27 were prepared from the appropriate starting materials in a manner analogous to the method of Preparation 21.

#### Preparation 22

##### 4-Pyridin-2-yl-benzaldehyde

20 <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 10.09 (s, 1H), 8.72 (s, 1H), 8.16 (s, 2H), 7.95 (s, 2H), 7.79 (s, 2H), 7.29 (m, 1H); MS 184 (M+1).

#### Preparation 23

##### 4-Pyridin-3-yl-benzaldehyde

- 25 <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 10.04 (s, 1H), 8.88 (s, 1H), 8.64 (s, 1H), 7.97 (s, 2H), 7.91 (m, 1H), 7.75 (m, 2H), 7.39 (m, 1H); MS 184 (M+1).

#### Preparation 24

##### 4-Pyridin-4-yl-benzaldehyde

- 30 <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 10.03 (s, 1H), 8.70 (s, 2H), 7.99 (s, 2H), 7.79 (s, 2H), 7.52 (s, 2H); MS 184 (M+1).

#### Preparation 25

##### 4-Thiazol-2-yl-benzaldehyde

MS 189 (M+).

-127-

Preparation 264-Pyrimidin-5-yl-benzaldehyde

<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 10.03 (s, 1H), 9.26 (s, 1H), 9.00 (s, 2H), 8.03 (m, 2H), 7.76 (m, 2H).

5

Preparation 274-Pyrazin-2-yl-benzaldehyde

<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 10.03 (s, 1H), 9.10 (s, 1H), 8.69 (s, 1H), 8.59 (s, 1H), 8.21 (d, 2H), 8.03 (d, 2H).

Preparation 28

- 10 1-(2-Bromo-ethoxy)-3,5-dichloro-benzene. To a solution of NaOH (2.45 g, 61.3 mmol) in water (20 mL) was added 3,5-dichlorophenol (5 g, 30.7 mmol). The solution was heated at reflux for 1 h and was cooled to room temperature. 1,2-Dibromoethane (11.52 g, 61.3 mmol) was added and the reaction was heated at reflux for 24 h. The cooled solution was diluted with EtOAc and the organic solution
- 15 was washed sequentially with HCl (1N, 1x), water (1x), and brine (1x). The organic solution was dried (MgSO<sub>4</sub>), filtered, and concentrated. Purification by flash chromatography (hexanes to 5% EtOAc in hexanes ) provided the title compound (3.79 g). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 6.98 (m, 1H), 6.82 (m, 2H), 4.25 (t, 2H), 3.61 (t, 2H).

20

Preparation 29

1-(2-Bromo-ethoxy)-3-chlorobenzene. The compound of Preparation 29 was prepared from the appropriate starting materials in a manner analogous to the method of Preparation 28.

Preparation 30

25

4-[(1-Acetyloxy)-hexyl]-benzyl bromideSTEP A: Grignard Reaction and Protection

- 4-[(1-Acetyloxy)-hexyl]-toluene. Pentylmagnesium bromide (2.0 M in Et<sub>2</sub>O, 25 mL, 50 mmol) was added slowly to *p*-tolylbenzaldehyde (5.0 mL, 42.4 mmol) in THF (50 mL) at 0°C. The reaction was warmed to room temperature and was stirred for 3 h.
- 30 Aqueous 1N HCl was added and the aqueous solution was extracted with EtOAc. The organic solution was washed with brine, dried over MgSO<sub>4</sub>, filtered, and concentrated. The residue was dissolved in pyridine (35 mL) and Ac<sub>2</sub>O (10 mL) was added. The reaction was stirred for 24 h and was diluted with water. The product



-128-

was extracted into EtOAc (3x) and the organic solution was washed with 1N HCl followed by brine, dried over MgSO<sub>4</sub>, filtered, and concentrated. The product was purified by flash chromatography (10% EtOAc/hexanes) to afford 4-((1-acetyloxy)-hexyl)-toluene (2.082 g). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.12-7.28 (m, 4H), 5.69 (t, 1H), 2.33 (s, 3H), 2.04 (s, 3H), 1.88 (m, 1H), 1.74 (m, 1H), 1.27 (m, 6H), 0.86 (m, 3H); MS 252 (M+18).

#### STEP B: Benzylic Bromination

A mixture of 4-[(1-acetyloxy)-hexyl]-toluene prepared of Preparation 30, Step A (2.082 g, 8.89 mmol), N-bromosuccinimide (1.58 g, 8.89 mmol), and catalytic 2,2'-azobisisobutyronitrile in carbon tetrachloride (30 mL) was heated at reflux for 2 h. The reaction was cooled and was washed with aqueous NaHCO<sub>3</sub> (saturated), dried over MgSO<sub>4</sub>, filtered, and concentrated. The product was purified by flash chromatography (5% EtOAc/hexanes) to afford the title compound of Preparation 30 (2.67 g). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.34-7.40 (m, 4H), 5.70 (t, 1H), 4.47 (s, 2H), 2.06 (s, 3H), 1.86 (m, 1H), 1.73 (m, 1H), 1.27 (m, 6H), 0.85 (m, 3H).

#### Preparation 31

1-Methyl-1H-indole-2-carbaldehyde. The title compound can be prepared using the method described by Comins and coworkers in J. Org. Chem., 52, 1, 104-9, 1987.

#### Preparation 32

5-Phenyl-furan-2-carbaldehyde. The title compound can be prepared using the method described by D'Auria and coworkers in Heterocycles, 24, 6, 1575-1578, 1986.

#### Preparation 33

4-Phenethylsulfanyl-benzaldehyde. The title compound can be prepared using the method described by Clark and coworkers in EP 332331.

#### Preparation 34

3-Hydroxy-4-propoxy-benzaldehyde. The title compound can be prepared using the method described by Beke in Acta Chim. Acad. Sci. Hung., 14, 325-8, 1958.

#### Preparation 35

4-Formyl-N-methyl-benzenesulfonamide. The title compound can be prepared using the method described by Koetschet in Helv. Chim. Acta., 12, 682, 1929.

-129-

Preparation 36

4-Chloro-thiophene-2-carbaldehyde. The title compound can be prepared using the method described by Raggon and coworkers in Org. Prep. Proced. Int.; EN, 27, 2, 233-6, 1995.

5

Preparation 37

4-Cyclohexyl-benzylamine. The title compound can be prepared using the method described by Meglio and coworkers in Farmaco Ed. Sci.; IT; 35, 3, 191-202, 1980.

Preparation 38

10 4-Imidazol-1-yl-benzaldehyde. The title compound can be prepared using the method described by Sircar and coworkers in J. Med. Chem. 30, 6, 1023-9, 1987.

Preparation 39

4-(2-Oxo-pyrrolidin-1-yl)-benzaldehyde. The title compound can be prepared using the method described by Kukalenko in Chem. Heterocycl. Compd. (Engl. Transl.), 8, 43, 1972.

15

Preparation 40

2-(3-Chloro-phenylsulfanyl)-ethylamine. The title compound can be prepared using the method described by Elz and coworkers in Fed. Rep. Ger. Sci. Pharm., 56, 4, 229-234, 1988.

Preparation 41

20 Prop-2-ynyl-carbamic acid t-butyl ester. The title compound can be prepared using the method described in J. Chem. Soc. Perkin Trans. I, 1985, 2201-2208.

Unless otherwise specified, all reactions were performed under an inert atmosphere such as nitrogen (N<sub>2</sub>).

25 NMR spectra were recorded on a Varian XL-300 (Varian Co., Palo Alto, California), a Bruker AM-300 spectrometer (Bruker Co., Billerica, Massachusetts) or a Varian Unity 400 at about 23°C at 300 or 400 MHz for proton and 75.4 MHz for carbon nuclei. Chemical shifts are expressed in parts per million downfield from trimethylsilane. The peak shapes are denoted as follows: s, singlet; d, doublet; t, triplet; q, quartet; m, multiplet; bs=broad singlet. Resonances designated as exchangeable did not appear  
30 in a separate NMR experiment where the sample was shaken with several drops of D<sub>2</sub>O in the same solvent. Atmospheric pressure chemical ionization (APCI) mass spectra were obtained on a Fisons Platform II Spectrometer. Chemical ionization

-130-

mass spectra were obtained on a Hewlett-Packard 5989 instrument (Hewlett-Packard Co., Palo Alto, California) (ammonia ionization, PBMS). Where the intensity of chlorine or bromine-containing ions are described the expected intensity ratio was observed (approximately 3:1 for  $^{35}\text{Cl}/^{37}\text{Cl}$ -containing ions) and 1:1 for  $^{79}\text{Br}/^{81}\text{Br}$ -containing ions) and the intensity of only the lower mass ion is given.

Column chromatography was performed with either Baker Silica Gel (40  $\mu\text{m}$ ) (J.T. Baker, Phillipsburg, N.J.) or Silica Gel 60 (EM Sciences, Gibbstown, N.J.) in glass columns under low nitrogen pressure. Radial Chromatography was performed, using a Chromatotron<sup>®</sup> (model 7924T, Harrison Research). Medium pressure chromatography was performed on a Flash 40 Biotage System (Biotage Inc, Dyax Corp., Charlottesville, Virginia). Unless otherwise specified, reagents were used as obtained from commercial sources. Dimethylformamide, 2-propanol, acetonitrile, methanol, tetrahydrofuran, and dichloromethane, when used as reaction solvents, were the anhydrous grade supplied by Aldrich Chemical Company (Milwaukee, Wisconsin). The terms "concentrated" and "coevaporated" refer to removal of solvent at water aspirator pressure on a rotary evaporator with a bath temperature of less than 45°C. Reactions conducted at "0-20°C" or "0-25°C" were conducted with initial cooling of the vessel in an insulated ice bath which was allowed to warm to room temperature over several hours. The abbreviation "min" and "h" stand for "minutes" and "hours" respectively. DTT means dithiothreitol. DMSO means dimethyl sulfoxide. EDTA means ethylenediamine tetraacetic acid.

Some of the preparation methods useful for the preparation of the compounds described herein may require protection of remote functionality (e.g., primary amine, secondary amine, carboxyl in Formula I precursors). The need for such protection will vary depending on the nature of the remote functionality and the conditions of the preparation methods. The need for such protection is readily determined by one skilled in the art. The use of such protection/deprotection methods is also within the skill in the art. For a general description of protecting groups and their use, see T.W. Greene, Protective Groups in Organic Synthesis, John Wiley & Sons, New York, 1991.

-131-

Biological AssaysDetermination of cAMP Elevation in 293-S Cell Lines Stably Overexpressing  
Recombinant Human EP<sub>2</sub> and EP<sub>4</sub> Receptors.

cDNAs representing the complete open reading frames of the human EP<sub>2</sub> and EP<sub>4</sub> receptors are generated by reverse transcriptase polymerase chain reaction using oligonucleotide primers based on published sequences (1, 2) and RNA from primary human kidney cells (EP<sub>2</sub>) or primary human lung cells (EP<sub>4</sub>) as templates. cDNAs are cloned into the multiple cloning site of pcDNA3 (Invitrogen Corporation, 3985B Sorrento Valley Blvd., San Diego, CA 92121) and used to transfect 293-S human embryonic kidney cells via calcium phosphate co-precipitation. G418-resistant colonies are expanded and tested for specific [<sup>3</sup>H]PGE<sub>2</sub> binding. Transfectants demonstrating high levels of specific [<sup>3</sup>H]PGE<sub>2</sub> binding are further characterized by Scatchard analysis to determine B<sub>max</sub> and K<sub>d</sub>s for PGE<sub>2</sub>. The lines selected for compound screening have approximately 338,400 receptors per cell and a K<sub>d</sub> = 12 nM for PGE<sub>2</sub> (EP<sub>2</sub>), and approximately 256,400 receptors per cell and a K<sub>d</sub> = 2.9 nM for PGE<sub>2</sub> (EP<sub>4</sub>). Constitutive expression of both receptors in parental 293-S cells is negligible. Cells are maintained in RPMI supplemented with fetal bovine serum (10% final) and G418 (700 ug/ml final).

cAMP responses in the 293-S/EP<sub>2</sub> and 293-S/EP<sub>4</sub> lines are determined by detaching cells from culture flasks in 1 ml of Ca<sup>++</sup> and Mg<sup>++</sup> deficient PBS via vigorous pounding, adding serum-free RPMI to a final concentration of 1 X 10<sup>6</sup> cells/ml, and adding 3-isobutyl-1-methylxanthine (IBMX) to a final concentration of 1 mM. One milliliter of cell suspension is immediately aliquoted into individual 2 ml screwcap microcentrifuge and incubated for 10 minutes, uncovered, at 37 °C, 5% CO<sub>2</sub>, 95% relative humidity. The compound to be tested is then added to cells at 1:100 dilutions such that final DMSO or ethanol concentrations is 1%. Immediately after adding compound, the tubes are covered, mixed by inverting two times, and incubated at 37 °C for 12 minutes. Samples are then lysed by incubation at 100 °C for 10 minutes and immediately cooled on ice for 5 minutes. Cellular debris is pelleted by centrifugation at 1000 X g for 5 minutes, and cleared lysates are transferred to fresh tubes. cAMP concentrations are determined using a commercially available cAMP radioimmunoassay kit RIA (NEK-033, DuPont/NEN Research Products, 549 Albany St., Boston, MA 02118) after diluting cleared lysates 1:10 in cAMP RIA assay buffer

-132-

(included in kit). Typically, one treats cells with 6-8 concentrations of the compound to be tested in 1 log increments. EC50 calculations are performed on a calculator using linear regression analysis on the linear portion of the dose response curves.

#### References

- 5           1. Regan, J.W. Bailey, T.J. Pepperl, D.J. Pierce, K.L. Bogardus, A.M. Donello, J.E. Fairbairn, C.E. Kedzie, K.M. Woodward, D.F. and Gil, D.W. 1994 Cloning of a Novel Human Prostaglandin Receptor with Characteristics of the Pharmacologically Defined EP<sub>2</sub> Subtype. Mol. Pharmacology 46:213-220.
2. Bastien, L., Sawyer, N., Grygorczyk, R., Metters, K., and Adam, M. 1994  
10 Cloning, Functional Expression, and Characterization of the Human Prostaglandin E<sub>2</sub> Receptor EP<sub>2</sub> Subtype. J. Biol. Chem. Vol 269, 16:11873-11877.

#### Assay for Binding to Prostaglandin E<sub>2</sub> Receptors

##### Membrane Preparation:

- 15           All operations are performed at 4 °C. Transfected cells expressing prostaglandin E<sub>2</sub> type 1 receptors (EP<sub>1</sub>), type 2 (EP<sub>2</sub>), type 3 (EP<sub>3</sub>) or type 4 (EP<sub>4</sub>) receptors are harvested and suspended to 2 million cells per ml in Buffer A [50 mM Tris-HCl (pH 7.4), 10 mM MgCl<sub>2</sub>, 1 mM EDTA, 1 mM Pefabloc peptide, (Boehringer Mannheim Corp., Indianapolis, IN), 10 uM Phosphoramidon peptide, (Sigma, St. Louis,  
20 MO), 1 uM pepstatin A peptide, (Sigma, St. Louis, MO), 10 uM elastatinal peptide, (Sigma, St. Louis, MO), 100 uM antipain peptide, (Sigma, St. Louis, MO)]. The cells are lysed by sonification with a Branson Sonifier (Model #250, Branson Ultrasonics Corporation, Danbury, CT) in 2 fifteen second bursts. Unlysed cells and debris are removed by centrifugation at 100 x g for 10 min. Membranes are then harvested by  
25 centrifugation at 45,000 x g for 30 minutes. Pelleted membranes are resuspended to 3-10 mg protein per ml, protein concentration being determined of the method of Bradford [Bradford, M., Anal. Biochem., 72, 248 (1976)]. Resuspended membranes are then stored frozen at -80 °C until use.

##### Binding Assay:

- 30           Frozen membranes prepared as above are thawed and diluted to 1 mg protein per ml in Buffer A above. One volume of membrane preparation is combined with 0.05 volume test compound or buffer and one volume of 3 nM <sup>3</sup>H-prostaglandin E<sub>2</sub> ( #TRK 431, Amersham, Arlington Heights, IL) in Buffer A. The mixture (205 µL total volume)

-133-

is incubated for 1 hour at 25°C. The membranes are then recovered by filtration through type GF/C glass fiber filters ( #1205-401, Wallac, Gaithersburg, MD ) using a Tomtec harvester ( Model Mach II/96, Tomtec, Orange, CT). The membranes with bound  $^3\text{H}$ -prostaglandin  $\text{E}_2$  are trapped by the filter, while the buffer and unbound  $^3\text{H}$ -  
5 prostaglandin  $\text{E}_2$  pass through the filter into waste. Each sample is then washed 3 times with 3 ml of [50 mM Tris-HCl (pH 7.4), 10 mM  $\text{MgCl}_2$ , 1 mM EDTA]. The filters are then dried by heating in a microwave oven. To determine the amount of  $^3\text{H}$ -prostaglandin bound to the membranes, the dried filters are placed into plastic bags with scintillation fluid and counted in a LKB 1205 Betaplate reader (Wallac,  
10 Gaithersburg, MD).  $\text{IC}_{50}$ s are determined from the concentration of test compound required to displace 50% of the specifically bound  $^3\text{H}$ -prostaglandin  $\text{E}_2$ .

The full length  $\text{EP}_1$  receptor is made as disclosed in Funk et al., Journal of Biological Chemistry, **1993**, **268**, 26767-26772. The full length  $\text{EP}_2$  receptor is made as disclosed in Regan et al., Molecular Pharmacology, **1994**, **46**, 213-220. The full  
15 length  $\text{EP}_3$  receptor is made as disclosed in Regan et al., British Journal of Pharmacology, **1994**, **112**, 377-385. The full length  $\text{EP}_4$  receptor is made as disclosed in Bastien, Journal of Biological Chemistry, **1994**, **269**, 11873-11877. These full length receptors are used to prepare 293S cells expressing the  $\text{EP}_1$ ,  $\text{EP}_2$ ,  $\text{EP}_3$  and  $\text{EP}_4$  receptors.

20 293S cells expressing either the human  $\text{EP}_1$ ,  $\text{EP}_2$ ,  $\text{EP}_3$  or  $\text{EP}_4$  prostaglandin  $\text{E}_2$  receptors are generated according to methods known to those skilled in the art. Typically, PCR (polymerase chain reaction) primers corresponding to the 5' and 3' ends of the published full length receptor are made according to the well known methods disclosed above and are used in an RT-PCR reaction using the total RNA  
25 from human kidney (for  $\text{EP}_1$ ), human lung (for  $\text{EP}_2$ ), human lung (for  $\text{EP}_3$ ) or human lymphocytes (for  $\text{EP}_4$ ) as a source. PCR products are cloned by the TA overhang method into pCR2.1 (Invitrogen, Carlsbad, CA) and identity of the cloned receptor is confirmed by DNA sequencing.

293S cells (Mayo, Dept. of Biochemistry, Northwestern Univ.) are transfected  
30 with the cloned receptor in pcDNA3 by electroporation. Stable cell lines expressing the receptor are established following selection of transfected cells with G418.

Clonal cell lines expressing the maximal number of receptors are chosen following a whole cell  $^3\text{H}$ -PGE $_2$  binding assay using unlabeled PGE $_2$  as a competitor.

-134-

Example Number	EP <sub>1</sub> (nM)		Example Number	EP <sub>2</sub> (nM)
1	395		1al	17
1a	2200		1an	94
1b	160		2	45
1c	24		2a	37
1d	885		2b	385
1e	760		3	650
1f	52		3a	440
1g	109		3b	7.83
1h	745		4	225
1i	72		4a	795
1j	385		4b	440
1k	1755		4c	885
1l	600		4d	175
1m	245		4e	410
1n	140		4f	705
1o	550		4g	720
1p	44		4h	210
1q	39		5	980
1r	308		5a	355
1s	165		5b	640
1t	8		6	315
1u	7		6a	325
1v	9		6c	300
1w	24		6d	170
1x	160		6e	325
1y	13		6f	490
1z	19		6g	350
1aa	335		6h	140
1ab	110		6i	270
1ac	335		6j	375
1ad	195		7a	400
1ae	121		7b	140
1af	106.5		7c	260
1ag	169		7d	52
1ah	180.5		7f	344
1ai	125		7g	412
1aj	25.5		7h	172
1ak	6.19		8	71

-135-

Example Number	EP <sub>2</sub> ( nM)		Example Number	EP <sub>2</sub> (nM)
8a	180		12i	16
8b	266		12k	31
9	17		12m	38
9a	84.5		12n	10
9b	94.5		12o	10
9c	490		12p	29
9d	202.5		12q	10
11a	8.2		12r	10
11b	12.75		12s	10
11c	13.2		12t	57
11d	45		12u	63
11e	6.3		12v	10
11f	25.68		12w	47
11g	25.04		12x	10
11h	7.5		12y	55
11i	89		12z	200
11m	10		13a	10
11n	10		13b	22
11o	10		13c	10
11p	10		13d	10
11q	10		13e	10
11r	10		13f	10
11s	10		13g	340
11t	10		13h	10
11u	10.5		13i	10.4
11v	10		13j	26
11w	36.5		13k	99
11x	63.5		13l	47.5
11y	10		13m	155
11z	13		13o	10
12a	65		13r	10
12b	14.3		13s	10
12c	25		13t	230
12d	515		13u	7.99
12e	118.5		13v	25.83
12f	10		13w	12
12g	10		13x	15.5
12h	10		13y	10



-136-


Example Number	EP <sub>2</sub> (nM)
13z	24.5
14a	13.5
14b	10
14c	68
14d	10
14e	13
15a	10
15b	10
15c	35
15d	10
15e	39
15f	10
15g	10
16a	110
16b	215

-137-

### Claims

What is claimed is:

1. A method of treating pulmonary hypertension, facilitating joint fusion,  
 5 facilitating tendon and ligament repair, reducing the occurrence of secondary fracture, treating avascular necrosis, facilitating cartilage repair, facilitating bone healing after limb transplantation, facilitating liver regeneration, facilitating wound healing, reducing the occurrence of gastric ulceration, treating hypertension,  
 10 facilitating the growth of tooth enamel or finger or toe nails, treating glaucoma, treating ocular hypertension or repairing damage caused by metastatic bone disease, the method comprising administering to a patient in need thereof a therapeutically effective amount of an EP<sub>2</sub> selective receptor agonist.
2. A method of treating pulmonary hypertension, facilitating joint fusion,  
 15 facilitating tendon and ligament repair, reducing the occurrence of secondary fracture, treating avascular necrosis, facilitating cartilage repair, facilitating bone healing after limb transplantation, facilitating liver regeneration, facilitating wound healing, reducing the occurrence of gastric ulceration, treating hypertension,  
 20 facilitating the growth of tooth enamel or finger or toe nails, treating glaucoma, treating ocular hypertension or repairing damage caused by metastatic bone disease, the method comprising administering to a patient in need thereof a therapeutically effective amount of compound of Formula I



Formula I

or a prodrug thereof, or a pharmaceutically acceptable salt thereof, wherein

A is SO<sub>2</sub> or CO;

G is Ar, Ar<sup>1</sup>-V-Ar<sup>2</sup>, Ar-(C<sub>1</sub>-C<sub>6</sub>)alkylene, Ar-CONH-(C<sub>1</sub>-C<sub>6</sub>)alkylene, R<sup>1</sup>R<sup>2</sup>-amino,  
 oxy(C<sub>1</sub>-C<sub>6</sub>)alkylene, amino substituted with Ar, or amino substituted with Ar(C<sub>1</sub>-  
 30 C<sub>4</sub>)alkylene and R<sup>11</sup>, wherein R<sup>11</sup> is H or (C<sub>1</sub>-C<sub>8</sub>)alkyl, R<sup>1</sup> and R<sup>2</sup> may be taken separately and are independently selected from H and (C<sub>1</sub>-C<sub>8</sub>)alkyl, or R<sup>1</sup> and R<sup>2</sup>

-138-

are taken together with the nitrogen atom of the amino group to form a five- or six-membered azacycloalkyl, said azacycloalkyl optionally containing an oxygen atom and optionally mono-, di- or tri-substituted independently with up to two oxo, hydroxy, (C<sub>1</sub>-C<sub>4</sub>)alkyl, fluoro or chloro;

5 B is N or CH;

Q is

-(C<sub>2</sub>-C<sub>6</sub>)alkylene-W-(C<sub>1</sub>-C<sub>3</sub>)alkylene-, said alkylenes each optionally substituted with up to four substituents independently selected from fluoro or (C<sub>1</sub>-C<sub>4</sub>)alkyl,

10 -(C<sub>4</sub>-C<sub>8</sub>)alkylene-, said alkylene optionally substituted with up to four substituents independently selected from fluoro or (C<sub>1</sub>-C<sub>4</sub>)alkyl,

-X-(C<sub>1</sub>-C<sub>5</sub>)alkylene-, said alkylene optionally substituted with up to four substituents independently selected from fluoro or (C<sub>1</sub>-C<sub>4</sub>)alkyl,

15 -(C<sub>1</sub>-C<sub>5</sub>)alkylene-X-, said alkylene optionally substituted with up to four substituents independently selected from fluoro or (C<sub>1</sub>-C<sub>4</sub>)alkyl,

-(C<sub>1</sub>-C<sub>3</sub>)alkylene-X-(C<sub>1</sub>-C<sub>3</sub>)alkylene-, said alkylenes each optionally substituted with up to four substituents independently selected from fluoro or (C<sub>1</sub>-C<sub>4</sub>)alkyl,

-(C<sub>2</sub>-C<sub>4</sub>)alkylene-W-X-(C<sub>0</sub>-C<sub>3</sub>)alkylene-, said alkylenes each optionally substituted with up to four substituents each independently selected from fluoro or (C<sub>1</sub>-C<sub>4</sub>)alkyl,

20 -(C<sub>0</sub>-C<sub>4</sub>)alkylene-X-W-(C<sub>1</sub>-C<sub>3</sub>)alkylene-, said alkylenes each optionally substituted with up to four substituents each independently selected from fluoro or (C<sub>1</sub>-C<sub>4</sub>)alkyl,

25 -(C<sub>2</sub>-C<sub>5</sub>)alkylene-W-X-W-(C<sub>1</sub>-C<sub>3</sub>)alkylene-, wherein the two occurrences of W are independent of each other, said alkylenes each optionally substituted with up to four substituents each independently selected from fluoro or (C<sub>1</sub>-C<sub>4</sub>)alkyl,

-(C<sub>1</sub>-C<sub>4</sub>)alkylene-ethenylene-(C<sub>1</sub>-C<sub>4</sub>)alkylene-, said alkylenes and said ethenylene each optionally substituted with up to four substituents each independently selected from fluoro or (C<sub>1</sub>-C<sub>4</sub>)alkyl,

30 -(C<sub>1</sub>-C<sub>4</sub>)alkylene-ethenylene-(C<sub>0</sub>-C<sub>2</sub>)alkylene-X-(C<sub>0</sub>-C<sub>5</sub>)alkylene-, said alkylenes and said ethenylene each optionally substituted with up to four substituents each independently selected from fluoro or (C<sub>1</sub>-C<sub>4</sub>)alkyl,

-139-

-(C<sub>1</sub>-C<sub>4</sub>)alkylene-ethenylene-(C<sub>0</sub>-C<sub>2</sub>)alkylene-X-W-(C<sub>1</sub>-C<sub>3</sub>)alkylene-, said alkylenes and said ethenylene optionally each substituted with up to four substituents each independently selected from fluoro or (C<sub>1</sub>-C<sub>4</sub>)alkyl,

5       -(C<sub>1</sub>-C<sub>4</sub>)alkylene-ethynylene-(C<sub>1</sub>-C<sub>4</sub>)alkylene-, said alkylenes and said ethynylene each optionally substituted with up to four substituents each independently selected from fluoro or (C<sub>1</sub>-C<sub>4</sub>)alkyl, or

      -(C<sub>1</sub>-C<sub>4</sub>)alkylene-ethynylene-X-(C<sub>0</sub>-C<sub>3</sub>)alkylene-, said alkylenes and said ethynylene each optionally substituted with up to four substituents each independently selected from fluoro or (C<sub>1</sub>-C<sub>4</sub>)alkyl;

10       Z is carboxyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonyl, tetrazolyl, 1,2,4-oxadiazolyl, 5-oxo-1,2,4-oxadiazolyl, 5-oxo-1,2,4-thiadiazolyl, (C<sub>1</sub>-C<sub>4</sub>)alkylsulfonylcarbamoyl or phenylsulfonylcarbamoyl;

      K is a bond, (C<sub>1</sub>-C<sub>9</sub>)alkylene, thio(C<sub>1</sub>-C<sub>4</sub>)alkylene, (C<sub>1</sub>-C<sub>4</sub>)alkylenethio(C<sub>1</sub>-C<sub>4</sub>)alkylene, (C<sub>1</sub>-C<sub>4</sub>)alkyleneoxy(C<sub>1</sub>-C<sub>4</sub>)alkylene or oxy(C<sub>1</sub>-C<sub>4</sub>)alkylene, said (C<sub>1</sub>-C<sub>9</sub>)alkylene optionally mono-unsaturated and wherein, when K is not a bond, K is optionally mono-, di- or tri-substituted independently with chloro, fluoro, hydroxy or methyl;

      M is -Ar<sup>3</sup>, -Ar<sup>4</sup>-V<sup>1</sup>-Ar<sup>5</sup>, -Ar<sup>4</sup>-S-Ar<sup>5</sup>, -Ar<sup>4</sup>-SO-Ar<sup>5</sup>, -Ar<sup>4</sup>-SO<sub>2</sub>-Ar<sup>5</sup> or -Ar<sup>4</sup>-O-Ar<sup>5</sup>;

      Ar is a partially saturated or fully unsaturated five to eight membered ring optionally having one to four heteroatoms selected independently from oxygen, sulfur and nitrogen, or a bicyclic ring consisting of two fused independently partially saturated, fully saturated or fully unsaturated five or six membered rings, taken independently, optionally having one to four heteroatoms selected independently from nitrogen, sulfur and oxygen, or a tricyclic ring consisting of three fused independently partially saturated, fully saturated or fully unsaturated five or six membered rings, optionally having one to four heteroatoms selected independently from nitrogen, sulfur and oxygen, said partially or fully saturated ring, bicyclic ring or tricyclic ring optionally having one or two oxo groups substituted on carbon or one or two oxo groups substituted on sulfur; or Ar is a fully saturated five to seven membered ring having one or two heteroatoms selected independently from oxygen, sulfur and nitrogen;

      Ar<sup>1</sup> and Ar<sup>2</sup> are each independently a partially saturated, fully saturated or fully unsaturated five to eight membered ring optionally having one to four

-140-

heteroatoms selected independently from oxygen, sulfur and nitrogen, or a bicyclic ring consisting of two fused independently partially saturated, fully saturated or fully unsaturated five or six membered rings, taken independently, optionally having one to four heteroatoms selected independently from nitrogen, sulfur and oxygen, or a tricyclic ring consisting of three fused independently partially saturated, fully saturated or fully unsaturated five or six membered rings, optionally having one to four heteroatoms selected independently from nitrogen, sulfur and oxygen, said partially or fully saturated ring, bicyclic ring or tricyclic ring optionally having one or two oxo groups substituted on carbon or one or two oxo groups substituted on sulfur;

said Ar, Ar<sup>1</sup> and Ar<sup>2</sup> moieties are optionally substituted on carbon or nitrogen, on one ring if the moiety is monocyclic, on one or both rings if the moiety is bicyclic, or on one, two or three rings if the moiety is tricyclic, with up to three substituents per moiety independently selected from R<sup>3</sup>, R<sup>4</sup> and R<sup>5</sup> wherein R<sup>3</sup>, R<sup>4</sup> and R<sup>5</sup> are independently hydroxy, nitro, halo, carboxy, (C<sub>1</sub>-C<sub>7</sub>)alkoxy, (C<sub>1</sub>-C<sub>4</sub>)alkoxy(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxycarbonyl, (C<sub>1</sub>-C<sub>7</sub>)alkyl, (C<sub>2</sub>-C<sub>7</sub>)alkenyl, (C<sub>2</sub>-C<sub>7</sub>)alkynyl, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl(C<sub>1</sub>-C<sub>4</sub>)alkanoyl, formyl, (C<sub>1</sub>-C<sub>8</sub>)alkanoyl, (C<sub>1</sub>-C<sub>6</sub>)alkanoyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkanoylamino, (C<sub>1</sub>-C<sub>4</sub>)alkoxycarbonylamino, hydroxysulfonyl, aminocarbonylamino or mono-N-, di-N,N-, di-N,N'- or tri-N,N,N'-(C<sub>1</sub>-C<sub>4</sub>)alkyl substituted aminocarbonylamino, sulfonamido, (C<sub>1</sub>-C<sub>4</sub>)alkylsulfonamido, amino, mono-N- or di-N,N-(C<sub>1</sub>-C<sub>4</sub>)alkylamino, carbamoyl, mono-N- or di-N,N-(C<sub>1</sub>-C<sub>4</sub>)alkylcarbamoyl, cyano, thiol, (C<sub>1</sub>-C<sub>6</sub>)alkylthio, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfinyl, (C<sub>1</sub>-C<sub>4</sub>)alkylsulfonyl or mono-N- or di-N,N-(C<sub>1</sub>-C<sub>4</sub>)alkylaminosulfinyl;

Ar<sup>3</sup>, Ar<sup>4</sup> and Ar<sup>5</sup> are each independently a partially saturated, fully saturated or fully unsaturated five to eight membered ring optionally having one to four heteroatoms selected independently from oxygen, sulfur and nitrogen, or a bicyclic ring consisting of two fused independently partially saturated, fully saturated or fully unsaturated five or six membered rings, taken independently, optionally having one to four heteroatoms selected independently from nitrogen, sulfur and oxygen, or a tricyclic ring consisting of three fused independently partially saturated, fully saturated or fully unsaturated five or six membered rings, optionally having one to four heteroatoms selected independently from nitrogen,

-141-

sulfur and oxygen, said partially or fully saturated ring, bicyclic ring or tricyclic ring optionally having one or two oxo groups substituted on carbon or one or two oxo groups substituted on sulfur; said Ar<sup>3</sup>, Ar<sup>4</sup> and Ar<sup>5</sup> moieties are optionally substituted on carbon or nitrogen, on one ring if the moiety is monocyclic, on one or both rings if the moiety is bicyclic, or on one, two or three rings if the moiety is tricyclic, with up to three substituents per moiety independently selected from R<sup>31</sup>, R<sup>41</sup> and R<sup>51</sup> wherein R<sup>31</sup>, R<sup>41</sup> and R<sup>51</sup> are independently hydroxy, nitro, halo, carboxy, (C<sub>1</sub>-C<sub>7</sub>)alkoxy, (C<sub>1</sub>-C<sub>4</sub>)alkoxy(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxycarbonyl, (C<sub>1</sub>-C<sub>7</sub>)alkyl, (C<sub>2</sub>-C<sub>7</sub>)alkenyl, (C<sub>2</sub>-C<sub>7</sub>)alkynyl, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl(C<sub>1</sub>-C<sub>4</sub>)alkanoyl, formyl, (C<sub>1</sub>-C<sub>8</sub>)alkanoyl, (C<sub>1</sub>-C<sub>6</sub>)alkanoyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkanoylamino, (C<sub>1</sub>-C<sub>4</sub>)alkoxycarbonylamino, hydroxysulfonyl, aminocarbonylamino or mono-N-, di-N,N-, di-N,N'- or tri-N,N,N'-(C<sub>1</sub>-C<sub>4</sub>)alkyl substituted aminocarbonylamino, sulfonamido, (C<sub>1</sub>-C<sub>4</sub>)alkylsulfonamido, amino, mono-N- or di-N,N-(C<sub>1</sub>-C<sub>4</sub>)alkylamino, carbamoyl, mono-N- or di-N,N-(C<sub>1</sub>-C<sub>4</sub>)alkylcarbamoyl, cyano, thiol, (C<sub>1</sub>-C<sub>6</sub>)alkylthio, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfinyl, (C<sub>1</sub>-C<sub>4</sub>)alkylsulfonyl or mono-N- or di-N,N-(C<sub>1</sub>-C<sub>4</sub>)alkylaminosulfinyl;

W is oxy, thio, sulfino, sulfonyl, aminosulfonyl-, -mono-N-(C<sub>1</sub>-C<sub>4</sub>)alkyleneaminosulfonyl-, sulfonylamino, N-(C<sub>1</sub>-C<sub>4</sub>)alkylenesulfonylamino, carboxamido, N-(C<sub>1</sub>-C<sub>4</sub>)alkylenecarboxamido, carboxamidooxy, N-(C<sub>1</sub>-C<sub>4</sub>)alkylenecarboxamidooxy, carbamoyl, -mono-N-(C<sub>1</sub>-C<sub>4</sub>)alkylenecarbamoyl, carbamoyloxy, or -mono-N-(C<sub>1</sub>-C<sub>4</sub>)alkylenecarbamoyloxy, wherein said W alkyl groups are optionally substituted on carbon with one to three fluorines;

X is a five or six membered aromatic ring optionally having one or two heteroatoms selected independently from oxygen, nitrogen, and sulfur; said ring optionally mono-, di- or tri-substituted independently with halo, (C<sub>1</sub>-C<sub>3</sub>)alkyl, trifluoromethyl, trifluoromethyloxy, difluoromethyloxy, hydroxyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy, or carbamoyl;

R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, R<sup>4</sup>, R<sup>5</sup>, R<sup>11</sup>, R<sup>31</sup>, R<sup>41</sup> and R<sup>51</sup>, when containing an alkyl, alkylene, alkenylene or alkynylene moiety, are optionally mono-, di- or tri-substituted on carbon independently with halo or hydroxy; and

-142-

V and V<sup>1</sup> are each independently a bond, thio(C<sub>1</sub>-C<sub>4</sub>)alkylene, (C<sub>1</sub>-C<sub>4</sub>)alkylenethio, (C<sub>1</sub>-C<sub>4</sub>)alkyleneoxy, oxy(C<sub>1</sub>-C<sub>4</sub>)alkylene or (C<sub>1</sub>-C<sub>3</sub>)alkylene optionally mono- or di-substituted independently with hydroxy or fluoro;

with the provisos that:

- 5           a. when K is (C<sub>2</sub>-C<sub>4</sub>)alkylene and M is Ar<sup>3</sup> and Ar<sup>3</sup> is cyclopent-1-yl, cyclohex-1-yl, cyclohept-1-yl or cyclooct-1-yl then said (C<sub>5</sub>-C<sub>8</sub>)cycloalkyl substituents are not substituted at the one position with hydroxy; and
- b. when K is a bond; G is phenyl, phenylmethyl, substituted phenyl or substituted phenylmethyl; Q is (C<sub>3</sub>-C<sub>8</sub>)alkylene; and M is Ar<sup>3</sup> or Ar<sup>4</sup>-Ar<sup>5</sup>, then A is  
10       sulfonyl.

3. A method of treating pulmonary hypertension, facilitating joint fusion, facilitating tendon and ligament repair, reducing the occurrence of secondary fracture, treating avascular necrosis, facilitating cartilage repair, facilitating bone  
15       healing after limb transplantation, facilitating liver regeneration, facilitating wound healing, reducing the occurrence of gastric ulceration, treating hypertension, facilitating the growth of tooth enamel or finger or toe nails, treating glaucoma, treating ocular hypertension or repairing damage caused by metastatic bone disease, the method comprising administering to a patient in need thereof a  
20       therapeutically effective amount of (3-(((4-tert-butyl-benzyl)-(pyridine-3-sulfonyl)-amino)-methyl)-phenoxy)-acetic acid or a pharmaceutically acceptable salt thereof.

4. The method of any of claims 1-3 wherein the method is treating hypertension  
25       or treating pulmonary hypertension.

5. The method of any of claims 1-3 wherein the method is facilitating joint fusion.

6. The method of any of claims 1-3 wherein the method is facilitating tendon and  
30       ligament repair or facilitating cartilage repair.

7. The method of any of claims 1-3 wherein the method is reducing the occurrence of secondary fracture.

-143-

8. The method of any of claims 1-3 wherein the method is treating avascular necrosis.
- 5 9. The method of any of claims 1-3 wherein the method is facilitating bone healing after limb transplantation.
- 10 10. The method of any of claims 1-3 wherein the method is facilitating liver regeneration.
- 11 11. The method of any of claims 1-3 wherein the method is facilitating wound healing.
- 15 12. The method of any of claims 1-3 wherein the method is reducing the occurrence of gastric ulceration.
- 20 13. The method of any of claims 1-3 wherein the method is facilitating the growth of tooth enamel, fingernails or toenails.
- 20 14. The method of any of claims 1-3 wherein the method is treating glaucoma or treating ocular hypertension.
- 25 15. A method of any of claims 1-3 wherein the method is repairing damage caused by metastatic bone disease.



# INTERNATIONAL SEARCH REPORT

Int. Patent Application No.  
PCT/IB2004/000553

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A61K31/185 A61K31/415 A61K31/433 A61P19/00 A61P9/02  
A61P1/04 A61P27/06

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, EMBASE, CHEM ABS Data

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 99/19300 A (ROSATI ROBERT LOUIS ;PFIZER (US); CAMERON KIMBERLY O KEEFE (US); L) 22 April 1999 (1999-04-22)	1-3, 6, 7, 9
Y	claims 1, 36, 65, 69, 71	1-4, 12, 14
X	WO 98/27976 A (ROSATI ROBERT LOUIS ;KE HUA ZHU (US); PFIZER (US); CAMERON KIMBERL) 2 July 1998 (1998-07-02)	1, 2, 9
X	US 6 046 236 A (HAMANAKA NOBUYUKI ET AL) 4 April 2000 (2000-04-04)	1
Y	claim 1 column 7, line 26	1-4, 12
	-/--	



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

\* Special categories of cited documents:

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- \*Z\* document member of the same patent family

Date of the actual completion of the international search

24 May 2004

Date of mailing of the international search report

02/06/2004

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Beranová, P

# INTERNATIONAL SEARCH REPORT

In International Application No  
PCT/IB2004/000553

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 6 376 533 B1 (BURK ROBERT M ET AL)	1
Y	23 April 2002 (2002-04-23) claim 1 column 1, line 9 - line 12 -----	1-3, 14

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/IB2004/000553

## Box II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:  
  
Although claims 1 - 15 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the composition.
2. ☐ Claims Nos.:  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No  
PCT/IB2004/000553

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9919300	A	22-04-1999	AU 731509 B2	29-03-2001
			AU 9181598 A	03-05-1999
			BG 104315 A	29-12-2000
			BR 9813028 A	15-08-2000
			CA 2305548 A1	22-04-1999
			CN 1275123 T	29-11-2000
			EA 3396 B1	24-04-2003
			EP 1021410 A1	26-07-2000
			HR 20000201 A1	30-06-2000
			HU 0100101 A2	28-06-2001
			WO 9919300 A1	22-04-1999
			ID 24836 A	24-08-2000
			JP 2001519414 T	23-10-2001
			NO 20001754 A	07-06-2000
			NZ 503165 A	27-07-2001
			PL 340014 A1	15-01-2001
			SK 4752000 A3	10-05-2001
			TR 200000927 T2	21-11-2000
			US 2003078261 A1	24-04-2003
			US 6498172 B1	24-12-2002
			ZA 9809230 A	10-04-2000
WO 9827976	A	02-07-1998	AU 4881697 A	17-07-1998
			CA 2275595 A1	02-07-1998
			EP 0951282 A1	27-10-1999
			HU 0001836 A2	28-09-2000
			WO 9827976 A1	02-07-1998
			JP 2000507961 T	27-06-2000
			KR 2000057680 A	25-09-2000
			US 6426359 B1	30-07-2002
			ZA 9711435 A	21-06-1999
US 6046236	A	04-04-2000	AT 145897 T	15-12-1996
			CA 2113787 A1	30-07-1994
			DE 69401011 D1	16-01-1997
			DE 69401011 T2	07-05-1997
			DK 608847 T3	23-12-1996
			EP 0608847 A1	03-08-1994
			ES 2097557 T3	01-04-1997
			GR 3022470 T3	31-05-1997
			JP 6279395 A	04-10-1994
			US 5663417 A	02-09-1997
US 6376533	B1	23-04-2002	US 2002187961 A1	12-12-2002